

Journal of The Chemical Society, Chemical Communications

NUMBER 24/1980

Syntheses and Ring-opening Reactions of 5-Alkoxy- and 5-Acetoxy-3-oxo-2-azabicyclo[2.2.0]hex-5-enes


By CHIKARA KANEKO,* KAZUHIRO SHIBA, HARUE FUJII, and YŪ MOMOSE

(Faculty of Pharmaceutical Sciences, Kanazawa University, Takara-machi, Kanazawa, 920, Japan)

Summary The efficient photochemical synthesis of 5-alkoxy- and 5-acetoxy-3-oxo-2-azabicyclo[2.2.0]hex-5-enes (**2a—e** and **f,g**), and a new rearrangement reaction of the former compounds (**2a—e**) to 6-alkoxy-2-pyridones (**3a—e**), are reported.

COREY and STREITH first synthesized *N*-methyl-photo-2-pyridone *via* low temperature photoisomerization of a dilute solution of *N*-methyl-2-pyridone.¹ Although the relatively high stability of the photo-2-pyridones, including the parent compound 3-oxo-2-azabicyclo[2.2.0]hex-5-ene, was later demonstrated by two research groups,^{2,3} the syntheses of these photo-2-pyridones have been hampered owing to concomitant and more efficient formation of the photodimers.^{1–5}

De Selms *et al.*² reported a high-yield formation of the corresponding photo-2-pyridone by irradiation of recinine (3-cyano-4-methoxy-1-methyl-2-pyridone) with no dimer detected, and speculated that this exceptional photochemical behaviour of recinine might have some significance in phytochemical photosynthesis. Previously, we reported that irradiation of 4-methoxy-2-pyridone in acetone in the presence of olefins gave the intermolecular [2 + 2] cycloadducts with no dimer formation.⁶ These two experiments seem to suggest an inability of the 4-alkoxy-2-pyridones to photodimerize and, hence, the selective and high-yield formation of the photo-2-pyridones by irradiation of compounds (**1**) in an appropriate transparent solvent. Experiments along these lines have led us to find an efficient route to the 5-alkoxy- and 5-acetoxy-derivatives (**2a—e** and **2f,g**) of 3-oxo-2-azabicyclo[2.2.0]hex-5-ene from the 4-oxygenated



2-pyridones (**1a—g**), and a novel rearrangement of the 5-alkoxy derivatives (**2a—e**) to the corresponding 6-alkoxy-2-pyridones (**3a—e**).

TABLE. M.p.s and yields of the photo-2-pyridones (**2**).

	R ¹	R ²	R ³	R ⁴	M.p./ °C	Yield/ %
a ;	Me	H	Me	H	Oil	81.9
b ;	H	H	Me	H	94—95	83.4
c ;	H	H	CH ₂ CH=CH ₂	H	42—43	80.0
d ;	H	H	[CH ₂] ₂ CH=CH ₂	H	38—41	92.0
e ;	H	H	CH ₂ Ph	H	122—124	89.8
f ;	H	CH ₂ CH=CH ₂	COMe	H	92—94	85.5
g ;	H	H	COMe	Me	Oil	88.5


Syntheses were carried out by irradiation of 10^{–2}–10^{–3} M ethereal solutions (other transparent solvents can also be used)† with Pyrex filtered irradiation from a high-pressure mercury arc lamp (Toshiba 400P) and the products (**2a—g**) were isolated by direct recrystallization or column chromatographic separation of the concentrated residue. No photodimer was obtained from any of these 4-oxygenated

† In these solvents, the cycloadducts of pyridones (**1**) with olefins were not obtained.

2-pyridones (**1a—g**), and hence the photo-2-pyridones (**2a—g**) were obtained as the sole product in very high yields.

These photo-2-pyridones (**2**) are all liquids or low melting solids and were characterized by acceptable mass (molecular ions and substituted cyclobutadiene cations as the major fragment ions), i.r. (absorptions at 1725—1765 cm⁻¹: β -lactam carbonyl), and n.m.r. spectra.[†]

Thermal reactions of the photo-2-pyridones so far reported have been limited to cycloreversion leading to the original 2-pyridones (*e.g.*, path a).^{2,3} However, it was found that, while the 5-acetoxy derivatives (**2f,g**) reverted selectively to 4-acetoxy-2-pyridones (**1f,g**) irrespective of the conditions employed, thermolysis of the 5-alkoxy derivatives (**2a—e**) above 80 °C caused concomitant formation of both 4- and 6-alkoxy-2-pyridones⁷ (**1** and **3**), the yield ratio of which was dependent upon the solvent employed. Thus, for example, while (**2e**) afforded 4-benzoyloxy-2-pyridone (**1e**) (24.0%) and its 6-isomer (**3e**) (20.8%) in addition to recovered (**2e**) (45.4%) after refluxing in benzene for 60 h, the same two pyridones were obtained in the respective yields of 20.2 and 57.9% by refluxing in acetonitrile (30 h), with 6.5% recovery of (**2e**). Preferential formation of the 6-alkoxy-2-pyridones (**3a—e**) compared to the 4-isomers (**1a—e**) was also observed in acetone (sealed tube at 100 °C), but not in *o*-dichlorobenzene (reflux). These facts[§] imply the intermediacy of zwitterionic species (**4a—e**) in the rearrangement of (**2a—e**) to (**3a—e**). In accordance with this explanation,[¶] the



addition of AgBF₄⁸ to these reaction media led to almost selective formation of the 4-alkoxy-2-pyridones (**2a—e**).

Since these 4-oxygenated 2-pyridones never photo-dimerize under any condition,** a general synthetic route to photo-2-pyridones having an oxygen function at the 5-position seems to be established.

(Received, 28th July 1980; Com. 811.)

[†] N.m.r. spectra were measured in CDCl₃: δ (**2b**) 3.18 (s, 1- and 4-H), 3.80 (s, Me), 5.00 (s, 6-H), and 6.37br (s, NH); δ (**2f**) 2.17 (s, Me), 4.13 (s, 1-H), 5.73 (s, 6-H), 6.80br (s, NH), and allyl group signals.

[§] The presence of the alkoxy-function at the 5-position of (**2a—e**) may facilitate the heterolytic cleavage of the C(1)–N(2) bond to give (**4**). The weaker electron-donating character of the acetoxy-function is probably not enough to accelerate this cleavage. Dielectric constants (ϵ) for these aprotic solvents (MeCN, acetone, *o*-dichlorobenzene, and benzene) are 37.5, 20.7, 9.93, and 2.28, respectively.

[¶] In order to verify the proposed mechanism, photochemical conversion of (**3b**) into (**5b**) was examined under the same conditions as for the conversion of (**1**) into (**2**). However, within a comparable irradiation period, the starting pyridone was recovered and neither the photopyridone nor the dimer was detected.

** The photolyses of (**1**) under sensitization conditions also did not produce any photodimer.⁶

¹ E. J. Corey and J. Streith, *J. Am. Chem. Soc.*, 1964, **86**, 950.

² R. C. De Selms and W. R. Schleigh, *Tetrahedron Lett.*, 1972, 3563.

³ H. Furrer, *Chem. Ber.*, 1972, **105**, 2780.

⁴ E. C. Taylor, R. O. Kan, and W. W. Paudler, *J. Am. Chem. Soc.*, 1961, **83**, 4484.

⁵ Y. Nakamura, T. Kato, and Y. Morita, *J. Chem. Soc., Chem. Commun.*, 1978, 620.

⁶ H. Fujii, K. Shiba, and C. Kaneko, *J. Chem. Soc., Chem. Commun.*, 1980, 537.

⁷ The structures of the 6-alkoxy-2-pyridones (**3a—e**) were assigned from their n.m.r. spectra *e.g.*, δ (**3e**) 5.73 (dd, 5-H), 6.16 (dd, 3-H), and 7.32 (dd, 4-H) ($J_{3,4}$ 8.6, $J_{4,5}$ 7.6, and $J_{3,5}$ 0.9 Hz), and benzyl signals at 5.07 (s) and 7.25br (s). The compound (**3b**) obtained from (**2b**) was identical with an authentic sample: A. R. Katritzky, F. D. Popp, and J. D. Rowe, *J. Chem. Soc. (B)*, 1966, 562.

⁸ Metal species such as AgBF₄ are known to promote orbital-symmetry forbidden disrotatory opening in molecules such as Dewar benzene: J. Wristers, L. Brener, and R. Pettit, *J. Am. Chem. Soc.*, 1970, **92**, 7499; W. Slegler, R. Case, J. S. McKennis, and R. Pettit, *ibid.*, 1974, **96**, 287.