

This article was downloaded by: [Fordham University]

On: 18 November 2012, At: 13:37

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office:

Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Synthetic Communications: An International Journal for Rapid Communication of Synthetic Organic Chemistry

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/lcyc20>

Selective Mono-BOC Protection of Diamines

Dae Woo Lee ^a, Hyun-Joon Ha ^a & Won Koo Lee ^b

^a Department of Chemistry, Hankuk University of Foreign Studies, Yongin, Kyunggi-Do, Korea

^b Department of Chemistry and Interdisciplinary Program of Integrated Biotechnology, Sogang University, Seoul, Korea

Version of record first published: 10 Mar 2007.

To cite this article: Dae Woo Lee, Hyun-Joon Ha & Won Koo Lee (2007): Selective Mono-BOC Protection of Diamines, *Synthetic Communications: An International Journal for Rapid Communication of Synthetic Organic Chemistry*, 37:5, 737-742

To link to this article: <http://dx.doi.org/10.1080/00397910601131403>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Selective Mono-BOC Protection of Diamines

Dae Woo Lee and Hyun-Joon Ha

Department of Chemistry, Hankuk University of Foreign Studies,
Yongin, Kyunggi-Do, Korea

Won Koo Lee

Department of Chemistry and Interdisciplinary Program of Integrated
Biotechnology, Sogang University, Seoul, Korea


Abstract: A facile route for mono-BOC protection of symmetrical and unsymmetrical diamines was developed by sequential additions of 1 mol of HCl and 1 mol of (BOC)₂O followed by neutralization.

Keywords: diamine, mono-BOC, protection

Monofunctionalization of symmetrical or unsymmetrical diamines is an essential step for the synthesis of biologically important pharmacores and materials.^[1–7] Recent progress on chemical and biological chip production requires monofunctionalized diamines to hold functional molecules at the one end while the other end is free to be attached on the solid surface.^[8–10] Therefore, a facile and large-scale synthetic method is urgent for the preparation of monoprotected diamine.^[11–22] Among the various amine protecting groups, BOC is the most popular, and its protected form is normally in a good physical state to be handled for the next reaction.^[19] Herein we report a facile route for mono-BOC protection of symmetrical and unsymmetrical diamines whose procedure is cost-efficient and applicable to a multigram scale.

Received in Japan

Address correspondence to Hyun-Joon Ha, Department of Chemistry, Hankuk University of Foreign Studies, Yongin, Kyunggi-Do 449719, Korea. E-mail: hjha@hufs.ac.kr




Scheme 1.

Two amine groups are differentiated as an acid salt and a free base, which is ready for further functionalization. One mole of hydrogen chloride was added to one molecule of a diamine bearing two basic amino groups in 50% aqueous methanol. The solution was stirred for 30 min to reach full equilibrium. Three different species would possibly exist including starting diamine **1** with two free amines, **A** as one free amine and one HCl salt, and **B** with two amine salts in aqueous methanol. Among them, **A** would be dominant in the solution after complete equilibrium (Scheme 1).

Addition of 1 mol of (BOC)₂O into this solution would lead to the reaction of the free amine in the dominant species **A** to yield mono-BOC-protected diamine. This procedure was proven to be highly efficient with good yields compared to the conventional method that requires tight control of the substrate concentration and very slow addition of the (BOC)₂O solution for a long time to prevent di-BOC formation.^[14,20–22] After the reaction was completed, MeOH was evaporated and diethyl ether was added to remove unreacted diamine. Then NaOH solution was added to let one HCl amine salt free. The reaction product was extracted by organic solvent. This procedure is applicable to various diamines **1** to give mono-BOC-protected diamines **2** without tedious chromatographic procedure in high yield (Scheme 2).








We have found that this was good for producing mono-BOC-protected diamines on a scale of several hundred grams, which is exemplified in Table 1. Acyclic symmetric diamines **1a–1d** yielded the expected products **2a–2d** in 65–87% yields. This worked for the cyclic diamine **1e** to provide the mono-BOC product **2e** in 80% yield. This procedure was also good for unsymmetric diamine substrates, **1f** and **1g**, to yield mon-BOC products, **2f** and **2g**, in 72 and 95% yields, respectively.

In conclusion, we developed a facile route for mono-BOC protection of symmetrical and unsymmetrical diamines by sequential additions of 1 mol of HCl and 1 mol of (BOC)₂O followed by neutralization.



Scheme 2.

Table 1. Mono-BOC protection of diamines

Diamines	Products	Yield (%) ^{a,b}
1a		87 ^[12,14]
1b		75 ^[12,14]
1c		65 ^[12,14]
1d		74 ^[9]
1e		80
1f		72
1g		95

^aAll products characterized by ¹H NMR, ¹³C NMR, and mass spectrometry.^bIsolated yields after neutralization.

EXPERIMENTAL

General Procedure (for 2a)

To the solution of 150 mL of MeOH with cooling at 0°C, HCl gas (17 g) was added with stirring for 15 min. The mixture was stirred for 15 min at rt and was carefully added to ethylenediamine (28 g, 0.466 mol) at 0°C. The mixture was stirred for 15 min at room temperature before adding 50 mL of H₂O and stirring for another 0.5 h. To the solution (BOC)₂O (101 g, 0.466 mol) in 200 mL of MeOH was added at room temperature for 10 min, and the resultant solution was stirred for 1 h. (A little better yield would be obtained by adding 1.5 mol equivalents of (BOC)₂O.) The mixture was concentrated in vacuo. Unreacted diamine was removed by diethyl ether (300 mL × 2). The residue was treated with 2 N NaOH (500 mL) solution. The product in the organic layer was extracted with CH₂Cl₂ (300 mL × 3). The combined extracts were washed with 300 mL of brine, dried over anhydrous MgSO₄, and concentrated in vacuo to yield 64.6 g (87%) of mono-BOC product as a colorless oil with more than 97% purity by HPLC.

Data

2a. Oil, ¹H NMR (200 MHz, CDCl₃) δ 1.42 (s, 9H), 1.92 (br s, 2H), 2.80 (t, *J* = 6.0 Hz, 2H), 3.14–3.19 (q, *J* = 6.0 Hz, 2H), 5.13 (br s, 1H). ¹³C

NMR (CDCl₃, 50.3 MHz) δ 28.5, 41.8, 43.2, 79.2, 156.3. MS (FAB): m/z = 161.1 (MH⁺).

2b. Oil, ¹H NMR (200 MHz, CDCl₃) δ 1.42 (s, 9H), 1.57–1.69 (m, 2H), 2.65 (br s, 2H), 2.73 (br s, 2H), 3.14–3.24 (m, 2H), 5.15 (br s, 1H). ¹³C NMR (CDCl₃, 50.3 MHz) δ 28.4, 32.6, 38.2, 39.3, 78.7, 156.2. MS (FAB): m/z = 175.1 (MH⁺).

2c. Oil, ¹H NMR (200 MHz, CDCl₃) δ 1.41 (s, 9H), 1.36–1.49 (m, 4H), 2.12 (br s, 2H), 2.65–2.69 (m, 2H), 2.97–3.08 (m, 2H), 4.87 (br s, 1H). ¹³C NMR (CDCl₃, 50.3 MHz) δ 27.4, 28.3, 30.2, 40.5, 41.7, 78.8, 156.1. MS (FAB): m/z = 189.1 (MH⁺).

2d. Oil, ¹H NMR (200 MHz, CDCl₃) δ 1.46 (s, 9H), 1.93 (br s, 2H), 2.60 (br s, 2H), 2.94–3.11 (m, 2H), 3.20–3.41 (m, 8H), 5.40 (br s, 1H). ¹³C NMR (CDCl₃, 50.3 MHz) δ 28.3, 40.2, 41.4, 70.7, 73.0, 78.7, 156.0. MS (FAB): m/z = 248.2 (MH⁺).

2e. Oil, ¹H NMR (200 MHz, CDCl₃) δ 1.03–1.54 (m, 5H), 1.43 (s, 9H), 1.60–1.69 (m, 2H), 1.85–1.96 (m, 2H), 2.30 (t, J = 10 Hz, 1H), 3.00 (t, J = 9.8 Hz, 1H), 3.07–3.14 (m, 1H), 4.06 (d, J = 3.4 Hz, 1H). ¹³C NMR (CDCl₃, 50.3 MHz) δ 25.2, 25.3, 28.5, 33.0, 35.2, 55.7, 57.7, 79.3, 156.2. MS (FAB): m/z = 215.1 (MH⁺).

2f. Oil, ¹H NMR (200 MHz, CDCl₃) δ 0.97–1.08 (m, 3H), 1.38 (s, 9H), 2.51–2.73 (m, 4H), 3.12–3.18 (m, 2H), 5.01 (broad s, 1H). ¹³C NMR (CDCl₃, 50.3 MHz) δ 15.0, 28.2, 40.0, 43.5, 48.9, 78.6, 156.1. MS (FAB): m/z = 189.1 (MH⁺).

2g. Oil, ¹H NMR (200 MHz, CDCl₃) δ 1.01 (d, J = 6.7 Hz, 6H), 1.37 (s, 9H), 2.61 (t, J = 6.6 Hz, 2H), 2.78 (sept, J = 6.0 Hz, 1H), 2.93 (t, J = 6.6 Hz, 2H), 5.62 (broad s, 1H). ¹³C NMR (CDCl₃, 50.3 MHz) δ 22.5, 28.3, 44.5, 48.7, 53.4, 78.5, 156.2. MS (FAB): m/z = 203.1 (MH⁺).

ACKNOWLEDGMENTS

We gratefully acknowledge the financial support from the Korea Science and Engineering Foundation (R01-2005-000-10032-0 and the Center for Bioactive Molecular Hybrids to H. J. H. and KRF-2002-070-C00060 to W. K. L.). The partial support from the Hufs research grant (2006) is also gratefully acknowledged.

REFERENCES

1. Tripathia, R. P.; Tiwaria, V. K.; Tewari, N.; Katiyar, D.; Saxena, N.; Sinha, S.; Gaikwad, A.; Srivastava, A.; Chaturvedi, V.; Manju, Y. K.; Srivastav, R.; Srivastav, B. S. Synthesis and antitubercular activities of bis-glycosylated diamino alcohols. *Bioorg. Med. Chem.* **2005**, *13*, 5668–5679.
2. Fonta, M.; Sanmartinb, C.; Garcias, H.; Contreras, S.; Paeilec, C.; Bilbenyc, N. A new polyamine derivative, a structural analog of spermine, with in vivo activity as an inhibitor of ethanol appetite. *Bioorg. Med. Chem.* **2005**, *13*, 4375–4382.
3. Lavrador, K.; Murphy, B.; Saunders, J.; Struthers, S.; Wang, X.; Williams, J. A screening library for peptide activated G-protein coupled receptors, 1: The test set. *J. Med. Chem.* **2004**, *47*, 6864–6974.
4. Gardner, R. A.; Delcros, J.-G.; Konate, F.; Breitbeil, F., III; Martin, B.; Sigman, M.; Huang, M.; Phanstiel, O., IV. N¹-Substituent effects in the selective delivery of polyamine conjugates into cells containing active polyamine transporters. *J. Med. Chem.* **2004**, *47*, 6055–6069.
5. Melchiorre, C.; Bolognesi, M. L.; Budriesi, R.; Ghelardini, C.; Chiarini, A.; Minarini, A.; Rosini, M.; Tumiatti, V.; Wade, E. J. Design, synthesis, and biological activity of methoctramine-related polyamines as putative G_i protein activators, 1: The test set. *J. Med. Chem.* **2001**, *44*, 4035–4038.
6. Reddy, V. K.; Valasinas, A.; Sarkar, A.; Basu, H. S.; Marton, L. J.; Frydman, B. Conformationally restricted analogues of ¹N,¹²N-bisethylspermine: Synthesis and growth inhibitory effects on human tumor cell lines. *J. Med. Chem.* **1998**, *41*, 4723–4732.
7. Bolognesi, M. L.; Minarini, A.; Budriesi, R.; Cacciaguerra, S.; Chiarini, A.; Spampinato, S.; Tumiatti, V.; Melchiorre, C. Universal template approach to drug design: Polyamines as selective muscarinic receptor antagonists. *J. Med. Chem.* **1998**, *41*, 4150–4160.
8. Kohn, M.; Wacker, R.; Peters, C.; Schroder, H.; Souler, L.; Breinbauer, R.; Niemeyer, C. M.; Waldmann, H. Staudinger ligation: A new immobilization strategy for the preparation of small-molecule arrays. *Angew. Chem. Int. Ed.* **2003**, *42*, 5830–5834.
9. Kanoh, N.; Kumashiro, S.; Simizu, S.; Kondoh, Y.; Hatakeyama, S.; Tashiro, H.; Osada, H. Immobilization of natural products on glass slides by using a photoaffinity reaction and the detection of protein–small molecule interactions. *Angew. Chem. Int. Ed.* **2003**, *42*, 5584–5587.
10. Oda, Y.; Owa, T.; Sato, T.; Boucher, B.; Daniels, S.; Yamanaka, H.; Shinohara, Y.; Yokoi, A.; Kuromitsu, J.; Nagasu, T. Quantitative chemical proteomics for identifying candidate drug targets. *Anal. Chem.* **2003**, *75*, 2159–2165.
11. Nagle, A. S.; Salvatore, R. N.; Cross, R. M.; Kapxhiu, E. A.; Sahab, S.; Yoon, C. H.; Jung, K. H. Selective mono protection of diols, diamines, and amino alcohols using cesium bases. *Tetrahedron Lett.* **2003**, *44*, 5695–5698.
12. Pittelkow, M.; Lewinsky, R.; Christensen, J. B. Selective synthesis of carbamate protected polyamines using alkyl phenyl carbonates. *Synlett* **2002**, 2159–2202.
13. Krapcho, A. P.; Maresch, M. J.; Lunn, J. Mono-(BOC)-protected diamines: Synthesis of tert-butyl-N-alkyl-N-(2-aminoethyl)carbamates and tert-butyl-N-[2-(alkylamino)ethyl]carbamates. *Synth. Commun.* **1993**, *23*, 2443–2449.
14. Krapcho, A. P.; Kuell, C. S. Mono-protected diamines: N-tert-butylcarbonyl α,ω -alkanediamines from α,ω -alkanediamines. *Synth. Commun.* **1990**, *20*, 2559–2564.

15. Favre-Reguillon, A.; Segat-Dioury, F.; Nait-Bouda, L.; Cosma, C.; Siaugue, G.-M.; Foos, J.; Guy, A. A highly chemoselective protection and activation of primary amines in polyamine. *Synlett* **2000**, 868–870.
16. Ham, J.-Y.; Kang, H.-S. A new convenient method for the monoprotection of α,ω -alkanediamines. *Bull. Kor. Chem. Soc.* **1994**, *15*, 1025–1026.
17. Jacobson, A. R.; Makris, A. N.; Sayre, L. M. Monoacylation of symmetrical diamine. *J. Org. Chem.* **1987**, *52*, 2592–2594.
18. Stahl, G.; Walter, R.; Smith, C. W. Mono-N-acylated 1,6-diaminohexanes. *J. Org. Chem.* **1978**, *43*, 2285–2286.
19. Greene, T. W.; Wuts, P. G. M. *Protective Groups in Organic Synthesis*, 3rd ed.; John Wiley & Sons: New York, 1999, pp. 494–653.
20. Eisenfuhr, A.; Arora, P. S.; Sengle, G.; Takaoka, L. R.; Nowich, J. S.; Famulok, M. A ribozyme with Michaelase activity: Synthesis of substrate precursors. *Bioorg. Med. Chem. Lett.* **2003**, *11*, 235–249.
21. Chhabra, S. R.; Mahajan, A.; Chan, W. C. Homochiral 4-azalysine building blocks: Syntheses and applications in solid-phase chemistry. *J. Org. Chem.* **2002**, *67*, 4017–4029.
22. Chan, P. C. M.; Roon, R. J.; Koerner, J. F.; Taylor, N. J.; Honek, J. F. A 3-amino-4-hydroxy-3-cyclobutene-1,2-dione-containing glutamate analogue exhibiting high affinity to excitatory acid receptor. *J. Med. Chem.* **1995**, *38*, 4433–4438.