3.31 (1 H, d, J = 13.8 Hz), 3.21 (1 H, d, J = 13.8 Hz), 2.64–2.41 (5 H, m), 2.43 (3 H, s), 2.31–2.14 (2 H, m), 2.13 (3 H, s), 1.42 (2 H, s), 0.38 (3 H, d, J = 7.2 Hz), -0.06 (9 H, s).

Ring Expansion of Allylic Chloride 19 to Sulfur-Bridged Carbocycle 23. To a solution of allylic chloride 19 (335 mg, 531 µmol) in CH<sub>3</sub>CN (52 mL) was added anhydrous  $K_2CO_3$  (flame-dried under vacuum immediately prior to use, 0.42 g, 3.0 mmol) and a solution of NaI in CH<sub>3</sub>CN (prepared by dissolving vacuum dried NaI in CH<sub>3</sub>CN, 0.47 M, 1.13 mL, 531  $\mu$ mol). The resulting heterogeneous mixture was heated at 68-70 °C for 17 h, cooled to room temperature and then partitioned between ether (150 mL) and brine (75 mL). The layers were separated, and the organics were washed with brine (75 mL). After the combined aqueous layers were extracted with ether (2  $\times$  75 mL), the combined organics were dried (MgSO<sub>4</sub>) and filtered, and the solvents were evaporated. Flash silica gel chromatography (column 1 in. × 8 in., 2:2:6 ether/CH2Cl2/hexane) provided first the sulfur-bridged carbocycles 23 + 24, 225 mg (71%), followed by a mixture of allylic halides (X = Cl, I), ca. 20 mg, and finally the divinylcyclopropane 25, 26 mg (9%). Sul-fur-bridged carbocycle 23 (major diastereomer): White crystals from ether/hexane; mp 159-161 °C dec; analytical TLC (silica gel F254), 1:1:2 ether/CH<sub>2</sub>Cl<sub>2</sub>/hexane,  $R_f = 0.37$ ; MS exact mass calcd for C<sub>33</sub>- $H_{43}O_5NSSi = 593.2631$ , found = 593.2604, error = 4.6 ppm; IR (neat, cm<sup>-1</sup>) C=O 1735, C=O 1730, C=O 1700; 270-MHz <sup>1</sup>H NMR (CD-Cl<sub>3</sub>)  $\delta$  7.38–7.11 (5 H, m); 6.15 (1 H, dd, J = 10.4, 15.8 Hz), 6.01–5.84 (1 H, m), 5.30 (1 H, s), 5.28 (1 H, d, J = 6.5 Hz), 4.25-4.13 (1 H, m),3.99 (1 H, dt, J = 10.7, 2.1 Hz), 3.62 (1 H, dd, J = 3.6, 11.9 Hz), 3.34(2 H, dd, J = 2.8, 12.5 Hz), 2.95-2.78 (1 H, m), 2.65-1.97 (8 H, m),2.44 (3 H, s), 2.25 (3 H, s), 1.38 (2 H, s), 0.34 (3 H, d, J = 7.1 Hz), -0.07 (9 H, s). Sulfur-bridged carbocycle 24 (minor diastereomer): oil, analytical TLC (silica gel F254), 1:1:2 ether/CH<sub>2</sub>Cl<sub>2</sub>/hexane,  $R_f = 0.38$ ; 270-MHz <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.38-7.09 (5 H, m), 6.19 (1 H, dd, J =10.6, 16.0 Hz), 5.27 (1 H, ddd, J = 4.7, 11.0, 16.0 Hz), 5.26 (1 H, d, J = 10.7 Hz), 5.16 (1 H, s), 3.76 (1 H, dt, J = 3.0, 9.5 Hz), 3.71 (1 H, dd, J = 1.8, 5.6 Hz), 3.34 (1 H, ddd, J = 1.8, 4.2, 12.2 Hz), 3.24 (1 H, dd, J = 2.6, 12.8 Hz), 3.18 (1 H, dd, J = 2.6, 12.8 Hz), 3.09 (1 H, d, J = 10.7 Hz), 2.81 (1 H, ddd, J = 4.5, 10.7, 13.0 Hz), 2.51–2.22 (3 H, m), 2.46 (3 H, s), 2.16-2.02 (4 H, m), 2.13 (3 H, s), 1.44 (2 H, s), 0.37 (3 H, d, J = 7.1 Hz), 0.06 (9 H, s). Vinylcyclopropane 25: white crystals from ether/hexane; mp 166–167 °C; analytical TLC (silica gel F254), 1:1:2 ether/CH<sub>2</sub>Cl<sub>2</sub>/hexane,  $R_f = 0.19$ ; MS exact mass calcd for C<sub>30</sub>-H<sub>35</sub>O<sub>5</sub>NS = 521.2236, found = 521.2235, error = 0.1 ppm; IR (near cm<sup>-1</sup>) C=O 1755, C=O 1740, C=O 1720, C=O 1700; 270-MHz <sup>1</sup>H NMR (CDCl<sub>3</sub>) & 7.35-7.18 (3 H, m), 7.12-7.04 (2 H, m), 5.42 (1 H,

d, J = 7.7 Hz), 5.35 (1 H, dd, J = 10.1, 17.0 Hz), 5.29 (1 H, s), 5.11 (1 H, s), 4.94 (1 H, d, J = 17.0 Hz), 4.88 (1 H, d, J = 10.1 Hz), 3.93–3.84 (1 H, m), 3.47 (1 H, dd, J = 5.6, 14.0 Hz), 3.28–3.17 (1 H, m), 3.20 (2 H, s), 2.80 (1 H, dd, J = 2.7, 14.0 Hz), 2.59–2.49 (2 H, m), 2.50 (3 H, s); 2.44–2.01 (5 H, m), 1.61 (3 H, s), 1.25 (1 H, dd, J = 5.1, 3.3 Hz), 1.15 (3 H, d, J = 6.3 Hz), 0.91–0.87 (1 H, m).

Bridgehead Methylation of 23. Preparation of 30. Lithium diisopropylamide (LDA) was prepared at 0 °C by addition of n-BuLi (2.00 M in hexanes, 0.20 mL, 400  $\mu$ mol) to a solution of diisopropylamine (Aldrich, distilled from CaH<sub>2</sub>; 59  $\mu$ L, 420  $\mu$ mol) in THF (3.0 mL). The LDA solution was cooled to -78 °C, and a THF solution of the ketone 23 (217 mg, 366 µm, in 7 mL) was added dropwise via cannula over 25 min. The ketone-containing flask was rinsed with additional THF (2.0 mL) which was added to the reaction. The solution was stirred at -78 °C for 1 h before iodomethane (Aldrich, freshly distilled from  $P_2O_5$ ; 0.23 mL, 3.7 mmol) was added all at once via syringe. The reaction was stirred at -78 °C for 50 min and was then warmed to 0 °C with an ice bath for 45 min. The reaction was quenched by addition of saturated NH<sub>4</sub>Cl (1 mL) and then partitioned between saturated NH<sub>4</sub>Cl (20 mL) and ether (60 mL), and the layers were separated. The organics were washed with saturated NH<sub>4</sub>Cl (20 mL), saturated NaHCO<sub>3</sub> (20 mL), and brine (20 mL). The combined aqueous layers were extracted with ether  $(2 \times 25 \text{ mL})$ , dried (MgSO<sub>4</sub>), and filtered, and the solvents were removed in vacuo. The residue was chromatographed on silica gel (1 in.  $\times$  8 in.) and eluted (1:1:2 ether/CH<sub>2</sub>Cl<sub>2</sub>/hexane) to give bridgehead methylated ketone 30, 220 mg (99%): white crystals from ether/hexane; mp 195-196 °C; analytical TLC (silica gel F254), 1:1:2 ether/ CH<sub>2</sub>Cl<sub>2</sub>/hexane,  $R_f = 0.47$ ; MS exact mass calcd for C<sub>14</sub>H<sub>45</sub>O<sub>5</sub>NSSi = 607.2788, found = 607.2792, error = 0.7 ppm; IR (CHCl<sub>3</sub>, cm<sup>-1</sup>) C=O 1730, C=O 1705, OAc 1300; 270-MHz <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  7.35-7.15 (5 H, m), 6.16 (1 H, dd, J = 10.1, 15.8 Hz), 6.06-5.92 (1 H, m), 5.28(1 H, s), 5.23 (1 H, d, J = 7.4 Hz), 4.29-4.19 (1 H, m), 3.98 (1 H, d, d)J = 9.8 Hz), 3.42-3.29 (2 H, m), 2.72-2.35 (6 H, m), 2.44 (3 H, s), 2.26 (3 H, s), 2.25-1.99 (3 H, m), 1.42 (3 H, s), 1.38 (2 H, s), 0.35 (3 H, d, J = 7.1 Hz), -0.07 (9 H, s).

Acknowledgment. This work was supported by the National Institutes of Health (Grant No. CA17918).

Supplementary Material Available: Table of mass spectral fragmentation patterns (1 page). Ordering information is given on any current masthead page.

# The Total Syntheses of dl-Zygosporin E and dl-C<sub>18</sub>-Desmethylcytochalasin D

### E. Vedejs\* and S. J. Wittenberger

Contribution from the Chemistry Department, University of Wisconsin, Madison, Wisconsin 53706. Received September 7, 1989

Abstract: The title syntheses are completed starting from the sulfur-bridged 11-membered carbocycles 1 and 2. Key steps include the reductive cleavage of the C-S bond from 11 to 14 and from 2 to 22 and the conversion of allylic silanes 15 and 22 into allylic alcohols 17 and 24, respectively. In the zygosporin synthesis, final sulfur removal is achieved by sulfoxide elimination from 18a to 19a, while in the desmethylcytochalasin D (32) synthesis, sulfur elimination occurs during conversion of allylic sulfoxide 29 to 31. Both synthetic routes depend on control of relative stereochemistry based on local conformational preferences and on the predictable consequences of reactions that introduce stereochemistry in the vicinity of C-S bonds.

The preceding paper describes the synthesis of sulfur-bridged [11]cytochalasans 1 and 2.<sup>1</sup> Five of the six asymmetric centers of the isoindolone subunit have been controlled by the Diels-Alder synthesis, and the acetoxy stereochemistry at cytochalasan  $C_{21}$  has been introduced by using methods that rely upon sulfur-

mediated control of relative stereochemistry. Procedures for the attachment of a hydroxyl group at  $C_7$  with the necessary  $\beta$  orientation have also been established. Further conversion into cytochalasins,<sup>2</sup> zygosporins,<sup>2</sup> or their analogues now requires control of  $C_{16}$  and  $C_{18}$  stereochemistry, as well as eventual removal of sulfur. The  $C_{16}$ —sulfur bond must be replaced by hydrogen,

<sup>(1)</sup> Vedejs, E.; Reid, J. G.; Rodgers, J. D.; Wittenberger, S. J. J. Am. Chem. Soc., preceding paper in this issue. For a preliminary account of the zygosporin E synthesis, see: Vedejs, E.; Rodgers, J. D.; Wittenberger, S. J. J. Am. Chem. Soc. 1988, 110, 4822.

<sup>(2)</sup> Tanenbaum, S. W. Ed. Cytochalasins: Biochemical and Cell Biological Aspects; North-Holland Publishing Company: Amsterdam, 1978.



Figure 1. NOE study of 10.

Scheme 1

ó ETHYLCYTOCHALASIN D ZYGOSPORIN E DES (DMCD) R- H (ŽE) CYTOCHALASIN D (CD) R- MA 3 R. M 5 2 B= H. B'= M R= H, X= H 6 R= H, R'= CO2CH2CCI3 R= H, X= SiMe<sub>2</sub>tBu 7 R= Me, R'= CO<sub>2</sub>CH<sub>2</sub>CCl<sub>3</sub> 10 R= Me, X= H

while the C<sub>20</sub> C-S linkage ideally would serve as the precursor of the  $C_{19} = C_{20}$  olefinic linkage.

The original plan called for enolate generation from 1 at  $C_{18}$ and methylation from the less hindered  $\beta$ -face to give 3. Subsequent reductive  $C_{16}$ -S bond cleavage to enolate 4 was expected to allow  $C_{16}$  methylation from the peripheral (desired) direction as shown in eq 1 (Scheme I). This stereochemical prediction assumes that the favored alkylation transition state will be determined by the preference for a pseudoequatorial  $C_{18}$  methyl group (local conformer control).<sup>3</sup> However, deprotonation of 1 at C<sub>18</sub> could not be achieved directly due to the surprisingly high kinetic acidity of the bridgehead proton (C<sub>16</sub>-H). All attempts to induce kinetically or thermodynamically controlled enolization afforded the enolate 5. It was possible to block the  $C_{16}$  position, as in the  $\beta$ -keto ester 6, but a practical overall yield for the alkylation at  $C_{18}$  with subsequent deblocking of 7 to 3 could not be achieved. The simplest alternative was to proceed with 2 and to control  $C_{16}$  stereochemistry by other means. This approach was eventually successful, but a series of problems had to overcome

Direct methylation of 2 at  $C_{18}$  was precluded by competing enolization at the N-acetyl group. The problem was solved by deacylation to 8 (K<sub>2</sub>CO<sub>3</sub>/MeOH) and temporary protection as the N-tert-butyldimethylsilyl amide 9 (TBSCI/DMAP, TBS =  $Me_2tBuSi$ , DMAP = (dimethylamino)pyridine). Enolization with lithium hexamethydisilazide (LiHMDS) then occurred without further difficulties, and  $C_{18}$  methylation gave a single product 10 (60% from 2) after cleavage of the N-silyl protecting group with







Scheme II



Et<sub>3</sub>NHF. The desired C<sub>18</sub> stereochemistry of 10 corresponds to alkylation from the less congested  $\beta$ -face of the enolate, subject to control by the  $C_{20}$  stereochemistry.<sup>1</sup> This assignment was supported by an NOE study, summarized in Figure 1, and by the eventual conversion into zygosporin E.

19b

Reductive cleavage of the C<sub>16</sub> C-S bond was the next obstacle. The sulfide was relatively unreactive toward zinc reduction, probably because the C-S bond is nearly orthogonal to the activating carbonyl  $\pi$ -orbitals. This problem had been overcome in model systems by sulfide activation via S-methylation and reductive cleavage of the sulfonium salt.<sup>4</sup> The best results were achieved by reacylating 10 at nitrogen to prevent competing lactam O-methylation (Ac<sub>2</sub>O/DMAP, 88%), followed by S-methylation with trimethyloxonium tetrafluorobrate in the presence of allyltrimethylsilane as an acid scavenger. The resulting sulfonium salt 12 (Scheme II) was then reductively cleaved by Rieke zinc<sup>5</sup> with acetic acid as the proton donor. This procedure afforded a mixture of two methyl sulfides 14a and 14b in a ratio of 1:2.6 (87% yield). Replacement of acetic acid by methanol as the proton donor in the reductive cleavage step gave ratios of 14a:14b as high as 1:4.3.

The stereochemistry of the two diastereomers 14a and 14b was not initially known with certainty, but the assignment follows from a consideration of enolate geometry. Due to the constraints imposed by the bicyclic framework, only the enolate isomer 13 can be formed from 12. The enolate double bond (formally, the E isomer) has the "Z" configuration with respect to medium ring substituents. It will therefore prefer a local geometry 13b that is controlled by the tendency of the  $C_{18}$  methyl group to occupy a pseudoequatorial orientation.<sup>3</sup> An alternative local conformation 13a encounters reduced eclipsing interactions between  $C_{18}$ -CH<sub>3</sub> and C17 oxygen, but this conformation suffers from increased transannular interactions with the pseudoaxial C<sub>18</sub>-CH<sub>3</sub> group. Assuming that the transition state for zinc enolate protonation resembles the enolates, protonation should occur preferentially

<sup>(4)</sup> Vedejs, E.; Reid, J. G. J. Am. Chem. Soc. 1984, 106, 4617. Vedejs,
E.; Fedde, C. L.; Schwartz, C. E. J. Org. Chem. 1987, 52, 4269.
(5) Arnold, R. T.; Kulenovic, S. T. Synth. Commun. 1977, 7, 223.

via 13b, resulting in 14b as the major product. Since the order of attachment of methyl and hydrogen substituents at  $C_{16}$  in the above scheme was inverted relative to the initial plan, the stereochemical outcome to favor the unnatural  $C_{16}$  epimer was no surprise. The assignment was eventually confirmed by the conversion of 14a into zygosporin E, as described shortly.

Even though good selectivity for 14a could not be achieved, the sequence allowed completion of a synthesis of *dl*-zygosporin E (19a, Scheme III). Both the natural (a) and  $C_{16}$ -epi (b) series were taken through subsequent steps without major problems. First, cleavage of the N-acetyl group with methanolic potassium carbonate produced 15 without complications from  $C_{21}$  acetate cleavage (a series, 70% b series, 82%). Conversion of the allyl silane into the allylic alcohol at C7 was then performed with the reagent prepared from diphenyl diselenide + trimethyloxonium tetrafluorobrate.<sup>6</sup> This source of electrophilic selenium generated the thermodynamically more stable allyl selenide 16 in excellent yield. An isomeric  $C_7$  selenide is probably the kinetic product and is rapidly isomerized to 16 under the reaction conditions. Subsequent treatment of the selenides 16 with sodium periodate resulted in selective oxidation and spontaneous selenoxide rearrangement to the allylic alcohols 17 (a series, 85%; b series, 99% from 15) with C–O bonding from the less hindered  $\beta$ -alkene face.

With 17a and 17b in hand, all that remained was to introduce the  $C_{19}$ ,  $C_{20}$  double bond by the thermal elimination of the sulfoxides 18, available by low-temperature oxidation of 17 with MCPBA (75-78%). Thermolysis of the  $C_{16}$ -epi isomer 18b proceeded smoothly at 135 °C in xylene in the presence of calcium carbonate buffer and gave the alkene 19b in 70% yield. A minor byproduct could not be characterized due to the small amount isolated, but the enol acetate structure 20b is a possibility by analogy to the behavior of 18a. In the natural series starting from 18a, similar thermolysis resulted in elimination of the sulfoxide to dl-zygosporin E (19a, 52%) together with the isomeric 20a (45%). Increased competition from the unusual elimination toward oxygen may be related to the formation of a ca. 1:1 mixture of diastereomers at sulfoxide sulfur in 18a, but the isomers could not be separated to test whether each sulfoxide was the unique precursor of one of the elimination products. Extensive analysis of NMR spectra and chromatographic properties established the identity of the sythetic zygosporin E by comparison with natural material (see Experimental Section).<sup>7</sup>

The successful completion of the zygosporin E synthesis confirmed the stereochemical assignment at C<sub>16</sub> that had been made earlier at the stage of the reductive C-S cleavage step. As already mentioned, the major product from 12 was the epimer 14b, presumably due to local conformer control in the enolate protonation step by the  $C_{18}$  methyl group. However, an alternative approach to control of the  $C_{16}$  stereocenter was possible where the reductive cleavage step was performed earlier, prior to functionalization at  $C_{18}$ . Thus, treatment of 2 with trimethyloxonium tetrafluoroborate (Scheme IV) followed by Rieke zinc as before gave a mixture of 22a and 22b (75% combined). NMR comparisons indicated the expected similarities with the related compounds 14a and 14b. In particular, the chemical shifts of the  $C_{16}$  methyl groups ( $\delta$  1.21 in 14a and 1.30 in 22a; 1.03 or 1.06 in 14b and 1.07 in 22b) supported the stereochemical assignment, as did similar comparisons in derived compounds. As in the zygosporin E series, the ratio of reductive cleavage products depended somewhat on the proton donor, and the optimum ratio of 3.7:1 was obtained by using Rieke zinc in THF containing a small amount of isopropanol. Apparently, the C18-unsubstituted enolate 21 prefers a transition state similar to the conformation **21a** and undergoes protonation from the  $\beta$ -face. Evaluation of alternative conformations of the relatively rigid ring indicates few viable options and suggests that 21b is a reasonable geometry for the transition state leading to the minor product. Since there are no allylic substituents to control local conformational preferences, this is a case where long-range effects have to be taken into Scheme IV



account. Due to the complex environment, only a qualitative evaluation is possible. However, the stability of enolate **21a** relative to **21b** should be improved compared to the methylated analogue **13a** vs **13b** because the unfavorable transannular interaction of  $C_{18}$ -CH<sub>3</sub> is no longer an issue. This factor may be responsible for the inversion in enolate face protonation selectivity to favor the natural epimer **22a**.

The above approach solves the problem of  $C_{16}$  stereochemistry and, in principle, allows entry into the most complex cytochalasin or zygosporin systems.<sup>2</sup> Among the most challenging target structures are the  $\alpha$ -hydroxy ketone members of these families. We have therefore explored the conversion of **22a** into **32**, the  $C_{18}$ desmethyl analogoue of cytochalasin D.<sup>2</sup>

Activation at C<sub>18</sub> could be achieved upon treatment of the major  $C_{16}$  methyl diastereomer 22a with LiHMDS + TMSCl in situ (THF, 78°) resulting in a  $C_{17}$ ,  $C_{18}$  enol silane. The NMR spectrum suggested that a single geometrical isomer of the  $C_{17}$ ,  $C_{18}$  enol ether was formed (geometry unknown), but the spectrum also indicated the presence of byproducts due to enolization of the  $C_{21}\ acetate.$ Reaction of 22a with DBU/TMSCI/CH<sub>3</sub>CN likewise produced a single enol ether isomer but in this case the  $C_6, C_7$  olefin proton signal was too small, indicating byproducts derived from isoindolone double bond migration to the tetrasubstituted position. In order to circumvent this problem, the allylic silane was elaborated into the allylic alchol 24 as in the zygosporin series by using the PhSeSe<sup>+</sup>(Me)Ph·BF<sub>4</sub><sup>-</sup> reagent.<sup>6</sup> After oxidation with NaIO<sub>4</sub> in dioxane/H<sub>2</sub>O and selenoxide rearrangement, the alcohol was protected as the tert-butyldimethylsilyl ether 25 (TBSCI/ DMAP) in ca. 80% overall yield from 22a. Trimethylsilyl enol ether formation under the DBU/TMSCl/CH3CN conditions now proved to be uneventful, and 25 provided one major  $C_{17}$ ,  $C_{18}$  enol silane 26. No evidence for the more substituted enol silane resulting from enolization at  $C_{16}$  was observed. The reasons for this selectivity are not known, but the result provides an opportunity to combine the ultimate sulfur removal step with the introduction of oxygen functionality at  $C_{18}$  via allylic sulfoxide rearrangement.

To set the stage for sulfur removal, 26 was reacted with the selenenylating reagent PhSeSe<sup>+</sup>(Me)Ph·BF<sub>4</sub><sup>-</sup> (used earlier for a different purpose) to yield a single major C<sub>18</sub> selenide 27 (93%). The keto selenide 27 was then treated with MCPBA at -78 °C to induce oxidation at both selenium and sulfur, resulting in the

<sup>(6)</sup> Vedejs, E.; Rodgers, J. D.; Wittenberger, S. J. Tetrahedron Lett. 1988, 29, 2287.

#### Table I. <sup>1</sup>H NMR Comparisons



proton(s)	chemical shifts (CDCl <sub>3</sub> , ppm)		
	$\frac{DMCD (32)}{(X = H, Y = OH)}$	$CD \\ (X = Me, Y = OH)$	ZE (19a) (X = Me, Y = H)
H <sub>21</sub>	5.55	5.62	5.52
H <sub>20</sub>	6.50	6.10	6.01
H	5.10	5.12	4.77
C <sub>18</sub> -Me		1.49 (s)	1.26 (d)
C <sub>16</sub> -Me	1.11*	0.91	1.00
His	2.86	ca. 2.8	2.85
H	5.37	5.32	5.28
Hin	5.77	5.69	5.73
Н,	3.82	3.79	3.80
CMe	1.15*	1.18	1.15

\* Indicates that assignments may be reversed.

selenoxide sulfoxide 28. Selenoxide elimination required some warming, but conversion occurred to afford a mixture of enones 29 (66%) and 30 (ca. 20%), each as a single sulfoxide diastereomer. The enones were reluctant to undergo the allylic sulfoxide-sulfenate rearrangement. Usually, the [2,3]-sigmatropic rearrangement occurs at ambient temperature and can be driven toward the allylic alcohol in the presence of thiophiles. In the specific case of cis-enone 29 the conditions required to cause the rearrangement were especially harsh, but heating in trimethylphosphite/n-propanol at 105-110 °C for 23 h provided the rearranged allylic alcohol 31 (53% yield) as the only isolable product. The  $\alpha$ -stereochemistry of the C<sub>18</sub> hydroxyl was assigned with hydroxyl configuration in 31 follows from the well known cyclic transition state governing the [2,3]-sigmatropic rearrangement of the allylic sulfoxide.

The isomeric *E*-enone **30** decomposed under the same forcing conditions without furnishing any isolable allylic alcohol products. Inspection of molecular models indicates that the presence of two *E* double bonds in addition to the trans ring fusion in **30** allows few conformational options, and the transition state for [2,3]-shift is highly destabilized by transannular interactions. The situation is not much better in the cis isomer **29**, but the greater flexibility of the molecule is a helpful factor.

Selective N-deacylation of 31 could be effected ( $K_2CO_3/MeOH-THF$ , -20 °C) followed by silyl ether cleavage with aqueous HF, 0 °C (70% yield) to provide dl-C<sub>18</sub>-desmethyl-cytochalasin D 32. Support for the structure 32 is provided by the spectral evidence. Key signals from the <sup>1</sup>H NMR spectra of relevant [11] cytochalasans are compared in Table I. Although the signals of the different compounds show some chemical shift variations depending on the presence or absence of the C<sub>18</sub>-OH group, signal appearance due to closely analogous coupling constants is qualitatively very similar.

The structure and 11-membered ring stereochemistry of 32 is also supported by characteristic fragmentation patterns in the mass spectrum. Comparison of desmethylcytochalasin D (32), cytochalasin D (CD), and zygosporin E (ZE) revealed interesting trends. As expected, the three compounds (32, CD, and ZE) have in common the loss of several neutral fragments: CO (cyclic ketone), OAc and HOAc (acetate),  $C_7H_7$  (benzyl), and combinations thereof (see supplemental material for a table of mass spectral data for 32, CD, and ZE). There are several homologous fragments that differ by the absence of a CH<sub>2</sub> unit in 32 compared to CD (240 and 254, 268 and 282, 314 and 328, 324 and 338), and there also are several identical fragments (172, 212, 240, 268) in both 32 and CD which underscore the similarity in structure. Most informative is the strong M - 18 fragment corresponding to facile dehydration of diols 32 and CD. This fragment was not unexpected, but the mass spectrum of ZE did not show any sig-



nificant M – 18 peak. Since all three compounds have an identical  $C_7$ -OH group, this implies that the  $C_{18}$  hydroxyls of 32 and CD are responsible for the M – 18 fragment. A possible pathway is depicted in Scheme V. Initial ionization of the  $C_{18}$  oxygen (33) triggers an intramolecular hydrogen transfer from  $C_{15}$  to the  $C_{18}$  oxygen via a six-membered transition state to give allylic radical 34 which loses water to give the observed ion 35. This is a common mechanism for decomposition of alcohol molecular ions.<sup>8</sup> Since the process involves an internal H-transfer, there should be a conformational and stereochemical dependence. The similarity in this key fragmentation pathway between 32 and cytochalasin D therefore provides strong support for the  $C_{18}$  stereochemical assignment for 32.

With the completion of the zygosporin E and  $C_{18}$ -desmethylcytochalasin D syntheses, techniques are established for the control of stereochemistry at all of the asymmetric centers of the [11]cytochalasans. The remaining challenge in this series is cytochalasin D, the most complex of the natural [11]cytochalasans. Further progress on this problem will be reported in due course.

#### **Experimental Section**

See Supplementary Material for details of the route to  $C_{16}$ -epi-zy-gosporin E.

Deacylation of 2 to Amide 8. To a solution of imide 2 (18 mg, 29.6  $\mu$ mol) in THF (0.50 mL) and MeOH (2.0 mL) at 0 °C was added anhydrous  $K_2CO_3$  (flame-dried in vacuo, 30 mg, 220  $\mu$ mol). The heterogeneous reaction mixture was stirred for 35 min before it was diluted with ether (20 mL) and washed with brine (2  $\times$  10 mL). The combined aqueous layers were extracted with ether  $(2 \times 5 \text{ mL})$  and the combined organics were dried (MgSO<sub>4</sub>) and filtered, and the solvents were removed in vacuo. Flash silica gel chromatography (0.5 cm × 15 cm, 40% Et-OAc/hexane) yielded amide 8, 16.5 mg (98%), sufficiently pure for the next step. White crystals from ether/hexane, mp 198-203 °C dec, analytical TLC (silica gel F254), 1:1:2 ether/CH<sub>2</sub>Cl<sub>2</sub>/hexane,  $R_f = 0.16$ ; IR (neat, cm<sup>-1</sup>) N--H 3300, C=O 1730, C=O 1700, C=O 1675; 270-MHz NMR (CDCl<sub>3</sub>)  $\delta$  7.36-7.09 (5 H, m), 6.33 (1 H, dd, J = 10.3, 15.7 Hz); 5.94 (1 H, ddd, J = 5.3, 10.7, 15.7 Hz), 5.42 (1 H, s), 5.34 (1 H, s), 5.18 (1 H, d, J = 7.1 Hz), 4.12 (1 H, ddd, J = 6.5, 7.1, 11.8Hz), 3.38 (1 H, d, J = 10.3 Hz), 3.21–3.12 (1 H, m), 2.82–2.73 (1 H, m), 2.81 (1 H, dd, J = 4.2, 13.0 Hz), 2.69–2.34 (6 H, m), 2.24 (3 H, s), 2.16-2.01 (2 H, m), 1.52 (2 H, s), 1.41 (3 H, s), 1.04 (3 H, d, J = 6.8 Hz), -0.04 (9 H, s).

Silvlation of Amide 8 to N-Silvl Amide 9. To a solution of amide 8 (146 mg, 258 µmol) in CH<sub>3</sub>CN (3.5 mL) were added tert-butyldimethylsilyl chloride (Petrarch; 130 mg, 862 µmol), (dimethylamino)pyridine (DMAP, Aldrich; 25 mg, 208 µmol), and diazabicycloundecene (DBU; Aldrich; distilled from CaH<sub>2</sub>, 0.15 mL, 1.0 mmol) in that order. The solution was stirred at room temperature for 4 h after which time it was diluted with hexane (20 mL) and washed with saturated NaHCO<sub>3</sub>  $(2 \times 10 \text{ mL})$ . The combined aqueous layers were extracted with hexane  $(2 \times 10 \text{ mL})$ . The combined organics were dried (MgSO<sub>4</sub>) and filtered and the solvents were removed in vacuo. The residual oil was flash chromatographed on silica gel (1 in.  $\times$  3 in., washed with 5% Et<sub>3</sub>N/ hexane followed by hexane), and eluted with hexane to remove siliconcontaining impurities, followed by 10% EtOAc/hexane to give the N-silvl amide 9 (133.5 mg, 76%), and then 50% EtOAc/hexane to recover starting material 8 (29.3 mg, 20%): oil; analytical TLC (silica gel F254), 1:4 ether/hexane,  $R_f = 0.22$ ; MS exact mass calcd for  $C_{38}H_{57}NO_5Si_2$ = 679.3547, found = 679.3522, error = 3.7 ppm; IR (neat  $cm^{-1}$ ) C=O 1730, C=O 1705, C=O 1665; 270-MHz NMR (CDCl<sub>3</sub>) δ 7.34-7.06 (5 H, m), 6.35 (1 H, dd, J = 10.4, 15.5 Hz), 5.93 (1 H, ddd, J = 4.5, 10.7, 15.5 Hz), 5.36 (1 H, d, J = 7.5 Hz), 5.23 (1 H, s), 4.28 (1 H, dt, dt)J = 12.8, 6.6 Hz), 3.25 (2 H, d, J = 9.8 Hz), 3.07 (1 H, dd, J = 2.9, 12.8 Hz), 2.94–2.55 (3 H, m), 2.57 (1 H, d, J = 10.8 Hz), 2.46 (1 H, d, J = 12.2 Hz), 2.41–2.29 (1 H, m), 2.24 (3 H, s), 2.18–1.99 (3 H, m), 1.41 (2 H, s), 1.40 (3 H, s), 0.99 (9 H, s), 0.36 (3 H, s), 0.25 (3 H, s), 0.21 (3 H, d, J = 6.8 Hz), -0.07 (9 H, s)

Alkylation of N-Silyl Amide 9 and Desilylation to 10. To a solution of lithium hexamethyldisilazide (64  $\mu$ mol, prepared from HN(SiMe<sub>3</sub>)<sub>2</sub>

(14 µL, 67 µmol) and n-BuLi (2.07 M, 31 µL, 64 µmol) in THF (0.56 mL) at 0 °C) at -78 °C was added N-silyl keto amide 9 (40 mg, 58 µmol) in THF (0.60 mL). The reaction was stirred at -78 °C for 5 min, warmed to -40 °C for 5 min, and then cooled back to -78 °C for 5 min before iodomethane (75  $\mu$ L, 1.2 mmol) was added. The cold bath was removed and the reaction flask was allowed to warm to room temperature over 10 min and quenched by addition of saturated NH<sub>4</sub>Cl (0.5 mL) and ether (2 mL). The layers were separated, and the organics were washed with saturated NH<sub>4</sub>Cl (3 mL), saturated NaHCO<sub>3</sub> (3 mL), and then brine (3 mL). The combined aqueous layers were extracted with ether  $(2 \times 3 \text{ mL})$  and the combined organic layers were dried (MgSO<sub>4</sub>) and filtered, and the solvents were evaporated (aspirator). The residual oil was applied to a flash silica gel plug (0.5 in.  $\times$  3 in.) and eluted with 1:1 EtOAc/hexane to give 38.5 mg of crude product which was dissolved in THF (3 mL) and MeOH (7 mL). To this solution was added Et<sub>3</sub>NHF (0.1 mL, 844  $\mu$ mol) and the reaction was stirred for 1.5 h at room temperature. The reaction was then diluted with ether (70 mL) and washed with brine  $(2 \times 30 \text{ mL})$ . The combined aqueous layers were extracted with ether  $(2 \times 20 \text{ mL})$ . The combined organics were dried (MgSO<sub>4</sub>) and filtered, and the solvents were removed in vacuo. The residual oil was applied to a flash silica gel column  $(1/2 \text{ in.} \times 5 \text{ in.})$  and eluted with 20% ether/hexane to remove silicon containing impurities and then with 50% EtOAc/hexane to give the methylated amide 10, 26.5 mg (79%): oil; analytical TLC (silica gel F254), 1:1 EtOAc/hexane,  $R_f$  = 0.47; MS exact mass calcd for  $C_{33}H_{45}NO_4SSi = 579.2839$ , found = 579.2823, error = 2.7 ppm; IR (neat,  $cm^{-1}$ ) N-H 3270, C=O 1735, aC=O 1705, C=O 1670; 270-MHz NMR (CDCl<sub>3</sub>) δ 7.37-7.09 (5 H, m), 6.37 (1 H, dd, J = 10.4, 15.5 Hz), 6.08 (1 H, ddd, J = 6.6, 8.6, 15.5 Hz); 5.38 (1 H, s), 5.37 (1 H, s), 5.17 (1 H, d, J = 7.1 Hz), 3.98 (1 H, ddd, J = 6.9, 7.1, 9.2 Hz), 3.41 (1 H, d, J = 10.4 Hz), 3.21-3.12 (1 H, m), 2.99 (1 H, ddq, J = 3.9, 6.5, 7.1 Hz), 2.83 (1 H, dd, J = 4.5, 13.7 Hz), 2.71 (1 H, ddd, J = 6.5, 8.9, 14.9 Hz), 2.58 (1 H, dd, J = 9.5, 13.4 Hz), 2.53-2.38 (1 H, m), 2.33-2.29 (2 H, m), 2.24 (3 H, s), 2.11-1.96 (2 H, m), 1.52 (2 H, s), 1.38 (3 H, s), 1.21 (3 H, d, J = 7.1 Hz), 1.05 (3 H, d, J = 7.4 Hz), 0.01 (9 H, s).

Acylation of Amide 10 to Imide 11. To a solution of the amide 10 (26 mg, 44  $\mu$ mol) in THF (1.90 mL) were added Et<sub>3</sub>N (0.11 mL, 0.80 mmol), Ac<sub>2</sub>O (0.21 mL, 2.24 mmol) and DMAP (12.6 mg, 0.10 mmol). The reaction was sealed and stirred at room temperature for 6 days. The reaction was quenched by addition of saturated NaHCO<sub>3</sub> (2 mL) and stirred 10 min before it was diluted with ether (10 mL) and the layers were separated. The organic portion was washed with saturated NaH-CO<sub>3</sub> (7 mL), saturated NH<sub>4</sub>Cl (7 mL), and brine (7 mL). After the combined aqueous layers were extracted with ether  $(2 \times 10 \text{ mL})$ , the combined organic portions were dried (MgSO<sub>4</sub>) and filtered, and the solvents were removed under reduced pressure. The residual oil was placed on a flash silica gel column (1 cm  $\times$  15 cm) and eluted with 2:1:7 ether/CH<sub>2</sub>Cl<sub>2</sub>/hexane to give imide 11, 24.1 mg (88%): oil; analytical TLC (silica gel F254), 2:2:6 ether/CH<sub>2</sub>Cl<sub>2</sub>/hexane,  $R_f = 0.43$ ; MS exact mass calcd for  $C_{35}H_{47}NO_5SSi = 621.2944$ , found = 621.2955, error = 1.7 ppm; IR (neat, cm<sup>-1</sup>) C=O 1730, C=O 1700; 270-MHz NMR (CDCl<sub>3</sub>) & 7.34-7.20 (5 H, m), 6.27-6.19 (2 H, m), 5.33 (1 H, s), 5.25 (1 H, d, J = 7.4 Hz), 4.09-4.01 (2 H, m), 3.48-3.39 (2 H, m), 3.25-3.12(1 H, m), 2.58-2.46 (2 H, m), 2.50 (3 H, s), 2.42-2.33 (2 H, m), 2.30 (3 H, s), 2.29-2.20 (1 H, m), 2.18-2.03 (2 H, m), 1.42 (2 H, s), 1.42 (3 H, s), 1.23 (3 H, d, J = 6.8 Hz), 0.37 (3 H, d, J = 7.1 Hz), -0.03 (9H, s).

**Preparation of Rieke Zinc.**<sup>5</sup> To a solution of naphthalene (106 mg, 828  $\mu$ mol) in THF (1.50 mL) was added several small pieces of freshly cut Na (18.4 mg, 800  $\mu$ mol). The reaction rapidly turned very dark green and was stirred for ca. 3 h to ensure complete reaction of the Na metal. The green solution was then added via cannula to a solution of anhydrous ZnCl<sub>2</sub> (fused in vacuo prior to use; 87 mg, 0.64  $\mu$ mol) in THF (2.0 mL) providing the zinc metal as a very finely divided black suspension (ca. 0.11 M) in THF.

Reductive C-S Cleavage: 14a and 14b. To a solution of keto sulfide 11 (30 mg, 48  $\mu$ mol) and allyltrimethylsilane (Aldrich; 140  $\mu$ L, 969  $\mu$ mol) in DME (1.5 mL) was added Me<sub>3</sub>O<sup>+</sup>BF<sub>4</sub><sup>-</sup> (Alfa; 50 mg, 340  $\mu$ mol) all at once. The heterogeneous reaction mixture was warmed to ca. 35 °C and stirred for 3.25 h at which time TLC analysis indicated complete reaction of starting material. The reaction was cooled to room temperature over 10 min and glacial acetic acid (17.5 M, 50  $\mu$ L, 880  $\mu$ mol) was added followed quickly by freshly prepared Rieke zinc (ca. 0.11 M in THF, 2.0 mL, 0.22 mmol). The black suspension was stirred 20 min at room temperature before it was partitioned between ether (10 mL) and saturated NaHCO<sub>3</sub> (10 mL). The layers were separated, and the organics were washed with brine (10 mL). The combined aqueous layers were extracted with ether (2 × 10 mL). The combined organic portions were dried (MgSO<sub>4</sub>) and filtered, and the solvents were removed. The oily crystalline residue was placed on a flash silica gel column (<sup>1</sup>/<sub>2</sub>) in.  $\times$  5 in.) and eluted first with hexane to remove naphthalene and then with ether. The ethereal fraction was divided into two portions and purified by ATLC (plate 20 cm × 10 cm, 1:1:4 ether/CH<sub>2</sub>Cl<sub>2</sub>/hexane, two elutions) to give as the faster moving component 14b (epi series; 19.3 mg, 63%) and as the lower  $R_f$  component 14a (natural series, 7.2 mg, 24%). 14b: oil; analytical TLC (silica gel F254), 1:1:2 ether/ CH<sub>2</sub>Cl<sub>2</sub>/hexane,  $R_f = 0.55$ ; MS exact mass calcd for C<sub>36</sub>H<sub>51</sub>NO<sub>5</sub>SSi = 637.3257, found = 637.3207, error = 7.9 ppm; IR (neat, cm<sup>-1</sup>) C=O 1730, C==O 1700; 270-MHz NMR (CDCl<sub>3</sub>) & 7.36-7.19 (5 H, m), 6.19 (1 H, dd, J = 10.1, 15.8 Hz), 5.63 (1 H, ddd, J = 5.6, 8.6, 15.8 Hz), 5.25(1 H, d, J = 7.7 Hz), 5.08 (1 H, s), 4.09-4.01 (2 H, m), 3.32 (1 H, dd, J)J = 3.0, 12.8 Hz, 3.12-2.87 (3 H, m), 2.55 (1 H, dd, J = 10.1, 12.8 Hz), 2.49 (3 H, s), 2.46-2.22 (2 H, m), 2.18 (3 H, s), 2.15-1.98 (3 H, m), 1.97 (3 H, s), 1.79 (1 H, ddd, J = 2.4, 9.5, 15.2 Hz), 1.38 (2 H, s), 1.06(3 H, d, J = 6.8 Hz), 1.03 (3 H, d, J = 6.9 Hz), 0.38 (3 H, d, J = 6.9 Hz)Hz), -0.08 (9 H, s). Natural series, 14a: oil; analytical TLC (silica gel F254), 1:1:2 ether/CH<sub>2</sub>Cl<sub>2</sub>/hexane,  $R_f = 0.53$ ; MS exact mass calcd for  $C_{36}H_{51}NO_5SSi = 637.3257$ , found = 637.3240, error = 2.8 ppm; IR (neat, cm<sup>-1</sup>) C=O 1750, C=O 1740, C=O 1735, C=O 1700, 270-MHz NMR (CDCl<sub>3</sub>)  $\delta$  7.39–7.16 (5 H, m), 6.11 (1 H, dd, J = 10.1, 15.7Hz), 5.49 (1 H, ddd, J = 3.8, 10.6, 15.7 Hz), 5.37 (1 H, d, J = 7.4 Hz), 5.12 (1 H, s), 4.08 (1 H, d, J = 10.0 Hz), 3.47-3.32 (3 H, m), 3.09-2.96(2 H, m), 2.91–2.78 (1 H, m), 2.59–2.49 (1 H, m), 2.47 (3 H, s), 2.23 (3 H, s), 2.09 (3 H, s), 2.06–1.92 (5 H, m), 1.34 (2 H, s), 1.21 (3 H, d, J = 6.5 Hz), 1.05 (3 H, d, J = 6.2 Hz), 0.32 (3 H, d, J = 6.8 Hz), -0.10 (9 H, s).

Deacetylation of 14a (Natural Series) to Amide 15a. To a solution of imide 14a (6.0 mg, 9.4 µmol) in THF (0.45 mL) and MeOH (0.75 mL) at -15 °C was added anhydrous K2CO3 (flame-dried in vacuo; 21 mg, 150  $\mu$ mol). The reaction was stirred for 2 h and then was partitioned between EtOAc (10 mL) and brine (10 mL). The layers were separated, and the organic layer was washed with brine (10 mL), extracted with EtOAc ( $2 \times 10$  mL), dried (MgSO<sub>4</sub>) and filtered, and the solvents were removed in vacuo. Flash silica gel chromatography (0.5 cm × 6 cm, 30% EtOAc/hexane) of the residual oil gave amide 15a, 3.9 mg (70%): oil; analytical TLC (silica gel F254), 1:2 EtOAc/hexane,  $R_f = 0.30$ ; MS exact mass calcd for  $C_{34}H_{49}NO_4SSi = 595.3152$ , found = 595.3137, error = 2.5 ppm; IR (neat,  $cm^{-1}$ ) N—H 3420, C=O 1740, C=O 1690; 270-MHz NMR (CDCl<sub>3</sub>)  $\delta$  7.35-7.11 (5 H, m), 6.21 (1 H, ddd, J = 1.2, 10.4, 15.4 Hz, 5.49 (1 H, s), 5.44 (1 H, ddd, J = 4.2, 10.9, 15.4 Hz),5.29 (1 H, d, J = 5.3 Hz), 5.16 (1 H, s), 3.53 (1 H, ddd, J = 1.5, 5.3, 12.7 Hz), 3.25-3.16 (1 H, m), 3.07-2.77 (3 H, m), 2.66 (1 H, dd, J =9.2, 13.0 Hz), 2.56-2.43 (1 H, m), 2.37-2.17 (2 H, m), 2.21 (3 H, s), 2.16 (3 H, s), 2.12–1.91 (4 H, m), 1.47 (2 H, s), 1.18 (3 H, d, J = 6.8Hz), 1.06 (3 H, d, J = 6.8 Hz), 0.98 (3 H, d, J = 7.1 Hz), -0.03 (9 H, s)

**Preparation of Selenylating Reagent PhSeSe<sup>+</sup>(Me)Ph·BF**<sub>3</sub><sup>-,6</sup> To a suspension of Me<sub>3</sub>O<sup>+</sup>·BF<sub>4</sub><sup>-</sup> (Alfa; 152 mg, 1.03 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (1.40 mL) was added PhSeSePh (Aldrich; 343 mg, 1.10 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (3.0 mL) via cannula. The diselenide containing flask was rinsed with additional solvent (1.0 mL, total solvent = 5.40 mL, 0.19 M). The yellow heterogeneous mixture was sealed and stirred at room temperature overnight (12-15 h) in the dark (the reagent darkens upon prolonged exposure to light) by which time all the solids had dissolved and the solution was orange-red in color. The reagent was used as a 0.19 M solution and was stored in a covered desiccator at ca. -30 °C in the freezer.

Selenylation of Allylic Silane 15a to Allylic Selenide 16a. To a solution of PhSeSe<sup>+</sup>(Me)Ph·BF<sub>4</sub><sup>-</sup> (0.19 M in CH<sub>i</sub>Cl<sub>2</sub>, 40  $\mu$ L, 7.5  $\mu$ mol) at -78 °C was added, via cannula, allylic silane 15a (3.9 mg, 6.5  $\mu$ mol) in CH<sub>2</sub>Cl<sub>2</sub> (0.3 mL) and the allylic silane containing flask was rinsed with additional CH<sub>2</sub>Cl<sub>2</sub> (0.2 mL). The reaction was stirred for ca. 7 min and then the cold reaction was poured into saturated NaHCO<sub>3</sub> (5 mL) with  $CH_2Cl_2$  (10 mL). The layers were separated, and the organic layer was washed with brine (5 mL). Following extraction of the aqueous layers  $(CH_2Cl_2, 3 \times 3 mL)$ , the combined organic portions were dried (MgSO<sub>4</sub>) and filtered, and the solvents were evaporated. The residue was placed on a flash silica gel column (0.5 cm  $\times$  5 cm) and eluted with 5% ether/hexane to remove PhSeMe and then with 50% EtOAc/hexane to give allylic selenide **16a**, 4.3 mg (97%): oil; analytical TLC (silica gel F254), 1:2 EtOAc/hexane,  $R_f = 0.23$ ; MS no peak match, parent, M - 59 (OAc) = 620.2102, calcd = 620.2101, error = 0.2 ppm; formula =  $C_{37}H_{45}NO_4SSe$ ; IR (neat, cm<sup>-1</sup>) N-H 3440, C=O 1740, C=O 1710, =O 1690; 270-MHz NMR (CDCl<sub>3</sub>) δ 7.37-7.11 (10 H, m), 6.15 (1 H, dd, J = 10.7, 15.4 Hz), 5.40 (1 H, ddd, J = 3.8, 10.7, 15.4 Hz), 5.35 (1 H, s), 5.29 (1 H, d, J = 5.6 Hz), 5.21 (1 H, s), 4.19-4.09 (1 H, m),3.54 (1 H, d, J = 11.8 Hz), 3.52-3.38 (1 H, m), 3.46 (1 H, d, J = 11.8 Hz), 3.06-2.81 (2 H, m), 2.82 (1 H, dd, J = 5.6, 13.6 Hz), 2.67 (1 H, Hz), 3.06-2.81 (2 H, m), 2.82 (1 H, dd), J = 5.6, 13.6 Hz), 2.67 (1 H, Hz), 3.06-2.81 (2 H, m), 3.82 (1 H, dd), J = 5.6, 13.6 Hz), 2.67 (1 H, Hz), 3.06-2.81 (2 H, m), 3.82 (1 H, dd), J = 5.6, 13.6 Hz), 3.66 (1 H, Hz), 3.66 (1 H, Hz), 3.66 (1 Hz), 3.6dd, J = 9.5, 13.0 Hz), 2.55–2.41 (1 H, m), 2.37–2.21 (2 H, m), 2.19 (3 H, s), 2.14 (3 H, s), 2.11-1.89 (2 H, m), 1.25 (2 H, s), 1.17 (3 H, d, J

= 6.8 Hz), 1.05 (3 H, d, J = 6.8 Hz), 0.86 (3 H, d, J = 6.8 Hz). Oxidation of Selenide 16a to Allylic Alcohol 17a. To a solution of allylic selenide 16a (4.3 mg, 6.3  $\mu$ mol) in dioxane (0.50 mL) and pH 7 buffer solution (1 part Aldrich pH 7 buffer concentrate to 24 parts water, 0.21 mL) at 0 °C was added NaIO<sub>4</sub> (0.168 M in water, 42 µL, 7.0 µmol). The cold bath was immediately removed and the reaction stirred for 2.6 h. The reaction was then partitioned between EtOAc (10 mL) and brine (5 mL). The layers were separated, and the organics were washed with brine (5 mL). The combined aqueous layers were extracted with CH<sub>2</sub>Cl<sub>2</sub>  $(2 \times 5 \text{ mL})$ , dried (MgSO<sub>4</sub>) and filtered, and the solvents were removed in vacuo. The residual oil was placed on a flash silica gel column (0.5  $cm \times 5 cm$ ) and eluted with 10% ether/hexane (PhSeSePh) then EtOAc to give allylic alcohol 17a, 3.0 mg (88%): oil; analytical TLC (silica gel F254), EtOAc,  $R_f = 0.49$ ; MS exact mass calcd for  $C_{31}H_{41}NO_5S = 539.2706$ , found = 539.2719, error = 2.5 ppm; IR (neat, cm<sup>-1</sup>) N—H 3440, O-H 3260, C=O 1650; 270-MHz NMR (CDCl<sub>3</sub>) & 7.37-7.11 (5 H, m), 6.08 (1 H, dd, J = 10.1, 15.4 Hz), 5.63 (1 H, dddd, J = 0.6,4.2, 10.9, 15.4 Hz), 5.50 (1 H, s), 5.36 (1 H, br s), 5.24 (1 H, s), 5.23 (1 H, d, J = 5.6 Hz), 5.04 (1 H, s), 3.95 (1 H, d, J = 10.4 Hz), 3.42 (1 Hz)H, ddd, J = 1.5, 5.6, 12.7 Hz), 3.26–3.19 (1 H, m), 3.09–2.97 (1 H, m), 2.87 (1 H, ddd, J = 2.4, 7.4, 12.2 Hz), 2.81-2.71 (3 H, m), 2.64-2.51 (2 H, m), 2.34 (1 H, t, J = 7.1 Hz), 2.21 (3 H, s), 2.16 (3 H, s), 2.07-1.92 (3 H, m), 1.19 (3 H, d, J = 6.8 Hz), 1.07 (3 H, d, J = 6.8 Hz), 0.88 (3 H, d, J = 6.8 Hz).

Oxidation of Methyl Sulfide 17a to Sulfoxide 18a. To a solution of sulfide 17a (3.0 mg, 5.6 µmol) in CH<sub>2</sub>Cl<sub>2</sub> (1.0 mL) with suspended NaHCO<sub>3</sub> (flame-dried in vacuo, ca. 10 mg, 120 µmol) at -78 °C was added MCPBA (Aldrich; 0.16 M in CH<sub>2</sub>Cl<sub>2</sub>, 40 µL, 6.6 µmol). The reaction was stirred for 30 min before methyl sulfide (Aldrich; 0.16 M in CH<sub>2</sub>Cl<sub>2</sub>, 40 µL, 6.6 µmol). The reaction was stirred for 30 min before methyl sulfide (Aldrich, 10  $\mu$ L, 136  $\mu$ mol) was added to quench excess peroxy acid. After stirring at -78 °C for 10 min the cold bath was removed and the reaction was allowed to warm to room temperature over 5 min and was added to saturated Na<sub>2</sub>CO<sub>3</sub> (5 mL) with CH<sub>2</sub>Cl<sub>2</sub> (10 mL). The layers were separated, and the organics were washed with saturated Na<sub>2</sub>CO<sub>3</sub> (5 mL) and brine (5 mL). The combined aqueous layers were extracted with  $CH_2Cl_2$  (2 × 5 mL), dried (MgSO<sub>4</sub>), and evaporated. The residual oil was placed on a flash silica gel column (0.5 cm  $\times$  5 cm) and eluted with 3:1 EtOAc/hexane to remove nonpolar impurities and then 5% MeOH/EtOAc to give sulfoxide 18a, 2.4 mg (77%) as a ca. 1:1 mixture of diastereomers: oil; analytical TLC (silica gel F254), 1:19 MeOH/EtOAc,  $R_f = 0.51$ ; IR (neat, cm<sup>-1</sup>) N—H 3420, 0-H 3250, C=0 1700, S=0 1040; 270-MHz NMR (CDCl<sub>3</sub>, partial)  $\delta$  7.41–7.08 (5 H, m), 6.10 (0.5 H, dd, J = 8.8, 14.8 Hz), 5.97 (0.5 H, dd, J = 10.0, 15.7 Hz), 5.72-5.56 (1 H, m), 5.67 (0.5 H, s), 5.66 (0.5 H, s), 5.46 (0.5 H, d, J = 6.8 Hz), 5.43 (0.5 H, s), 5.41 (0.5 H, d, J =5.6 Hz), 5.31 (0.5 H, s), 5.08 (1 H, s), 2.87 (1.5 H, s), 2.77 (1.5 H, s), 2.24 (1.5 H, s), 2.19 (1.5 H, s), 1.22 (1.5 H, d, J = 6.5 Hz), 1.21 (1.5 H, d, J = 6.8 Hz), 1.09 (1.5 H, d, J = 6.5 Hz), 1.07 (1.5 H, d, J = 6.8Hz), 0.93 (1.5 H, d, J = 6.8 Hz)

Sulfoxide 18a Elimination to dl-Zygosporin E (19a). To a solution of sulfoxide 18a (2.4 mg, 4.3 µmol) in xylenes (0.4 mL) was added CaCO<sub>3</sub> (flame-dried in vacuo; ca. 20 mg, 200 µmol). The reaction was placed in an oil bath at 135 °C for 55 min, cooled to room temperature, and then filtered through a pad of Celite (CHCl<sub>3</sub>). The solvents were removed in vacuo, and the residue was purified via ATLC (plate 5 cm  $\times$  10 cm, 100% EtOAc) to give two products; the higher R<sub>f</sub> component was identified as dl-zygosporin E (19a), 1.1 mg (52%), and the lower  $R_f$ component was tentatively assigned as enol acetate 20a, 1.0 mg (47%). dl-Zygosporin E (19a) (synthetic): white needles from acetone/hexane; mp 209-210 °C; analytical TLC (silica gel F254), EtOAc,  $R_f = 0.45$ ; MS exact mass calcd for  $C_{30}H_{37}NO_5 = 491.2672$ , found = 491.2695, error = 4.7 ppm; IR (neat, cm<sup>-1</sup>) N-H 3413, O-H 3266, C=O 1733, C=O 1701, C=O 1689; 270-MHz NMR (CDCl<sub>3</sub>) & 7.39-7.09 (5 H, m), 6.01 (1 H, ddd, J = 1.2, 2.4, 15.7 Hz), 5.73 (1 H, dd, J = 9.5, 16.0Hz), 5.52 (1 H, s), 5.45 (1 H, s), 5.34 (1 H, d, J = 1.2 Hz), 5.28 (1 H, ddd, J = 5.3, 10.7, 16.0 Hz), 5.11 (1 H, s), 4.77 (1 H, ddd, J = 2.1, 7.4, ddd, J = 5.3, 10.7, 16.0 Hz), 5.11 (1 H, s), 4.77 (1 H, ddd, J = 2.1, 7.4, ddd, J = 5.3, 10.7, 16.0 Hz), 5.11 (1 H, s), 4.77 (1 H, ddd, J = 5.3, 10.7, 16.0 Hz), 5.11 (1 H, s), 4.77 (1 H, ddd, J = 5.3, 10.7, 16.0 Hz), 5.11 (1 H, s), 4.77 (1 H, ddd, J = 5.3, 10.7, 16.0 Hz), 5.11 (1 H, s), 4.77 (1 H, ddd, J = 5.3, 10.7, 16.0 Hz), 5.11 (1 H, s), 4.77 (1 H, ddd, J = 5.3, 10.7, 16.0 Hz), 5.11 (1 H, s), 4.77 (1 H, ddd, J = 5.3, 10.7, 16.0 Hz), 5.11 (1 H, s), 4.77 (1 H, ddd, J = 5.3, 10.7, 16.0 Hz), 5.11 (1 H, s), 4.77 (1 H, ddd, J = 5.3, 10.7, 16.0 Hz), 5.11 (1 H, s), 4.77 (1 H, ddd, J = 5.3, 10.7, 16.0 Hz), 5.11 (1 H, s), 4.77 (1 H, ddd, J = 5.3, 10.7, 16.0 Hz), 5.11 (1 H, s), 4.77 (1 H, ddd, J = 5.3, 10.7, 16.0 Hz), 5.11 (1 H, s), 4.77 (1 H, ddd, J = 5.3, 10.7, 16.0 Hz), 5.11 (1 H, s), 4.77 (1 H, ddd, J = 5.3, 10.7, 16.0 Hz), 5.11 (1 H, s), 4.77 (1 H, ddd, J = 5.3, 10.7, 16.0 Hz), 5.11 (1 H, s), 4.77 (1 H, ddd, J = 5.3, 17.7, 15.7 Hz), 3.81 (1 H, d, J = 11.8 Hz), 3.28–3.21 (2 H, m), 2.89–2.73 (2 H, m), 2.86 (1 H, d, J = 13.3 Hz), 2.67–2.55 (2 H, m), 2.47–2.31 (1 H, m), 2.26 (3 H, s), 2.10 (1 H, t, J = 4.2 Hz), 2.06–1.90 (2 H, m), 1.26 (3 H, d, J = 6.8 Hz), 1.15 (3 H, d, J = 6.8 Hz), 1.00 (3 H, d, J = 6.8 Hz). Natural zygosporin E:<sup>7</sup> colorless needles from acetone/hexane; mp 218-223.5 °C; analytical TLC (silica gel F254), EtOAc,  $R_f = 0.45$ ; IR (neat cm<sup>-1</sup>) N-H 3414, O-H 3260, C=O 1740, C=O 1706, C=O 1685; 270-MHz NMR (CDCl3) & 7.38-7.09 (5 H, m), 6.01 (1 H, ddd,

J = 1.2, 10.8, 15.8 Hz), 5.73 (1 H, dd, J = 9.8, 15.5 Hz), 5.52 (1 H, dd, J = 2.4, 3.6 Hz, 5.44 (1 H, s), 5.34 (1 H, s), 5.28 (1 H, s), 5.28 (1 H, ddd, J = 4.8, 11.0, 15.5 Hz), 5.11 (1 H, s), 4.77 (1 H, ddd, J = 2.4, 7.5, 1.515.8 Hz), 3.80 (1 H, d, J = 11.0 Hz), 3.29-3.20 (2 H, m), 2.89-2.81 (1 Hz)H, m), 2.85 (1 H, d, J = 13.4 Hz), 2.79-2.58 (2 H, m), 2.64 (1 H, d, J = 13.4 Hz, 2.47–2.35 (1 H, m), 2.26 (3 H, s), 2.10 (1 H, t, J = 4.5Hz), 2.03-1.94 (1 H, m), 1.98 (1 H, d, J = 1.5 Hz), 1.26 (3 H, d, J = 6.8 Hz), 1.15 (3 H, d, J = 6.9 Hz), 1.00 (3 H, d, J = 6.8 Hz). Enol acetate 20a: very fine white needles from ether/hexane; mp 204-205 °C; analytical TLC (silica gel F254), EtOAc,  $R_f = 0.42$ ; MS exact mass calcd for  $C_{30}H_{37}NO_5 = 491.2672$ , found = 491.2637, error = 7.1 ppm; IR (neat,  $cm^{-1}$ ) N-H 3360, O-H 3254, C=O 1756, C=O 1704, C=O 1694; 270-MHz NMR (CDCl<sub>3</sub>) δ 7.38-7.09 (5 H, m), 5.88 (1 H, dd, J = 9.2, 14.2 Hz), 5.49-5.29 (2 H, m), 5.39 (1 H, s), 5.21 (1 H, s), 5.02 (1 H, s), 4.57-4.09 (2 H, m), 3.76-3.60 (2 H, m), 3.25-3.19 (1 H, m), 2.74-2.36 (6 H, m), 2.33 (3 H, s), 2.28-2.20 (2 H, m), 2.07-1.88 (1 H, m), 1.17 (3 H, d, J = 7.1 Hz), 1.14 (3 H, d, J = 7.4 Hz), 0.84 (3 H)H. d. J = 6.8 Hz).

Sulfur Alkylation/Reduction of 2. Preparation of 22a and 22b. To a solution of 2 (11 mg, 18.1 µmol) and allyltrimethylsilane (Aldrich, distilled; 50 µL, 315 µmol) in DME (0.6 mL) was added Me<sub>3</sub>O<sup>+</sup>·BF<sub>4</sub><sup>-</sup> (Alfa; 26.9 mg, 181 µmol) all at once. The heterogeneous reaction mixture was stirred at ambient temperature for 2.3 h by which time all of the sulfide had gone to a baseline spot by TLC. Methyl sulfide (Aldrich; 20  $\mu$ L, 272  $\mu$ mol) was added to quench the excess oxonium salt. The reaction was stirred for 2 min before isopropyl alcohol (EM Science, reagent grade; 50 µL) was added followed by freshly prepared Rieke zinc<sup>5</sup> (ca. 0.2 M in THF, 1.0 mL, 0.2 mol). The black reaction mixture was stirred for ca. 35 min before it was partitioned between ether (10 mL) and saturated  $NH_4Cl$  (5 mL). The layers were separated and the organics were washed with saturated NaHCO3 and then brine (5 mL each). The combined aqueous layers were extracted with ether  $(2 \times 5)$ mL), and the combined organics were dried (MgSO<sub>4</sub>) and filtered, and the solvents were evaporated. The oily crystalline residue was placed on a flash silica gel column (0.5 cm  $\times$  5 cm) and eluted with hexane (naphthalene) followed by 1:3 EtOAc/hexane to give 12.8 mg of crude product. The compound was purified by HPLC (15% EtOAc/hexane) to give the C<sub>16</sub> methyl diastereomers 22a (6.7 mg, 59% yield) and 22b (1.8 mg, 16% yield) in a 3.7:1 ratio. 22a: oil; analytical TLC (silica gel F254), 1:1:2 ether/CH<sub>2</sub>Cl<sub>2</sub>/hexane,  $R_f = 0.54$ ; MS exact mass calcd for  $C_{35}H_{49}NO_5SSi = 623.3101$ , found = 623.3120, error = 3.1 ppm; IR (neat, cm<sup>-1</sup>) C=O 1745, C=O 1725, C=O 1700; 270-MHz NMR  $(CDCl_3) \delta 7.38-7.19 (5 H, m), 6.41 (1 H, ddd, J = 1.2, 10.1, 15.2 Hz),$ 5.76 (1 H, ddd, J = 3.9, 10.7, 15.2 Hz), 5.38 (1 H, d, J = 9.5 Hz), 5.09(1 H, s), 4.09 (1 H, d, J = 10.1 Hz), 3.45-3.31 (2 H, m), 3.16-2.98 (2 H)H, m), 2.73–1.94 (9 H, m), 2.45 (3 H, s), 2.26 (3 H, s), 2.14 (3 H, s), 1.34 (2 H, s), 1.30 (3 H, d, J = 6.9 Hz), 0.29 (3 H, d, J = 7.1 Hz), -0.09 (9 H, s). 22b: white crystals from ether/hexane; mp 170-172 °C, analytical TLC (silica gel F254), 1:1:2 ether/CH<sub>2</sub>Cl<sub>2</sub>/hexane,  $R_f = 0.58$ ; MS exact mass calcd for  $C_{35}H_{49}NO_5SSi = 623.3101$ , found = 623.3101, error = 0.1 ppm; IR (neat, cm<sup>-1</sup>) C=O 1740, C=O 1715, C=O 1700, C=O 1685; 270-MHz NMR (CDCl<sub>3</sub>)  $\delta$  7.38-7.19 (5 H, m), 6.32 (1 H, dd, J = 9.6, 15.0 Hz), 5.59 (1 H, ddd, J = 6.9, 8.1, 15.0 Hz), 5.38 (1 H, d, J = 9.5 Hz), 5.09 (1 H, t, J = 2.4 Hz), 4.08 (1 H, d, J = 10.1 Hz), 3.47-3.27 (2 H, m), 3.07-2.92 (2 H, m), 2.87-2.77 (1 H, m), 2.54 (1 H, t, J = 11.3 Hz), 2.45 (3 H, s), 2.41–2.35 (1 H, m), 2.33–2.26 (2 H, m), 2.24 (3 H, s), 2.15 (3 H, s), 2.12–1.95 (3 H, m), 1.69–1.57 (1 H, m), 1.33 (2 H, s), 1.07 (3 H, d, J = 6.5 Hz), 0.28 (3 H, d, J = 6.9 Hz), -0.09 (9 H, s).

Conversion of Allylic Silane 22a to Silyl Ether 25. To a solution of PhSeSe<sup>+</sup>(Me)Ph BF<sub>4</sub><sup>-</sup> (prepared as described earlier, 125 mM, 285  $\mu$ L, 36 µmol) at -78 °C was added a solution of allylic silane 22a (18.5 mg, 29 µmol) in CH2Cl2 (1.0 mL) via cannula. The silane-containing flask was rinsed with additional solvent (0.4 mL) that was added to the reaction. After ca. 5 min the reaction was poured into a separatory funnel containing saturated NaHCO<sub>3</sub> (5 mL) with CH<sub>2</sub>Cl<sub>2</sub> (10 mL). The layers were separated, the organics were washed with brine (5 mL), and the combined aqueous layers were extracted with  $CH_2Cl_2$  (2 × 5 mL). The combined organic portions were dried (MgSO<sub>4</sub>) and filtered, and the solvents were removed in vacuo. The residue was placed on a flash silica gel plug (1 cm  $\times$  2 cm) and eluted with 5% ether/hexane (PhSeMe) followed by ether to give the allylic selenide 23, 21.5 mg (theoretical yield = 20.5 mg), which was carried on without further purification: oil; analytical TLC (silica gel F254), 1:1:2 ether/CH<sub>2</sub>Cl<sub>2</sub>/hexane,  $R_f = 0.49$ ; MS exact mass calcd for C<sub>38</sub>H<sub>45</sub>NO<sub>5</sub>SSe = 707.2183, found = 707.2178, error = 0.8 ppm; 270-MHz NMR (CDCl<sub>3</sub>) δ 7.42-7.17 (10 H, m), 6.39 (1 H, dd, J = 10.7, 15.1 Hz), 5.72 (1 H, ddd, J = 3.3, 10.7, 15.1 Hz),5.41 (1 H, d, J = 9.0 Hz), 5.39 (1 H, s), 4.27 (1 H, dd, J = 2.7, 11.6 Hz), 3.53-3.42 (2 H, m), 3.37-3.28 (2 H, m), 3.10-2.96 (2 H, m), 2.70-2.51 (3 H, m), 2.36 (3 H, s), 2.30-2.05 (5 H, m), 2.26 (3 H, s),

<sup>(7)</sup> Minato, H.; Katayama, T. J. Chem. Soc. C 1970, 45. We are grateful

to Dr. Minato for a comparison sample of zygosporin E. (8) McLafferty, F. W. Interpretation of Mass Spectra; University Science Books: Mill Valley, CA, 1980.

H, d, J = 6.8 Hz). The allylic selenide 23 from above (21.5 mg,  $29 \mu mol$ ) was dissolved in *p*-dioxane (1.50 mL) to which was added pH 7 buffer solution (Aldrich; 1 part concentrate to 24 parts water, 0.30 mL) and NaIO<sub>4</sub> (0.19 M in H<sub>2</sub>O, 0.46 mL, 89  $\mu$ mol). The reaction was stirred at ambient temperature during which time it gradually yellowed and deposited a white precipitate. After 70 min the excess periodate was quenched by the addition of Na<sub>2</sub>SO<sub>3</sub> (aqueous) and the mixture was partitioned between EtOAc (10 mL) and brine (5 mL). The layers were separated, the organics were washed with brine (5 mL), the aqueous layers were extracted with EtOAc ( $2 \times 5$  mL), the combined organics were dried (MgSO<sub>4</sub>) and filtered, and the solvents were removed. The residue was placed on a flash silica gel plug (1 cm  $\times$  2 cm) and eluted with 15% EtOAc/hexane to give a yellow forerun (PhSeSePh) followed by 3:1 EtOAc/hexane to give ca. 15 mg of crude product. This was then purified by flash silica gel chromatography (column 0.5 cm × 15 cm, 1:1 EtOAc/hexane) to give the rearranged allylic alcohol 24, 13.6 mg (81% from the allylic silane): oil; analytical TLC (silica gel F254), 1:1 Et-OAc/hexane,  $R_f = 0.28$ ; MS exact mass calcd for  $C_{32}H_{41}NO_6S = 567.2654$ , found = 567.2621, error = 5.9 ppm; IR (neat, cm<sup>-1</sup>) O-H 3450, C=O 1744, C=O 1723, C=O 1700; 270-MHz NMR (CDCl<sub>3</sub>)  $\delta$  7.39–7.18 (5 H, m), 6.27 (1 H, dd, J = 9.2, 15.1 Hz), 5.89 (1 H, ddd, J = 3.5, 10.9, 15.1 Hz), 5.36 (1 H, d, J = 9.5 Hz), 5.04 (1 H, s), 4.89 (1 H, s), 4.00 (1 H, dd, J = 1.2, 10.7 Hz), 3.87 (1 H, d, J = 9.5 Hz),3.42 (1 H, dd, J = 2.6, 12.4 Hz), 3.26 (1 H, ddd, J = 2.1, 9.5, 11.6 Hz), 3.08 (1 H, ddd, J = 1.5, 12.7, 14.5 Hz), 2.76 (1 H, dd, J = 9.2, 9.8 Hz),2.75-2.57 (3 H, m), 2.43 (3 H, s), 2.41-2.29 (3 H, m), 2.28 (3 H, s), 2.26-2.20 (1 H, m), 2.14 (3 H, s), 2.10 (1 H, d, J = 6.2 Hz), 1.70 (1 H, s), 1.65-1.55 (1 H, m), 1.31 (3 H, d, J = 7.1 Hz), 0.24 (3 H, d, J= 6.8 Hz).

To a solution of the allylic alcohol 24 (13.6 mg, 24  $\mu$ mol) in DMF (1.1 mL) were added imidazole (Aldrich; 33 mg, 486 µmol), DMAP (Aldrich; 6 mg, 49 μmol) and TBSCI (Petrarch; 28.4 mg, 188 μmol). The resulting solution was stirred at ambient temperature for 16 h and then was diluted with hexane (ca. 15 mL), and washed successively with saturated NaH-CO<sub>3</sub> (5 mL), saturated NH<sub>4</sub>Cl (2 × 5 mL), and brine (5 mL). The combined aqueous layers were extracted with hexane  $(2 \times 5 \text{ mL})$ , dried (MgSO<sub>4</sub>), and filtered, and the solvents were evaporated. The residue was placed on a flash silica gel column (1 cm  $\times$  5 cm) and eluted with hexane to remove silicon impurities followed by 25% EtOAc/hexane to give the crude silvl ether 25 (17.3 mg, theoretical yield = 16.3 mg), sufficiently pure for the next step: oil; analytical TLC (silica gel F254), 15% EtOAc/hexane,  $R_f = 0.19$ ; MS no peak match, parent, loss of Me<sub>3</sub>C, M - 57, 624.2887, calcd = 624.2815, error = 4.4 ppm, formula =  $C_{38}H_{55}NO_6SSi$ ; IR (neat, cm<sup>-1</sup>) C=O 1745, C=O 1723, C=O 1702; 270-MHz NMR (CDCl<sub>3</sub>)  $\delta$  7.41-7.18 (5 H, m), 6.14 (1 H, ddd, J = 0.9, 9.5, 15.4 Hz), 5.67 (1 H, ddd, J = 3.5, 10.3, 15.4 Hz), 5.35 (1 H, d, J = 9.8 Hz), 4.80 (1 H, s), 4.76 (1 H, s), 4.01 (1 H, ddd, J = 0.9, 2.1, 11.2Hz), 3.86 (1 H, d, J = 8.3 Hz), 3.42 (1 H, dd, J = 3.3, 13.0 Hz), 3.23(1 H, ddd, J = 1.5, 9.5, 11.0 Hz), 3.10 (1 H, ddd, J = 1.5, 12.7, 17.4Hz), 2.71-2.56 (5 H, m), 2.42 (3 H, s), 2.39-2.36 (2 H, m), 2.30 (3 H, s), 2.22-2.18 (1 H, m), 2.14 (3 H, s), 2.04 (1 H, dd, J = 0.9, 6.5 Hz), 1.64-1.50 (1 H, m), 1.30 (3 H, d, J = 6.8 Hz), 0.76 (9 H, s), 0.22 (3 H, d, J = 7.1 Hz), -0.08 (6 H, s)

Selenenylation of 25 to Selenide 27 (Major C<sub>16</sub> Methyl Series). A solution of the ketone 25 (17 mg, 24  $\mu$ mol), TMSCl (freshly distilled from CaH<sub>2</sub>; 60  $\mu$ L, 470  $\mu$ mol) and DBU (75  $\mu$ L, 500  $\mu$ mol) in CH<sub>3</sub>CN (2.0 mL) was heated at 70–75 °C for ca. 12 h. The very dark reaction was diluted with hexane (25 mL) and washed with saturated NaHCO<sub>3</sub> (2 × 10 mL) and then 0.1 N citric acid (10 mL). The combined aqueous layers were extracted with hexane (2 × 10 mL), dried (MgSO<sub>4</sub>), and filtered, and the solvents were removed in vacuo. The residue was placed on a short flash silica gel plug (0.5 cm × 5 cm) and eluted with 15% EtOAc/hexane to give the crude silyl enol ether 26, 21 mg (theoretical yield = 18.1 mg), that was used directly in the next step: oil; analytical TLC (silica gel F254), 15% EtOAc/hexane,  $R_f = 0.43$ ; 270-MHz NMR (CDCl<sub>3</sub>, partial)  $\delta$  6.00 (1 H, dd, J = 9.8, 14.5 Hz), 5.43 (1 H, d, J = 9.5 Hz), 5.35 (1 H, ddd, J = 3.5, 10.9, 14.5 Hz), 4.78 (1 H, s), 4.75 (1 H, s), 4.48 (1 H, t, J = 7.7 Hz), 4.03 (1 H, ddd, J = 1.2, 2.7, 10.9 Hz).

To a solution of PhSeSe<sup>+</sup>(Me)Ph·BF<sub>4</sub><sup>-</sup> prepared as described earlier (125 mM, 0.23 mL, 29  $\mu$ mol) at -78 °C was added the silyl enol ether 26 (21.3 mg, 24  $\mu$ mol) in CH<sub>2</sub>Cl<sub>2</sub> (0.7 mL) via cannula. The silyl enol ether containing flask was rinsed with additional solvent (0.3 mL) that was added to the reaction. After ca. 10 min the reaction was poured into a separatory funnel containg saturated NaHCO<sub>3</sub> (5 mL) with CH<sub>2</sub>Cl<sub>2</sub> (10 mL). The layers were separated, the organics were washed with brine (5 mL), aqueous layers were extracted with CH<sub>2</sub>Cl<sub>2</sub> (2 × 5 mL), organics were dried (MgSO<sub>4</sub>) and filtered, and solvents were removed. The residue was placed on a flash silica gel column (1 cm × 6 cm) and eluted with 5% ether/hexane (PhSeMe) followed by ether to give  $C_{18}$ -selenide 27, 18.6 mg (93% two steps): oil; analytical TLC (silica gel F254), 15% EtOAc/hexane,  $R_f = 0.23$ ; IR (neat, cm<sup>-1</sup>) C=O 1747, C=O 1723, C=O 1700, C=O 1690; 270-MHz NMR (CDCl<sub>3</sub>)  $\delta$  7.62-7.18 (10 H, m), 6.15 (1 H, ddd, J = 1.6, 10.4, 15.4 Hz), 5.66 (1 H, ddd, J = 3.9, 10.7, 15.4 Hz), 5.29 (1 H, d, J = 10.1 Hz), 4.78 (1 H, s), 4.75 (1 H, s), 4.03 (1 H, dd, J = 2.7, 11.6 Hz), 3.98 (1 H, ddd, J = 1.5, 2.7, 11.0 Hz), 3.85 (1 H, d, J = 8.3 Hz), 3.38 (1 H, ddd, J = 2.4, 12.4 Hz), 3.09 (1 H, ddd, J = 1.8, 8.3, 10.1 Hz), 2.92-2.78 (1 H, m), 2.71-2.56 (3 H, m), 2.49-2.24 (3 H, m), 2.37 (3 H, s), 2.06-1.93 (2 H, m), 2.00 (3 H, s), 1.96 (3 H, s), 1.63 (3 H, d, J = 7.1 Hz), 0.75 (9 H, s), 0.19 (3 H, d, J = 6.8 Hz), -0.08 (6 H, s).

Selenoxide Elimination of 27 to Enones 29 and 30. To a stirred suspension of NaHCO3 (flame-dried in vacuo; 3 mg, 35  $\mu mol)$  and C18selenide **27** (5.1 mg, 6.1  $\mu$ mol) in CH<sub>2</sub>Cl<sub>2</sub> (0.25 mL) at -78 °C was added MCPBA (Aldrich; 0.14 M in CH<sub>2</sub>Cl<sub>2</sub>, 94  $\mu$ L, 13.4  $\mu$ mol) via syringe. The reaction was stirred at -78 °C for 35 min before it was allowed to warm slowly to -20 °C over 35 min. At this point methyl sulfide (Aldrich; 5 µL, 68 µmol) and triethylamine (Aldrich, distilled from CaH<sub>2</sub>; ca. 15  $\mu$ L, ca. 100  $\mu$ mol) were added and the cold bath was removed. After warming to room temperature, chloroform (0.5 mL) was added and the reaction was warmed to ca. 50 °C for 70 min. After returning to room temperature, the reaction mixture was washed with saturated Na<sub>2</sub>CO<sub>3</sub> (2 × 5 mL), saturated NH<sub>4</sub>Cl (5 mL), and brine (5 mL). The combined aqueous layers were extracted with CHCl<sub>3</sub> ( $2 \times 5$ mL), the organics were dried (MgSO<sub>4</sub>) and filtered, and the solvents were removed. A flash silica gel plug (0.5 cm  $\times$  5 cm) eluted with 10% EtOAc/hexane (PhSeSePh) followed by EtOAc afforded 4.4 mg of crude product. Purification by ATLC (plate 5 cm × 10 cm, 2:1 EtOAc/hexane) gave as the major product cis-enone 29 (2.8 mg, 66%) and a minor product trans-enone 30 (ca. 0.8 mg, ca. 20%). 29: oil; analytical TLC (silica gel F254), 2:1 EtOAc/hexane,  $R_f = 0.26$ ; IR (neat, cm<sup>-1</sup>) C=O 1750, C=O 1715, C=O 1705, C=O 1683, S=O 1062; 270-MHz NMR (CDCl<sub>3</sub>)  $\delta$  7.43-7.17 (5 H, m), 6.74 (1 H, d, J = 12.1 Hz), 6.05 (1 H, d, J = 9.2 Hz), 5.85 (1 H, dd, J = 9.5, 14.2 Hz), 5.66 (1 H, ddd,J = 3.9, 9.8, 14.2 Hz, 5.62 (1 H, dd, J = 11.8, 12.1 Hz), 4.82 (1 H, s), 4.79 (1 H, s), 4.56 (1 H, dd, J = 9.2, 11.8 Hz), 4.07 (1 H, ddd, J = 2.1,3.0, 11.0 Hz), 3.87 (1 H, d, J = 8.9 Hz), 3.46 (1 H, dd, J = 3.2, 13.3 Hz), 2.85-2.63 (4 H, m), 2.68 (3 H, s), 2.49-2.38 (1 H, m), 2.40 (3 H, s), 2.27–2.23 (1 H, m), 2.25 (3 H), 2.21–2.09 (1 H, m), 1.40 (3 H, d, J = 7.4 Hz), 0.74 (9 H, s), 0.25 (3 H, d, J = 7.1 Hz), -0.07 (6 H, s). 30: oil; analytical TLC (silica gel F254), 2:1 EtOAc/hexane,  $R_f = 0.23$ ; 270-MHz NMR (CDCl<sub>3</sub>)  $\delta$  7.50-7.21 (5 H, m), 6.69 (1 H, m, dd, J = 7.4, 17.5 Hz), 6.30 (1 H, s), 6.15 (1 H, d, J = 17.5 Hz), 5.83 (1 H, dd, J = 9.8, 16.0 Hz), 5.42 (1 H, ddd, J = 7.4, 9.5, 16.0 Hz), 4.81 (1 H, s),4.79 (1 H, s), 4.124 (1 H, d, J = 7.4 Hz), 0.09 (1 H, dd, J = 9.2, 14.5 Hz), 3.82 (1 H, d, J = 8.3 Hz), 3.54 (1 H, d, J = 8.0 Hz), 3.43 (1 H, dd, J = 2.7, 13.6 Hz), 3.33-3.26 (1 H, m), 2.79-2.33 (4 H, m), 2.61 (3 H, m)H, s), 2.46 (3 H, s), 2.24 (3 H, s), 2.16-2.01 (1 H, m), 1.22 (3 H, d, J = 6.8 Hz), 0.74 (9 H, s), 0.30 (3 H, d, J = 6.5 Hz), -0.07 (6 H, s).

Allylic Sulfoxide 29 Rearrangement to Allylic Alcohol 31. A solution of 29 (1.8 mg, 2.6 µmol) and trimethylphosphite (Aldrich, freshly distilled from CaH<sub>2</sub>; 0.15 mL) and 1-propanol (EM Science, reagent grade; 0.05 mL) was heated at 100-105 °C. After ca. 5 h, additional trimethyl phosphite (0.15 mL) was added and heating continued 18 h longer. After the reaction was allowed to cool to room temperature, the volatiles were removed in vacuo. The residue was taken up in EtOAc (10 mL) and was washed with brine  $(2 \times 5 \text{ mL})$ . The combined aqueous layers were extracted with EtOAc  $(2 \times 5 \text{ mL})$ . The combined organic portions were dried (MgSO<sub>4</sub>) and filtered, and the solvents were removed in vacuo. The residue was passed through a flash silica gel plug (0.5 cm  $\times$  2 cm, 1:2 EtOAc/hexane) and further purified by ATLC (plate 5 cm × 10 cm, 1:2 EtOAc/hexane) to give rearranged allylic alcohol 31, 0.9 mg (53% yield): oil; analytical TLC (silica gel F254), 1:2 EtOAc/hexane,  $R_f = 0.36$ ; MS no peak match, parent, M - 57 (*tert*-butyl) = 592.2753, calcd = 592.2730, error = 3.4 ppm, formula =  $C_{37}H_{51}NO_7Si$ ; IR (neat, cm<sup>-1</sup>) O-O 3415, C=O 1745, C=O 1723, C=O 1705; 270-MHz NMR  $(CDCl_3) \delta 7.35-7.18 (5 H, m), 6.08 (1 H, dd, J = 0.9, 16.3 Hz), 5.71$ (1 H, dd, J 11.6, 14.8 Hz), 5.70 (1 H, dd, J = 0.9, 2.4 Hz), 5.29 (1 H, 1.4 Hz), 5.29 (1 Hz), 5ddd, J = 6.8, 8.3, 14.8 Hz), 5.09 (1 H, ddd, J = 2.4, 5.0, 14.8 Hz), 4.87 (1 H, s), 4.84 (1 H, dd, J = 3.3, 5.0 Hz), 4.81 (1 H, s), 3.97 (1 H, ddd,J = 1.5, 4.1, 10.3 Hz), 3.76 (1 H, d, J = 8.9 Hz), 3.27 (1 H, dd, J = 2.4, 12.1 Hz), 3.17 (1 H, ddq, J = 3.3, 8.9, 7.1 Hz), 2.88-2.82 (1 H, m),2.79 (1 H, dd, J = 8.6, 9.5 Hz), 2.59 (1 H, dd, J = 10.7, 12.2 Hz), 2.54-2.06 (4 H, m), 2.49 (3 H, s), 2.29 (3 H, s), 1.19 (3 H, d, J = 7.1 Hz), 0.76 (9 H, s), 0.31 (3 H, d, J = 6.8 Hz), -0.07 (6 H,s).

Deprotection of 31 to dl-C<sub>18</sub>-Desmethylcytochalasin D (32). To a solution of 31 (0.9 mg, 1.4  $\mu$ mol) in THF (0.1 mL) and MeOH (0.2 mL) at ca. -20 °C (CCl<sub>4</sub>/CO<sub>2</sub> bath) was added K<sub>2</sub>CO<sub>3</sub> (flame-dried in vacuo, 6.8 mg, 49  $\mu$ mol). The suspension was stirred at ca. -20 °C for 70 min

before it was partitioned between EtOAc (10 mL) and brine (5 mL). The layers were separated, and the organics were washed with brine (5 mL), and the combined aqueous layers were extracted with EtOAc (2  $\times$  5 mL). The organics were dried (MgSO<sub>4</sub>) and filtered, and the solvents were removed to give 0.7 mg of the deacylated amide as a slightly yellow oil that was used in the next step without further purification: oil; analytical TLC (silica gel F254), 1:1 EtOAc/hexane,  $R_f = 0.30$ ; 270-MHz NMR (CDCl<sub>3</sub>)  $\delta$  7.39–7.12 (5 H, m), 6.41 (1 H, ddd, J = 1.8, 3.9, 16.0 Hz), 5.63 (1 H, dd, J = 5.6, 15.4 Hz), 5.59 (1 H, d, J = 2.3 Hz), 5.46 (1 H, s), 5.34 (1 H, d, J = 5.3 Hz), 5.17 (1 H, ddd, J = 7.1, 7.7, 15.4 Hz), 5.15 (1 H, s), 5.05 (1 H, ddd, J = 4.4, 6.8, 16.0 Hz), 4.99 (1 H, s), 4.77-4.74 (1 H, m), 3.90 (1 H, d, J = 3.29 Hz), 3.29-3.19 (1 H, m), 2.92-2.73 (3 H, m), 2.61 (1 H, dd, J = 9.8, 13.3 Hz), 2.46-2.43 (1 H, m), 2.35-2.,26 (1 H, m), 2.23 (3 H, s), 2.13-1.96 (2 H, m), 1.15 (3 H, d, J = 7.1 Hz), 1.02 (3 H, d, J = 7.1 Hz), 0.81 (9 H, s), -0.03 (6 H, s).

The amide (0.7 mg) was dissolved in CH<sub>3</sub>CN (0.1 mL) and cooled to 0 °C then 48% (aqueous) HF was added (1 µL, 24 µmol) and the reaction was stirred for 100 min at 0 °C and then partitioned between EtOAc (5 mL) and brine (5 mL). The organics were washed with brine (5 mL), the combined aqueous layers were bakc-extracted with EtOAc  $(2 \times 5 \text{ mL})$ , and then the organic portion was dried (MgSO<sub>4</sub>) and filtered, and the solvents were evaporated. The residue was purified by

ATLC (5 cm  $\times$  10 cm, 3:1 EtOAc/hexane) to give *dl*-C<sub>18</sub>-desmethylcytochalasin D (32), 0.5 mg (ca. 70% for two steps): white crystals from acetone/hexane; mp 232-236 °C dec; analytical TLC (silica gel F254), 3:1 EtOAc/hexane,  $R_f = 0.41$ ; MS exact mass calcd for  $C_{29}H_{35}NO_6$ 493.2464, found 493.2472, error = 1.6 ppm; IR (neat, cm<sup>-1</sup>) N H and OH 3340, C=O 1740, C=O 1690, C=O 1685; 270-MHz NMR (CD-Cl<sub>3</sub>)  $\delta$  7.35-7.11 (5 H, m), 6.50 (1 H, dd, J = 1.8, 16.0 Hz), 5.77 (1 H, dd, J = 9.2, 15.4 Hz), 5.55 (1 H, dd, J = 2.4, 2.7 Hz), 5.51 (1 H, d, J= 0.9 Hz, 5.44 (1 H, s), 5.37 (1 H, ddd, J = 6.8, 7.4, 15.4 Hz), 5.10 (1 H, ddd, J = 2.7, 5.0, 16.0 Hz), 4.78-4.75 (1 H, m), 3.82 (1 H, dd, dd)J = 1.5, 10.9 Hz), 3.35-3.25 (2 H, m), 2.94 (1 H, dd, J = 3.88, 13.3 Hz), 2.80 (1 H, dd, J = 10.0, 11.0 Hz), 2.80–2.73 (1 H, m), 2.56 (1 H, dd, J = 9.8, 13.3 Hz, 2.38–2.28 (2 H, m), 2.21 (3 H, s), 2.19–2.05 (3 H, m), 1.97 (1 H, s), 1.15 (3 H, d, J = 7.1 Hz), 1.11 (3 H, d, J = 6.8 Hz).

Acknowledgment. This work was supported by the National Institutes of Health (Grant No. CA-17918).

Supplementary Material Available: Table of mass spectral fragmentation patterns for 32, CD, and ZE and details of the route to C<sub>16</sub>-epi-zygosporin E (5 pages). Ordering information is available on any current masthead page.

# Synthesis of Cyclobutanones by the Photolytic Reaction of Chromium Carbene Complexes with Olefins: Inter- and Intramolecular Reactions

## Björn C. Söderberg, Louis S. Hegedus,\* and Miguel A. Sierra

Contribution from the Department of Chemistry, Colorado State University, Fort Collins, Colorado 80523. Received October 5, 1989

Abstract: Cyclobutanones were synthesized in good yield and with a high degree of stereo- and regioselectivity by the photolytic reaction between a variety of chromium alkoxycarbene complexes and olefins. Bicyclic cyclobutanones were synthesized in good yield and with a high degree of stereo- and regioselectivity by the photolysis of chromium alkoxycarbenes having remote double bonds in the alkoxy group.

Photolysis reactions of heteroatom-stabilized (Fischer) chromium carbene complexes are becoming increasingly useful for the synthesis of novel organic compounds under exceptionally mild conditions. Thus photolysis, using visible light and a Pyrex reaction vessel, in solvents ranging from hexane through acetonitrile, of methoxycarbenes1 and aminocarbenes2 with imines produced  $\beta$ -lactams in excellent chemical yield. Use of optically active aminocarbene complexes<sup>3</sup> resulted in the production of optically active  $\beta$ -lactams in good yield and with high stereoselectivity. Photolysis of aminocarbene complexes in the presence of alcohols produced  $\alpha$ -amino acid esters.<sup>5</sup> During the development of these reactions it became clear that photolysis of chromium carbene complexes produced species that reacted as if they were ketenes, although no evidence for the generation of free ketenes was observed.<sup>6</sup> This suggested that other classes of reactions in which ketenes engaged should be examined.

Stereospecific [2 + 2] cycloaddition<sup>7</sup> reactions of ketenes and olefins to produce cyclobutanones8 have been extensively developed, although the use of electron-rich O- or N-containing ketenes is uncommon.9 In contrast, intramolecular versions of this process

Hegedus, L. S.; McGuire, M. A.; Schultze, L. M.; Yijun, C.; Anderson,
 P. J. Am. Chem. Soc. 1984, 106, 2680.
 Borel, C.; Hegedus, L. S.; Krebs, J.; Satoh, Y. J. Am. Chem. Soc. 1987,

<sup>109, 1101.</sup> 

<sup>(3)</sup> Inwinkelried, R.; Hegedus, L. S. Organometallics 1988, 7, 702.
(4) Hegedus, L. S.; Imwinkelried, R.; Alarid-Sargent, M.; Dvorak, D.; Satoh, Y. J. Am. Chem. Soc. 1990, 112, 1109.
(5) Hegedus, L. S.; Schwindt, M. A.; DeLombaert, S.; Imwinkelried, R. J. Am. Chem. Soc. 1990, 112, 2264.

<sup>(6) (</sup>a) Hegedus, L. S.; deWeck, G.; D'Andrea, S. J. Am. Chem. Soc. 1988, 110, 2122. (b) Cyclobutanones have also been produced by intramolecular trapping of chromium-vinyl ketene complexes, produced by the thermal re-action of chromium-carbene complexes with enynes, see: Wulff, W. D.; Kaisler, R. W. Organometallics 1985, 4, 1461. Korkowski, P. F.; Hoye, T. R.; Rydberg, D. B. J. Am. Chem. Soc. 1988, 110, 2676. (c) Reaction of (phenyl)(methoxy)carbene-chromium complexes with enamides under 150 atm of CO pressure also produced cyclobutanones: Dorrer, B.; Fischer, E. O. Chem. Ber. 1974, 107, 2683. (d) Treatment of (ethoxy)(triphenylsily)-tungsten carbenes under 50 atm of CO produced the free ketene which was isolated and/or allowed to react with enol ethers to form cyclobutanones: Schubert, U.; Uron, J.; Hörnig, H. J. Organomet. Chem. 1988, 355, 243. Uron, J.; Schubert, U. Ibid. 1989, 373, 203.

<sup>Uron, J.; Schubert, U. Ibid. 1989, 373, 203.
(7) (a) Ghosez, L.; O'Donnell, J. M. Pericyclic Reactions; Marchand, A. P., Lehr, R. E., Eds.; Academic Press: New York, 1977; Vol. II, pp 79-140.
(b) Brady, W. T. Tetrahedron 1981, 37, 2949.
(c) Brady, W. T. The Chemistry of Ketenes, Allenes, and Related Compounds; Patai, S., Ed.; Interscience Publications: New York, 1980; pp 278-308.
(8) Bellus, D.; Ernst, B. Angew. Chem., Int. Ed. Engl. 1988, 27, 797.
(9) (a) Reichen, W. Chem. Rev. 1978, 78, 568.
(b) Brady, W. T.; Gu, Y.</sup> 

Q. J. Org. Chem. 1989, 54, 2834.