

Total Synthesis of Chloropeptin II (Complestatin) and Chloropeptin I

Joie Garfunkle, F. Scott Kimball, John D. Trzupek, Shinobu Takizawa, Hiroyuki Shimamura, Masaki Tomishima, and Dale L. Boger*

Department of Chemistry and The Skaggs Institute for Chemical Biology, The Scripps Research Institute, 10550 North Torrey Pines Road, La Jolla, California 92037

Received August 25, 2009; E-mail: boger@scripps.edu

Complestatin (**1**, chloropeptin II) was first disclosed in 1980 as an inhibitor of the alternate pathway of human complement,¹ Figure 1. Nine years later, both its original isolation² from *Streptomyces lavendulae* at Sankyo and additional details of its complement activity were reported along with its structure elucidation by Seto³ that defined its connectivity and partial stereochemistry. Shortly thereafter, the first report of its activity against HIV infectivity and its cytopathic effects were disclosed.⁴ Five years later, Omura reported the isolation of both chloropeptin I (**2**) and chloropeptin II (**1**) from *Streptomyces* sp. WK-3419 as inhibitors of HIV gp120-CD4 binding, determined that chloropeptin II and complestatin are identical, and established their partial stereochemistry.⁵ A more detailed analysis of their NMR data provided the full structural and stereochemical assignment for chloropeptin I (**2**) including the axial atropisomer chirality.⁶ A remarkable acid-catalyzed rearrangement (TFA, 50 °C, >90%) of chloropeptin II (**1**, complestatin) to the less strained chloropeptin I (**2**) that proceeds with retention of the atropisomer stereochemistry subsequently established the full stereochemical assignments for **1**.^{1c,7} These later studies were conducted in the course of the additional isolations of the natural products at Merck⁸ and Schering-Plough,⁹ with the latter establishing that chloropeptin I (**2**) is an authentic natural product and not an acid-catalyzed artifact derived from chloropeptin II (**1**).

As a result of the challenging structural features and complexity of **1** and **2**, rivaling that of the glycopeptide antibiotics (e.g., vancomycin), combined with their equally important HIV activity derived through a unique site of action, they have attracted considerable interest. Although structurally similar to the glycopeptide antibiotics, one of the characteristic biaryl ether linkages is replaced with a biaryl linkage to C6 or C7 of a (*R*)-tryptophan indole embedded in the macrocyclic core adopting a single atropisomer stereochemistry (*R*) that is not capable of thermal interconversion. Snapper and Hoveyda reported the first, and to date only, total synthesis of a member of this unique class of natural products, chloropeptin I (**2**), confirming the structural and stereochemical assignments.¹⁰ Their approach, enlisting a late stage intramolecular biaryl Stille coupling for closure of the right-hand macrocyclic ring system (38–42%, three steps) provided exclusively the natural (*R*)-atropisomer of **2** as a single diastereomer when conducted on substrate containing the left-hand macrocycle. Remarkably, they later found that extending this approach to chloropeptin II (**1**), using an analogous late stage intramolecular biaryl Suzuki coupling, provided exclusively (63%) the unnatural (*S*)-atropisomer (isocomplestatin).¹¹

Complementary to these and related ongoing efforts,¹² herein we report the first total synthesis of chloropeptin II (**1**, complestatin), the more strained and challenging of the natural products. Key to the approach is the use of an intramolecular Larock indole synthesis¹³ for the initial macrocyclization, adopting conditions that permit utilization of a 2-bromoaniline¹⁴ and incorporating a removable terminal alkyne substituent (–SiEt₃) that sterically dictates the indole cyclization regioselectivity,^{13–15} Figure 1.

Not only did this key reaction provide the fully functionalized right-hand ring system of **1** in superb conversion (89%) and good




Figure 1. Natural products and key retrosynthetic disconnections.

atropiastereoselectivity (4:1), but it also represents the first reported example of what may prove to be a useful Larock macrocyclization strategy. Subsequent introduction of the left-hand ring system by enlisting an aromatic nucleophilic substitution reaction for ring closure with biaryl ether formation completed the assemblage of the core bicyclic structure of **1** and represents a macrocyclization order complementary to that of Snapper and Hoveyda. Intrinsic in the design of the approach and by virtue of the single-step acid-catalyzed conversion of **1** to chloropeptin I (**2**), the route also provides a total synthesis of **2**.

Although **1** is composed of seven amino acid subunits, each is extensively modified, four are racemization prone phenylglycines, five are incorporated with the unnatural D-configuration, and one is incorporated as a *N*-methyl derivative. Four of these (A, C, E, and G) are readily accessible from commercially available precursors,¹⁶ whereas the remaining three required more significant synthetic operations key to implementation of our approach. The B subunit **5**, substituted to accommodate its use in an intramolecular aromatic nucleophilic substitution reaction for macrocyclic biaryl ether formation, simply required *N*-methylation of the Boc derivative of (*S*)-4-fluoro-3-nitrophenylalanine prepared by a four-step synthesis previously utilized to access its enantiomer,¹⁷ Scheme 1. The F subunit **6** bearing the TES-substituted alkyne required for the Larock indole annulation was accessed by diastereoselective alkylation (dr >99:1) of (*S*)-**8** with the propargylic diphenylphosphate **7**¹⁵ following a protocol detailed

treatment with Boc₂O providing **4**. Notably, the indole *N*-acetamide was unaffected by this treatment, and the intrinsically strained ring system did not undergo rearrangement to the more stable C7 (vs C6) biaryl indole linkage. The assigned structure and stereochemistry were reaffirmed, through a full spectroscopic characterization of **4**, not only with observation of key NOEs¹⁶ and the diagnostic chemical shifts of the Trp α -CH (THF-*d*₈: δ 3.86, app t) and the diastereotopic Trp β -CH₂ (δ 2.82, d and 3.43, app t) as well as their multiplicity¹¹ but also simply through the indole coupling pattern where C7-H remains a singlet (δ 8.30, s) while C4-H and C5-H appear as coupled doublets in **4**.

This set the stage for introduction of the left-hand ring system. Coupling (EDCI, HOAt, DMF/CH₂Cl₂ 1:3, -5 °C, 6 h, 59%) of **4** with the tripeptide **3**, prepared by the sequential couplings and N-terminus deprotections of (*R*)-H₂N-Hpg-OMe (**24**)¹⁶ with **5** (PyBOP, 80%; 4 N HCl, dioxane) and (*R*)-FmocHN-3,5-Cl₂Hpg-OH¹⁶ (**17**, DEPBT, NaHCO₃, THF, 0 °C, 24 h, 83%, 9:1 dr; Bu₄NF,²¹ THF, 0–23 °C, <1 h),¹⁶ provided **25**. Macrocyclization²² of **25** to provide **26** as predominantly a single atropisomer of an inconsequential mixture of atropisomers was accomplished upon treatment with K₂CO₃ in THF (0.5 mM, 60 °C, 48–64 h) in the presence of 18-c-6 and 4 Å MS in conversions as high as 81%, provided rigorous anhydrous conditions were maintained to prevent competitive methyl ester hydrolysis. Two-step removal of the activating nitro group (H₂, Ra-Ni, MeOH, 0–23 °C, 6 h, 87%; *t*-BuONO, H₃PO₂, THF, 0 °C, 3 h, 72%)²³ afforded **28**. Boc deprotection (4 N HCl, dioxane, 23 °C, 1–3 h) and coupling of the amine with 2-(3,5-dichloro-4-hydroxyphenyl)-2-oxoacetic acid (**29**,^{10,16} EDCI, HOAt, DMF/CH₂Cl₂ 1:5, 0 °C, 2 h, 55%) provided the penultimate precursor **30**. Deprotection of **30** to provide **1** was accomplished with LiOH (THF/H₂O, 0 °C, 3 h, 60%) in a reaction where the indole *N*-acetyl group was removed faster (<30 min) than the methyl ester hydrolysis. Finally, and although we did not conduct the reaction on a preparative scale providing an isolated yield, the clean acid-catalyzed conversion of **1** to **2** was conducted on a small scale with both synthetic and authentic **1** and monitored by LCMS. The two samples behaved in the same manner providing only **2** and was most conveniently conducted with 50% TFA/H₂O at 50 °C progressing at a rate that is easily monitored (5 h, vs <5–15 min with neat TFA at 50 °C).²⁴

Continued efforts on the optimization and definition of the scope of the Larock macrocyclization reaction, the examination of the reverse macrocyclization order, and the extension of the approach to additional natural products and their key analogues are in progress and will be disclosed in due course.

Acknowledgment. In memory of David S. Lewy. We gratefully acknowledge the financial support of the National Institutes of Health (CA041101) and the Skaggs Institute for Chemical Biology. We wish to thank Dr. S. B. Singh (Merck) for authentic samples of **1** and **2**. We wish to especially thank M. Tichenor for introducing improvements to the Larock macrocyclization and subsequent elaboration to the DEF ring system; Dr. J. Cottell for initiating studies on the ABCD ring system; and D. S. Lewy, Drs. A. Pichota, L. Resnick, and W. Han for exploration of early stage routes to the DEF ring system in which the syntheses of the amino acid subunits were first developed. J.G. and J.D.T. were Skaggs fellows.

Supporting Information Available: Full experimental details are provided. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) (a) Kaneko, I.; Fearon, D. T.; Austen, K. F. *J. Immunol.* **1980**, *124*, 1194–1198. (b) Tachikawa, K.; Hasumi, K.; Endo, A. *Thromb. Haemost.* **1997**, *77*, 137–142. (c) Tachikawa, K.; Hasumi, K.; Endo, A. *Thrombosis Res.* **1997**, *87*, 571–576.
- (2) Kaneko, I.; Kamoshida, K.; Takahashi, S. *J. Antibiot.* **1989**, *42*, 236–241.
- (3) Seto, H.; Fujioka, T.; Furihata, K.; Kaneko, I.; Takahashi, S. *Tetrahedron Lett.* **1989**, *30*, 4987–4990.
- (4) Momota, K.; Kaneko, I.; Kimura, S.; Mitamura, K.; Shimada, K. *Biochem. Biophys. Res. Commun.* **1991**, *179*, 243–250.
- (5) (a) Matsuzaki, K.; Ikeda, H.; Ogino, T.; Matsumoto, A.; Woodruff, H. B.; Tanaka, H.; Omura, S. *J. Antibiot.* **1994**, *47*, 1173–1174. (b) Tanaka, H.; Matsuzaki, K.; Nakashima, H.; Ogino, T.; Matsumoto, A.; Ikeda, H.; Woodruff, H. B.; Omura, S. *J. Antibiot.* **1997**, *50*, 58–65. (c) Matsuzaki, K.; Ogino, T.; Sunazuka, T.; Tanaka, H.; Omura, S. *J. Antibiot.* **1997**, *50*, 66–69.
- (6) Gouda, H.; Matsuzaki, K.; Tanaka, H.; Hirono, S.; Omura, S.; McCauley, J. A.; Sprengeler, P. A.; Furst, G. T.; Smith, A. B., III. *J. Am. Chem. Soc.* **1996**, *118*, 13087–13088.
- (7) (a) Jayasuriya, H.; Salituro, G. M.; Smith, S. K.; Heck, J. V.; Gould, S. J.; Singh, S. B.; Homnick, C. F.; Holloway, M. K.; Pitzenger, S. M.; Patane, M. A. *Tetrahedron Lett.* **1998**, *39*, 2247–2248. (b) Hegde, V. R.; Dai, P.; Patel, M.; Gullo, V. P. *Tetrahedron Lett.* **1998**, *39*, 5683–5684.
- (8) (a) Singh, S. B.; Jayasuriya, H.; Salituro, G. M.; Zink, D. L.; Shafiee, A.; Heimbuch, B.; Silverman, K. C.; Lingham, R. B.; Genilloud, O.; Teran, A.; Vilella, D.; Felock, P.; Hazuda, D. J. *Nat. Prod.* **2001**, *64*, 874–882. (b) Singh, S. B.; Jayasuriya, H.; Hazuda, D. L.; Felock, P.; Homnick, C. F.; Sardana, M.; Patane, M. A. *Tetrahedron Lett.* **1998**, *39*, 8769–8770.
- (9) (a) Hegde, V. R.; Puar, M. S.; Dai, P. D.; Patel, M.; Gullo, V. P.; Chan, T.-M.; Silver, J.; Pramanik, B. N.; Jenh, C.-H. *Bioorg. Med. Chem. Lett.* **2003**, *13*, 573–575. (b) Hegde, V. R.; Puar, M. S.; Dai, P.; Patel, M.; Gullo, V. P.; Pramanik, B. N.; Jenh, C.-H. *Tetrahedron Lett.* **2002**, *43*, 5339–5341.
- (10) Deng, H.; Jung, J.-K.; Liu, T.; Kuntz, K. W.; Snapper, M. L.; Hoveyda, A. H. *J. Am. Chem. Soc.* **2003**, *125*, 9032–9034.
- (11) Shinohara, T.; Deng, H.; Snapper, M. L.; Hoveyda, A. H. *J. Am. Chem. Soc.* **2005**, *127*, 7334–7336.
- (12) (a) Jia, Y.; Bois-Choussy, M.; Zhu, J. *Angew. Chem., Int. Ed.* **2008**, *47*, 4167–4172. (b) Jia, Y.; Bois-Choussy, M.; Zhu, J. *Org. Lett.* **2007**, *9*, 2401–2404. (c) Yamada, Y.; Akiba, A.; Arima, S.; Okada, C.; Yoshida, K.; Itou, F.; Kai, T.; Satou, T.; Takeda, K.; Harigaya, Y. *Chem. Pharm. Bull. Jpn.* **2005**, *53*, 1277–1290. (d) Smith, A. B., III; Chruma, J. J.; Han, Q.; Barbosa, J. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 1697–1702. (e) Elder, A. M.; Rich, D. H. *Org. Lett.* **1999**, *1*, 1443–1446. (f) Roussi, G.; Zamora, E. G.; Carbonnelle, A.-C.; Beugelmans, R. *Heterocycles* **1999**, *51*, 2041–2063. (g) Kai, T.; Kajimoto, N.; Konda, Y.; Harigaya, Y.; Takayanagi, H. *Tetrahedron Lett.* **1999**, *40*, 6289–6292. (h) Gurjar, M. K.; Tripathy, N. K. *Tetrahedron Lett.* **1997**, *38*, 2163–2166.
- (13) (a) Larock, R. C.; Yum, E. K. *J. Am. Chem. Soc.* **1991**, *113*, 6689–6690. (b) Larock, R. C.; Yum, E. K.; Refvik, M. D. *J. Org. Chem.* **1998**, *63*, 7652–7662.
- (14) Shen, M.; Li, G.; Lu, B. Z.; Hossain, A.; Roschhangar, F.; Farina, V.; Senanayake, C. H. *Org. Lett.* **2004**, *6*, 4129–4132.
- (15) (a) Ma, C.; Liu, X.; Li, X.; Flippen-Anderson, J.; Yu, S.; Cook, J. M. *J. Org. Chem.* **2001**, *66*, 4525–4542. See also: (b) Jeschke, T.; Wensbo, D.; Annby, U.; Gronowitz, S.; Cohen, L. A. *Tetrahedron Lett.* **1993**, *34*, 6471–6474.
- (16) Details are provided in Supporting Information.
- (17) (a) Boger, D. L.; Kim, S. H.; Mori, Y.; Weng, J.-H.; Rogel, O.; Castle, S. L.; McAtee, J. J. *J. Am. Chem. Soc.* **2001**, *123*, 1862–1871. (b) Boger, D. L.; Kim, S. H.; Miyazaki, S.; Strittmatter, H.; Weng, J.-H.; Mori, Y.; Rogel, O.; Castle, S. L.; McAtee, J. J. *J. Am. Chem. Soc.* **2000**, *122*, 7416–7417.
- (18) Reddy, K. L.; Sharpless, K. B. *J. Am. Chem. Soc.* **1998**, *120*, 1207–1217.
- (19) Miyauro, N.; Suzuki, A. *Chem. Rev.* **1995**, *95*, 2457–2483.
- (20) Trzupke, J. D. Ph.D. Dissertation, 2006.
- (21) Jiang, W.; Wanner, J.; Lee, R. J.; Bounaud, P.-Y.; Boger, D. L. *J. Am. Chem. Soc.* **2002**, *124*, 5288–5290 and **2003**, *125*, 1877–1887.
- (22) (a) Boger, D. L. *Med. Res. Rev.* **2001**, *21*, 356–381. (b) Boger, D. L.; Miyazaki, S.; Kim, S. H.; Wu, J. H.; Loiseleur, O.; Castle, S. L. *J. Am. Chem. Soc.* **1999**, *121*, 3226–3227 and 10004–10011. (c) Boger, D. L.; Castle, S. L.; Miyazaki, S.; Wu, J. H.; Beres, R. T.; Loiseleur, O. *J. Org. Chem.* **1999**, *64*, 70–80. (d) Boger, D. L.; Borzilleri, R. M.; Nukui, S.; Beres, R. T. *J. Org. Chem.* **1997**, *62*, 4721–4736. (e) Boger, D. L.; Borzilleri, R. M.; Nukui, S. *Bioorg. Med. Chem. Lett.* **1995**, *5*, 3091–3096. (f) Boger, D. L.; Borzilleri, R. M. *Bioorg. Med. Chem. Lett.* **1995**, *5*, 1187–1190. (g) Crowley, B. M.; Boger, D. L. *J. Am. Chem. Soc.* **2006**, *128*, 2885–2892.
- (23) Crowley, B. M.; Mori, Y.; McComas, C. C.; Tang, D.; Boger, D. L. *J. Am. Chem. Soc.* **2004**, *126*, 4310–4317.
- (24) Abbreviations: DEPBT, 3-(diethoxyphosphoryloxy)-1,2,3-benzotriazin-4-one; DtBPT, 1,1'-bis(di-*tert*-butylphosphino)ferrocene; EDCI, 1-[3-(dimethyl-amino)propyl]-3-ethylcarbodiimide hydrochloride; HOAt, 1-hydroxy-7-azabenzotriazole; PyBOP, (benzotriazol-1-yl)tripyrrolidinophosphonium hexafluorophosphate.

JA907193B