

Tetrahedron Letters 39 (1998) 8373-8376

TETRAHEDRON LETTERS

A Convenient Synthesis of Racemic 2'-Deoxy Carbocyclic Thymidines Lacking the 5'-Methylene Group

Jinglan Zhou and Philip B. Shevlin*

Department of Chemistry, Auburn University, Auburn, AL 36830

Received 11 August 1998; accepted 10 September 1998

Abstract. A convenient synthesis of racemic 2'-deoxy carbocyclic thymidines lacking the 5'-methylene group is reported. Thus, appropriately protected nucleic acid bases are coupled to 3-cyclopentenol, 1, via the Mitsunobu reaction. Epoxidation of the protected thymine substituted cyclopentene gives the corresponding syn and anti-epoxides in a 4:1 ratio. Ring opening of the syn-epoxide by a variety of nucleophiles leads to the racemic 2'-deoxy carbocyclic thymidines lacking the 5'-methylene group. © 1998 Elsevier Science Ltd. All rights reserved.

We wish to report a convenient and potentially general method for synthesis of 2'-deoxy carbocyclic nucleoside analogs lacking the 5'-methylene group.¹ As outlined in eq 1, our synthetic strategy involves an early stage attachment of an appropriately protected nucleic acid base to 3-cyclopentenol, 1, followed by epoxidation. Ring opening of the epoxides by various types of nucleophiles leads to the desired 2'-deoxy carbocyclic nucleosides lacking the 5'methylene with a variety of functional groups at the 3'- position. Direct coupling





of the nucleic acid bases with cyclopentenol under Mitsunobu conditions and minimum use of protecting groups make the synthetic route very short, simple and economical. Reversing the order of Mitsunobu coupling and epoxidation allows the synthesis of the anti base substituted epoxide.

An ideal starting material, 3-cyclopentenol, 1, can be prepared from cyclopentadiene in large scale by a two-step epoxidation-reduction sequence in an overall yield of 35%.² The Mitsunobu reaction of 1 with protected nucleic acid bases was carried out in dioxane in the presence of triphenylphosphine and diethyl azodicarboxylate (DEAD) at room temperature to give **2a-d**. Thymine or uracil were protected 0040-4039/98/\$ - see front matter © 1998 Elsevier Science Ltd. All rights reserved. *PII:* S0040-4039(98)01929-7

as 3-benzoylthymine (p-T) or 3-benzoyluracil (p-U) which were prepared from thymine or uracil by a two step benzoylation-hydrolysis.³ Adenine and guanine were protected as 6-Isobutyrylaminopurine (p-A)⁴ and N₂acetyl-O₆-(2-(p-nitrophenyl)-ethyl)guanine (p-G)⁵ respectively.



Oxidation of 2a using *m*-CPBA in methylene chloride gave a 75% yield of syn and anti-3a, in a 4 to 1 ratio (eq 2), which were easily separable by column chromatography.⁶ The assignment of stereochemistry to syn and anti-3a relies on their independent synthesis. Thus the known syn-4 was prepared in 70% yield by hydroxy directed epoxidation of 1 with *t*-butyl hydroperoxide in benzene in the presence of a catalytic amount of VO(acac)₂.⁷ Reaction of syn-4 with **p**-T under Mitsunobu conditions (eq 3) gave a compound in 28% isolated yield which was identical to anti-3a. In order to further confirm the configuration of syn and anti-3a, 1 was oxidized with m-CPBA in cyclopentane to give the known⁸ syn and anti-4 in a 3:1 ratio. Anti-4 was separated by chromatography and allowed to undergo Mitsunobu coupling with **p**-T to generate a compound identical to syn-3a in 67% yield (eq 4).



In order to test the synthetic utility of **3a** in generating 3,4-disubstituted carbocyclic nucleoside analogs, several ring opening reactions were carried out. Thus, synthesis of (\pm) -3-benzoyl- $((3'\alpha, 4'\beta)$ -bishydroxy-

cyclopentyl)-1H-thymine, 5, was accomplished by treating syn or anti-3a with dilute sulfuric acid in THF.^{1b} Ring opening of syn-3a with sodium azide in dioxane-H₂O under reflux was accompanied by simultaneous removal of 3-benzoyl group to give (\pm) - $(3'-\alpha$ -azido-4' β -hydroxy cyclopentyl)-1H-thymine 6 in an isolated yield of 71%.⁹ Treatment of syn-3a with PhSH/PhSNa in DMF at room temperature¹⁰ gave the protected thymine, (\pm) -3-benzoyl-3'- α -phenylthio-4' β -hydroxycyclopentyl)-1H-thymine, 7, and the deprotected (\pm) -3' α -phenylthio-4' β - hydroxycyclopentyl)-1H-thymine, 8 in a 1:2.8 ratio.



As illustrated in eqs 3 and 4, these procedures permit the preparation of either syn or anti-3a as the major product depending upon the order of Mitsunobu coupling and epoxidation. The fact that syn-4 is the only product of stereospecific epoxidation⁷ allows the synthesis of anti-3a. The preparation of anti-3b-d via this route should also be straightforward. Although amine oxide formation may complicate the preparation of syn-3c,d, syn-3b should be available via the process in eq 2. These synthetic transformations illustrate the potential versatility of these compounds in leading to carbocyclic nucleoside analogs.

References and Notes:

(a) Patil, S. D.; Koga, M.; Schneller, S. W. J. Med. Chem. 1992, 35, 2191-2195. (b) Kim, C. U.; Luh, B. Y.;
Martin, J. C. J. Org, Chem. 1991, 56, 2642-2647. (c) Koga, M.; Schneller, S. W. Tetrahedron Lett. 1990, 31, 5861-5864. (d) For a discussion of furan analogs see: Elliott, R. D.; Rener, G. A.; Riordan, J. M.; Secrist, J. A. III; Parker, W. B.; Montgomery, J. A. J. Med. Chem. 1994, 37, 739-744.

- 2. Crandall, J. K.; Banks, D. B.; Colyer, R. A.; Watkins, R. J.; Arrington, J. P. J Org. Chem. 1968, 33, 423-425.
- 3. Cruickshank, K. A.; Jiricny, J.; Reese, C. B. Tetrahedron Lett. 1984, 25, 681-684.
- 4. Zhou, J. L.; Shevlin, P. B. Tetrahedron Lett. 1997, 38, 4037-4038.
- 5. Zhou, J. L. The Synthesis of DNA Analogs as Potential Antisense Agents, Auburn University, 1997.
- 6. Selected spectroscopic data for 2a-d, syn and anti-3a:

2a: ¹H NMR (CDCl₃, 250 Mhz) δ 1.94 (s, 3H), 2.41 (dd, J = 3.0, 16.4 Hz, 2H), 2.92 (dd, J = 8.9, 16.7 Hz, 2H), 5.32-5.42 (m, 1H), 5.87 (s, 2H), 7.05 (s, 1H), 7.49 (t, J = 7.8 Hz, 2H), 7.62-7.67 (m, 1H), 7.91-7.95 (m, 2H); ¹³C NMR (CDCl₃, 62.9 MHz) δ 12.5, 39.7, 52.4, 111.6, 129.1, 129.5, 130.5, 131.7, 135.0. 136.4, 149.8, 162.9, 169.4; HRMS calcd for C₁₇H₁₇N₂O₃ 297.1239, found 297.1242.

2b: ¹H NMR (CDCl₃, 300 Mhz) δ 2.42 (dd, J = 3.0, 16.8 Hz, 2H), 2.95 (dd, J = 8.7, 16.8 Hz, 2H), 5.32-5.38 (m, 1H), 5.81 (d, J = 7.9 Hz, 1H), 5.87 (s, 2H), 7.25 (d, J = 7.9 Hz, 1H), 7.51 (t, J = 7.6 Hz, 2H), 7.63-7.68 (m, 1H), 7.93-7.96 (m, 2H); ¹³C NMR (CDCl₃, 75.5 MHz) δ 39.4, 52.9, 102.5, 129.0, 129.2, 130.3, 131.3, 135.0, 140.9, 149.6, 162.0, 169.1; MS *m/z* 283 (M+1, 100); HRMS calcd for C₁₆H₁₅N₂O₃ 283.1083, found 283.1078. **2c**: ¹H NMR (CDCl₃, 250 MHz) δ 1.32 (d, J = 9.0 Hz, 6H), 2.65 (dd, J = 3.3, 15.7 Hz, 2H), 3.04 (dd, J = 8.0, 16.0 Hz, 2H), 3.19-3.30 (m, 1H), 5.37-5.46 (m, 1H), 5.95 (s, 2H), 8.05 (s, 1H), 8.74 (s, 1H), 9.13 (br s, 1H); ¹³C NMR (CDCl₃, 62.9 MHz) δ 19.4, 36.1, 40.5, 53.1, 122.3, 129.2, 141.0, 149.5, 151.5, 152.4, 176.6; HRMS calcd for C₁₄H₁₈N₅O 272.1511, found 272.1520.

2d: ¹H NMR (CDCl₃, 300 MHz) δ 2.58 (s, 3H), 2.65 (dd, J = 3.6, 15.6 Hz, 2H), 3.00 (dd, J = 8.1,15.9 Hz, 2H), 3.31 (t, J = 6.9 Hz, 2H), 4.78 (t, J = 6.9 Hz, 2H), 5.19-5.28 (m, 1H), 5.92 (s, 2H), 7.49 (d, J = 8.7 Hz, 2H), 7.83 (br s, 2H), 8.16 (d, J = 8.7 Hz, 2H); ¹³C NMR (CDCl₃, 75.5 MHz) δ 25.0, 34.8, 39.9, 52.9, 66.5, 117.7, 123.6, 128.9, 129.9, 139.9, 145.7, 146.8, 151.8, 152.7, 160.5, 171.2; HRMS calcd for C₂₀H₂₁N₆O₄ 409.1624, found 409.1643.

Syn-3a: Yield 60%; ¹H NMR (CDCl₃, 300 MHz) δ 1.98 (d, J = 1.2 Hz, 3H), 2.12 (dd, J = 1.8, 16.2 Hz, 2H), 2.49 (dd, J = 10.2, 16.2 Hz, 2H), 3.64 (s, 2H), 5.36-5.43 (m, 1H), 7.48 (t, J = 7.5 Hz, 2H), 7.60-7.65 (m, 1H), 7.76 (d, J = 1.2 Hz, 1H), 7.89-7.92 (m, 2H); ¹³C NMR (CDCl₃, 75.5 MHz) δ 12.5, 35.0, 52.8, 58.3, 110.9, 129.1, 130.4, 131.7, 134.9, 138.3, 150.5, 162.8, 169.4; HRMS calcd for C₁₇H₁₇N₂O₄ 313.1188, found 313.1186. **Anti-3a**: Yield 15%; ¹H NMR (CDCl₃, 300 MHz) δ 1.96 (d, J = 1.2 Hz, 3H), 2.11 (dd, J = 9.3, 14.1 Hz, 2H), 2.43 (dd, J = 7.8, 13.8 Hz, 2H), 3.60 (s, 2H), 4.42-4.53 (m, 1H), 7.04 (d, J = 1.2 Hz, 1H), 7.50 (t, J = 7.5 Hz, 2H), 7.63-7.66 (m, 1H), 7.90-7.93 (m, 2H); ¹³C NMR (CDCl₃, 75.5 MHz) δ 12.3, 31.3, 54.4, 55.4, 111.1, 129.2, 130.4, 131.5, 135.2, 138.6, 149.7, 162.8, 169.3; HRMS calcd for C₁₇H₁₇N₂O₄ 313.1188, found 313.1185. 7. Asami, M. *Bull. Chem. Soc. Jpn.* **1990**, *63*, 1402-1408.

8. Derby, A. C; Henbest, H. B.; McClenagham, I. Chem. Ind. (London) 1962, 462; Hornback, J. M. J. Org. Chem. 1973, 38, 4122; David, F. J. Org. Chem. 1981, 46, 3512-3519.

9. Creger, P. L. J. Org. Chem. 1972, 37, 1907-1918.

10. Strekoroski, L.; Battiste, M. A. Tetrahedron Lett. 1981, 22, 279-282.