


Dearomatized BIAN Alkaline-Earth Alkyl Catalysts for the Intramolecular Hydroamination of Hindered Aminoalkenes

Merle Arrowsmith, Michael S. Hill,* and Gabriele Kociok-Köhn

Department of Chemistry, University of Bath, Claverton Down, Bath BA2 7AY, U.K.


Supporting Information

ABSTRACT: Reaction of a sterically encumbered bis(imino)acenaphthene (dipp-BIAN) with either potassium alkyl or the heavier alkaline-earth dialkyl $[M\{CH(SiMe_3)_2\}_2(THF)_2]$ ($M = Mg, Ca, Sr$) reagents results in dearomatization of the aromatic ligand. The heteroleptic alkaline-earth alkyl species show enhanced stability toward Schlenk-type redistribution but undergo solution exchange when the bis(trimethylsilyl)methyl substituent is replaced by an anionic ligand of lower overall steric demands. In contrast, analogous reactions performed with $[Ba\{CH(SiMe_3)_2\}_2(THF)_2]$ evidenced facile solution redistribution and resulted in an unusual C–C coupling reaction which is suggested to result from a sterically induced reductive process. An assessment of the Mg, Ca, and Sr alkyl compounds as precatalysts for the intramolecular hydroamination of aminoalkenes evidenced enhanced reactivity, which is ascribed to the greater solution stability of the catalytically active species. Most notably the calcium species may even be applied to the high-yielding cyclization of substrates bearing alkyl substitution at either of the alkenyl positions.



INTRODUCTION

In the catalogue of readily available coordinating α -diimines, bis(imino)acenaphthene (BIAN) ligands such as **I** have gained some prominence over the last two decades. Although first reported in the 1960s,¹ the use of this class of ligand only became popular in the late 1990s.



The ability of neutral BIAN ligands to act as strong and rigid chelators for metals in a variety of oxidation states has given rise to a rich transition-metal coordination and catalytic chemistry.² In particular, Elsevier and co-workers have reported the use of palladium and platinum aryl-BIAN complexes for catalytic C–C bond formation, hydrogenation, and hydrosilylation reactions,³ while Brookhart⁴ and Coates⁵ have demonstrated the efficiency of cationic nickel(II), palladium(II), and platinum(II) aryl-BIAN precatalysts in the polymerization of olefins. Because of the presence of the naphthalene backbone,

BIAN ligands may also act as electron acceptors in the presence of strongly reducing metals. For example the addition of 1, 2, 3 or 4 equiv of elemental sodium to **I** respectively yields the mono-, di-, tri-, and tetraanionic salts $[\{I\}^{n-}\{Na_n(S)_m\}^{n+}]$ ($n = 1-4$, $S = Et_2O, THF$).⁶ This chemistry, extended to *s*- and *p*-block metals⁷ and, more recently, to *f*-block elements,⁸ has been extensively explored by the group of Fedushkin.


The redox properties of the BIAN ligand have also led to some unusual reactivity for main-group and, of most relevance to the current work, group 2 complexes. The magnesium species $[\{I\}Mg(THF)_3]$ (**II**), for example, has been shown to act as a reducing agent for nonenolizable ketones via single-electron transfer to form the corresponding magnesium alkoxide radical anion complex.⁹ Oxidation of complex **II** by halogenated tin, copper, mercury, or silicon reagents led to the isolation of radical-anion magnesium halides, such as compound **III**.¹⁰ Furthermore, compound **II** and its strontium and barium analogues have also been shown to deprotonate acidic substrates such as enolizable ketones,¹¹ phenylacetylene,¹² and α -acidic nitriles¹³ to give the corresponding enolate, acetylides, and keteniminate group 2 complexes of the protonated monoanionic amine-amido ligand **I-H**⁻. Alkylation of the BIAN imine carbon has also been observed in the reaction between **II** and ethyl halides, which resulted in the formation of complex **IV** via ethylation of the carbon adjacent to the amide functionality.¹⁴ The mechanism proposed by the authors for the formation of this compound involves a single-electron transfer from **II** to the ethyl halide, yielding the

Received: September 26, 2013

Published: December 11, 2013

intermediate radical anion magnesium halide complex **III**, and Et^\bullet , followed by radical attack of Et^\bullet to form complex **IV**. A similar alkyl transfer has also been observed in the reaction between ${}^n\text{BuLi}$ or MR_3 (M = group 13 metal, R = alkyl substituent) and **I** in noncoordinating solvents, leading to the formation of a imine-amido complex bearing an alkyl substituent on the carbon adjacent to the amide functionality.¹⁵ More recently Cowley and co-workers also reported the double alkylation of the acenaphthene backbone of **I** in the C^4 and C^5 positions by *tert*-butyllithium, via a radical dearomatization, two-electron reduction pathway, leading to the formation of complex **V**.¹⁶

Our studies have focused on the coordination chemistry and catalytic properties of hitherto understudied alkaline-earth complexes.¹⁷ While rapid progress has been made over the past decade, one major challenge remains the stabilization of heteroleptic species of the type $[\text{LAeR}]$ (L = monoanionic ligand, Ae = alkaline-earth dication, R = reactive coligand), as the heavier alkaline-earth-metal complexes are prone to Schlenk-type ligand redistribution processes in solution, which hamper catalytic activity.¹⁸ Examples of heteroleptic group 2 alkyl complexes remain particularly rare, as many conventional ligand systems, such as the ubiquitous β -diketiminato ligand $[\text{HC}\{\text{C}(\text{Me})\text{Ndipp}\}_2]^-$ (dipp = 2,6-diisopropylphenyl), are prone to deprotonation reactions by the highly reactive alkaline-earth-bound alkyl functionality.¹⁹ Thus, only the calcium β -diketiminato bis(trimethylsilyl)methyl derivative **VI** could be prepared from the reaction between $[\{\text{HC}\{\text{C}(\text{Me})\text{Ndipp}\}_2\}\text{H}]$ and the homoleptic precursor $[\text{Ca}\{\text{CH}(\text{SiMe}_3)_2(\text{THF})_2\}]$.^{19b} Harder and co-workers have also reported several heteroleptic calcium, strontium, and barium benzyl complexes for the catalytic polymerization of styrene.²⁰ More recently, Carpentier and Sarazin have described a series of analogous heteroleptic heavier alkaline-earth alkyl complexes, compounds **VIIa–c**, supported by a sterically demanding iminoanilide ligand, and reported their successful application as precatalysts for intra- and intermolecular hydroamination reactions.²¹



We have reported that bis(imino)pyridine ligands undergo double deprotonation of the β -imino-methyl substituents in the presence of the alkaline-earth dialkyls $[\text{M}\{\text{CH}(\text{SiMe}_3)_2\}_2(\text{THF})_2]$ (M = Mg, Ca, Sr, Ba). In the case of the calcium, strontium, and barium precursors this reaction proceeded via the formation of unstable, intermediate, heteroleptic alkaline-earth alkyl complexes, such as **VIII** and

IX, supported by a *meta*- or *para*-alkylated dearomatized (imine-amido)dihydropyridine ligand.²² A similar strategy had already been employed to obtain heteroleptic dihydroterpyridide lanthanide alkyl complexes in a selective alkylation/dearomatization reaction.²³

We have previously described, in preliminary form, dearomatization reactions of *dipp*-substituted bis(imino)-acenaphthenes with group 1 and group 2 derivatives of the sterically demanding bis(trimethylsilyl)methyl anion.²⁴ In this contribution we provide a complete description of this reactivity and the application of the resultant heteroleptic group 2 species for the intramolecular hydroamination of hindered aminoalkenes.

EXPERIMENTAL SECTION

General Procedures. All glassware was dried for 24 h in an oven at 150 °C prior to use. Solvents for air- and moisture-sensitive reactions were provided by an Innovative Technology Solvent Purification System. Hexane and toluene were used as collected. THF and diethyl ether were dried by distillation over sodium-benzophenone ketyl. All solvents were stored under argon over molecular sieves activated at 150 °C in vacuo. C_6D_6 and d_8 -toluene were purchased from Fluorochem and dried over molten potassium prior to vacuum transfer into a sealed ampule and storage in the glovebox under argon. Tetrakis(trimethylsilyl)silane (TMSS) was purchased from Goss Scientific Instruments Ltd. and used as received. After drying over CaH_2 all aminoalkenes were distilled in vacuo and either vacuum-transferred into sealed ampules or freeze-dried prior to transfer into the glovebox. The aminoalkene substrates,²⁵ the ligand precursor **I**,²⁸ the potassium precursors $[\text{KCH}_2\text{Ph}]$ ²⁶ and $[\text{KCH}(\text{SiMe}_3)_2]$,²⁷ and the homoleptic group 2 alkyls $[\text{M}\{\text{CH}(\text{SiMe}_3)_2\}_2(\text{THF})_2]$ (M = Mg, Ca, Sr, Ba)²⁸ were synthesized following literature procedures. Reactions were carried out under an argon atmosphere using standard Schlenk line and glovebox techniques in an MBraun Labmaster glovebox at O_2 , H_2O <0.1 ppm. NMR experiments were conducted in Youngs tap NMR tubes prepared and sealed in a glovebox under argon. NMR data was acquired at room temperature on a Bruker 300 Ultrashield (^{13}C , 75.48 MHz) and spectra were referenced using residual solvent resonances. Elemental analyses were performed by Stephen Boyer at London Metropolitan Enterprises.

Synthesis of Compounds 1–9. The numbering scheme used for the NMR assignments of compounds 1–9 follows that in Schemes 1 and 2.

Synthesis of 1. A 25 mg portion of $[\text{KCH}(\text{SiMe}_3)_2]$ (99.8 μmol) and 50 mg of **I** (99.8 μmol) were dissolved in C_6D_6 . The reaction mixture instantly turned dark green (NMR yield >99%). Green crystals suitable for X-ray crystallography were isolated upon crystallization from toluene at room temperature. The reaction was later scaled up in toluene with $[\text{KCH}(\text{SiMe}_3)_2]$ (396 mg, 2.00 mmol) and **I** (1.00 g, 2.00 mmol): 95% isolated yield (1.32 g, 1.90 mmol). ^1H NMR ppm (C_6D_6): 7.00–7.21 (m, 6H, Ar-H), 6.51 (d, 1H, H-3', $^3J = 7.5$ Hz), 6.03 (t, 1H, H-4', $^3J = 7.5$ Hz), 5.65 (d, 1H, H-5', $^3J = 7.5$ Hz), 5.60 (d, 1H, H-3, $^3J_{\text{cis}} = 9.8$ Hz), 4.65 (dd, 1H, H-4, $^3J_{\text{cis}} = 9.8$ Hz, $^3J = 4.1$ Hz), 4.41 (broad, 1H, H-5), 3.59–3.60 (m, 2H, $^i\text{Pr-CHMe}_2$), 3.06 + 2.98 (two sept, 1H each, $^i\text{Pr-CHMe}_2$, $^3J = 6.7$ Hz), 1.51 + 1.44 (two d, 3H each, $^i\text{Pr-CH}_3$, $^3J = 6.7$ Hz), 1.23 + 1.17 + 1.13 (three d, 9H:3H:6H, $^i\text{Pr-CH}_3$, $^3J = 6.7$ Hz), 0.55 (s, 1H, CCH(SiMe_3)₂), 0.18 + 0.04 (two s, 9H each, CCH(SiMe_3)₂). $^{13}\text{C}\{^1\text{H}\}$ NMR ppm (C_6D_6): 175.7 (C-1'), 153.9, 152.5, 148.7, 148.5, 141.8, 141.6, 136.6, 136.4, 129.9, 128.2, 127.9, 125.4, 124.4, 124.2, 123.7 (C-5' + C-3), 123.1, 123.0, 122.9, 121.2 (C-4'), 120.4, 116.8 (C-4), 102.7, 38.5 (C-5), 29.1 + 28.9 + 28.6 + 28.5 ($^i\text{Pr-CHMe}_2$), 25.2 (CCH(SiMe_3)₂), 24.9 + 24.7 + 24.6 + 24.4 + 23.9 + 23.8 ($^i\text{Pr-CH}_3$), 2.3 + 2.1 (CCH(SiMe_3)₂). Anal. Calcd for $\text{C}_{43}\text{H}_{59}\text{KN}_2\text{Si}_2$ (699.2): C, 73.86; H, 8.51; N, 4.01. Found: C, 73.79; H, 8.59; N, 3.96. ESI-MS for the protonated dearomatized ligand $[\text{C}_{43}\text{H}_{60}\text{N}_2\text{Si}_2 - \text{Na}]^+$: calcd 683.41928, found 683.4846.

Synthesis of 2. [$K\{CH(SiMe_3)_2(2-NMe_2-Ph)\}$] (25 mg, 83.5 μ mol) and **I** (42 mg, 83.5 μ mol) were dissolved in d_8 -toluene. The solution instantly turned dark green. Crystallization after 1 day at room temperature afforded dark green crystals of the R,S + S,R diastereomer of complex **2** (51 mg, 68.3 μ mol, 82%). 1H NMR ppm (d_8 -tol): major R,S + S,R diastereomer (88%), 6.88–7.20 (m, 10H, Ar-H), 6.04 (d, 1H, $H-3'$, $^3J = 7.7$ Hz), 5.60 (t, 1H, $H-4'$, $^3J = 7.7$ Hz), 5.44–5.50 (m, 2H, $H-3/S'$), 4.63–4.67 (m, 1H, $H-4$), 3.81 (broad s, 1H, $H-5$), 3.49 (sept, 2H, $^iPr-CHMe_2$, $^3J = 6.7$ Hz), 3.35 (s, 1H, $CH(SiMe_3)$), 2.95 (sept, 1H, $^iPr-CHMe_2$, $^3J = 6.6$ Hz), 2.85 (broad m, 1H, $^iPr-CHMe_2$), 2.33 (s, 6H, NMe_2), 1.52 (d, 3H, $^iPr-CH_3$, $^3J = 6.6$ Hz), 1.30–1.44 (m, 6H, $^iPr-CH_3$), 0.98–0.19 (m, 12H, $^iPr-CH_3$), 0.81 (d, 3H, $^iPr-CH_3$, $^3J = 6.7$ Hz), –0.03 (s, 9H, $SiMe_3$); minor R,R + S,S diastereomer (12%), 6.66 (d, 1H, $H-3'$, $^3J = 7.7$ Hz), 5.85–5.88 (m, 1H, $H-4'$), 3.46–3.49 (m, 1H, $H-3/S'$), 4.78 (broad d, 1H, $H-4$, $^3J_{cis} = 10.1$ Hz), 3.81 (broad s, 1H, $H-5$), 3.53 (m, 2H, $^iPr-CHMe_2$ + $CH(SiMe_3)$, identified by COSY), 2.95 + 2.85 (two broad sept, 1H each, $^iPr-CHMe_2$), 2.45 (s, 6H, NMe_2), 0.99–1.57 (m, 24H, $^iPr-CH_3$), 0.00 (s, 9H, $SiMe_3$). $^{13}C\{^1H\}$ NMR ppm (d_8 -tol): mixture of both diastereomers, 174.1 (C-1), 153.2, 152.6, 148.0, 141.1, 140.8, 140.1, 137.8, 137.7, 137.6, 137.2, 137.1, 136.1, 135.1, 132.1 (C-3'), 128.1, 125.3, 123.8, 123.7, 123.5, 123.0 (C-3/S'), 122.5 (C3/S'), 120.7, 119.3 (C-4'), 111.8 (C-4), 45.1 (NMe_2), 42.1 (C-5), 40.1 ($CH(SiMe_3)$), 29.2 + 29.1 + 28.8 + 28.4 ($^iPr-CHMe_2$), 25.7 + 25.2 + 25.1 + 24.5 + 24.4 + 24.2 + 23.5 + 23.4 + 23.2 ($^iPr-CH_3$), –0.04 ($SiMe_3$). Anal. Calcd for $C_{288}H_{360}K_6N_{18}Si_6$ (4477.14): C, 77.26; H, 8.10; N, 5.63. Found: C, 77.22; H, 8.19; N, 5.58.

Synthesis of 3. A 50 mg portion of [$Mg\{CH(SiMe_3)_2\}_2(THF)_2$] (103.9 μ mol) and 52 mg of **I** (103.9 μ mol) were dissolved in C_6D_6 . The reaction mixture was heated to 60 °C overnight and slowly turned dark green. Analysis by 1H NMR spectroscopy showed the clean formation of complex **3** as the sole product of the reaction (NMR yield >99%). The reaction was scaled up ([$Mg\{CH(SiMe_3)_2\}_2(THF)_2$], 271 mg, 0.599 mmol; **I**, 300 mg, 0.599 mmol), and the mixture was heated for 24 h at 60 °C in toluene prior to removal of the solvent in vacuo. As the product refused to crystallize in 0.2 mL of pentane at –30 °C over a period of several weeks, complex **3** was used as a crude green solid, whose purity was confirmed by NMR and elemental analysis data. 1H NMR ppm (C_6D_6): 7.15–7.25 (m, 6H, Ar-H), 6.45 (d, 1H, $H-3'$, $^3J = 7.7$ Hz), 6.05 (t, 1H, $H-4'$, $^3J = 7.7$ Hz), 5.47 (dd, 1H, $H-3$, $^3J_{cis} = 10.2$ Hz, $^4J = 2.3$ Hz), 5.44–5.47 (m, 1H, $H-5'$), 5.05 (dd, 1H, $H-4$, $^3J_{cis} = 10.2$, $^3J = 4.4$ Hz), 4.07 (m, 1H, $H-5$), 3.78 (broad m, THF), 3.55 + 3.11 (two sept, 2H each, $^iPr-CHMe_2$, $^3J = 6.8$ Hz), 1.41–1.44 (m, 14H, $^iPr-CH_3$ + THF), 1.39 + 1.36 + 1.31 + 1.29 + 1.16 + 1.13 (six d, 3H each, $^iPr-CH_3$, $^3J = 6.8$ Hz), 0.46 (d, 1H, $CCH(SiMe_3)_2$, $^3J = 1.6$ Hz), 0.00 (s, 18H, $MgCH(SiMe_3)_2$), 0.01 + –0.03 (two s, 9H each, $CCH(SiMe_3)_2$), –1.59 (s, 1H, $MgCH(SiMe_3)_2$). $^{13}C\{^1H\}$ NMR ppm (C_6D_6): 181.9 (C-1'), 149.5, 148.8, 148.2, 146.5, 144.5, 139.5, 139.4, 133.1, 133.0 (C-3'), 131.8, 127.2, 126.0, 125.8 (C-4), 125.2, 124.9, 124.8 (C-5'), 124.6, 124.5, 124.0, 123.4 (C-4'), 122.6 (C-3), 122.1, 111.8 (C-2), 68.9 (THF, very broad), 38.4 (C-5), 29.1 + 28.9 + 28.6 + 28.6 ($^iPr-CHMe_2$), 25.9 + 25.33 + 25.29 + 25.27 + 25.1 + 24.9 + 24.8 + 24.7 ($^iPr-CH_3$), 23.9 (THF), 22.7 ($CCH(SiMe_3)_2$), 5.6 ($MgCH(SiMe_3)_2$), 2.1 + 1.0 ($CCH(SiMe_3)_2$), –3.1 ($MgCH(SiMe_3)_2$). Anal. Calcd for $C_{54}H_{86}MgN_2OSi_4$ (915.9): C, 70.81; H, 9.46; N, 3.06. Found: C, 70.79; H, 9.36; N, 2.95.


Synthesis of 4. A 50 mg portion of [$Ca\{CH(SiMe_3)_2\}_2(THF)_2$] (99.8 μ mol) and 50 mg of **I** (99.8 μ mol) were dissolved in C_6D_6 . The reaction mixture instantly turned dark green. 1H NMR data showed the clean formation of complex **4** as the sole product of the reaction (NMR yield >99%). The reaction was scaled up ([$Ca\{CH(SiMe_3)_2\}_2(THF)_2$], 1.00 g, 2 mmol; **I**, 1.00 g, 2 mmol), and the mixture was stirred for 10 min at room temperature in toluene prior to removal of the solvent in vacuo. As the product refused to crystallize in 0.2 mL of pentane at –30 °C over a period of several weeks, complex **4** was used as a crude green solid, whose purity was confirmed by NMR spectroscopy and elemental analysis data. 1H NMR ppm (C_6D_6): 7.15–7.26 (m, 6H, Ar-H), 6.46 (d, 1H, $H-3'$, $^3J = 7.7$ Hz), 6.06 (t, 1H, $H-4'$, $^3J = 7.7$ Hz), 5.43–5.47 (m, 2H, $H-5' + H-3$), 4.96

(dd, 1H, $^3J_{cis} = 10.2$, $^3J = 4.4$ Hz), 4.12 (m, 1H, $H-5$), 3.62 (sept, 2H, $^iPr-CHMe_2$, $^3J = 6.8$ Hz), 3.51 (broad m, THF), 3.24 (sept, 2H, $^iPr-CHMe_2$, $^3J = 6.7$ Hz), 1.49 (d, 6H each, $^iPr-CH_3$, $^3J = 6.7$ Hz), 1.41 + 1.34 (two d, 3H each, $^iPr-CH_3$, $^3J = 6.8$ Hz), 1.18–1.31 (m, 20H, THF + $^iPr-CH_3$), 0.48 (d, 1H, $CCH(SiMe_3)_2$, $^3J = 1.6$ Hz), 0.21 (s, 18H, $CaCH(SiMe_3)_2$), 0.04 + 0.00 (two s, 9H each, $CCH(SiMe_3)_2$), –1.72 (s, 1H, $CaCH(SiMe_3)_2$). $^{13}C\{^1H\}$ NMR ppm (C_6D_6): 181.7 (C-1'), 150.1, 149.7, 148.2, 146.1, 143.3, 142.7, 138.8, 137.0, 132.5 (C-3'), 131.2, 126.4, 126.2, 124.5, 124.5 (C-3), 124.4, 124.0, 123.8, 123.5 (C-4), 123.1, 122.7 (C-5'), 122.6 (C-4'), 110.7 (C-2), 69.6 (THF, very broad), 38.3 (C-5), 29.3 + 29.2 + 28.9 + 28.5 ($^iPr-CHMe_2$), 26.0 + 25.6 + 25.3 + 25.2 + 25.2 + 24.7 + 24.4 + 24.3 ($^iPr-CH_3$), 25.3 (THF), 23.4 ($CCH(SiMe_3)_2$), 14.3 ($CaCH(SiMe_3)_2$), 6.1 ($CaCH(SiMe_3)_2$), 2.1 + 1.0 ($CCH(SiMe_3)_2$). Anal. Calcd for $C_{58}H_{94}CaN_2O_2Si_4$ (1003.8): C, 69.40; H, 9.44; N, 2.79. Found: C, 69.34; H, 9.34; N, 2.73.

Synthesis of 5. A 50 mg portion of [$Sr\{CH(SiMe_3)_2\}_2(THF)_2$] (90.8 μ mol) and 45 mg of **I** (90.8 μ mol) were dissolved in C_6D_6 . The reaction mixture instantly turned dark green. Analysis by 1H NMR spectroscopy showed the clean formation of complex **5** as the sole product of the reaction (NMR yield >99%). The reaction was scaled up ([$Sr\{CH(SiMe_3)_2\}_2(THF)_2$], 551 mg, 1 mmol; **I**, 0.50 g, 1 mmol), and the mixture was stirred for 10 min at room temperature in toluene prior to removal of the solvent in vacuo. As the product refused to crystallize in 0.2 mL of pentane at –30 °C over a period of several weeks, complex **5** was used as a crude green solid, whose purity was confirmed by NMR spectroscopy and elemental analysis data. 1H NMR ppm (C_6D_6): 6.97–7.16 (m, 6H, Ar-H), 6.45 (d, 1H, $H-3'$, $^3J = 7.7$ Hz), 6.06 (t, 1H, $H-4'$, $^3J = 7.7$ Hz), 5.41 (dd, 1H, $H-3$, $^3J_{cis} = 10.2$ Hz, $^4J = 2.1$ Hz), 5.40–5.42 (m, 1H, $H-5'$), 4.89 (dd, 1H, $H-4$, $^3J_{cis} = 10.2$, $^3J = 4.6$ Hz), 4.15 (m, 1H, $H-2$), 3.68 (sept, 2H, $^iPr-CHMe_2$, $^3J = 6.8$ Hz), 3.37 (broad m, THF), 3.27 (sept, 2H, $^iPr-CHMe_2$, $^3J = 6.7$ Hz), 1.53 + 1.52 (two d, 3H each, $^iPr-CH_3$, $^3J = 6.7$ Hz), 1.38 + 1.36 (two d, 3H each, $^iPr-CH_3$, $^3J = 6.8$ Hz), 1.25–1.31 (m, 20H, $^iPr-CH_3$ + THF), 0.49 (d, 1H, $CCH(SiMe_3)_2$, $^3J = 1.6$ Hz), 0.30 (s, 18H, $SrCH(SiMe_3)_2$), 0.06 + –0.01 (two s, 9H each, $CCH(SiMe_3)_2$), –1.86 (s, 1H, $SrCH(SiMe_3)_2$). $^{13}C\{^1H\}$ NMR ppm (C_6D_6): 181.1 (C-1'), 151.1, 150.2, 149.0, 146.9, 143.3, 142.6, 138.6, 138.4, 132.4 (C-3'), 131.0, 126.3, 125.9, 124.5 (C-5'), 124.2, 124.3, 123.7, 123.1, 123.0, 122.9 (C-3), 122.5 (C-4), 122.2 (C-4'), 109.5 (C-2), 69.4 (THF, very broad), 38.4 (C-5), 29.5 + 29.3 + 29.0 + 28.6 ($^iPr-CHMe_2$), 26.0 + 25.7 + 25.4 + 25.3 + 25.2 + 24.9 + 24.4 + 24.3 ($^iPr-CH_3$), 25.3 (THF), 23.6 ($CCH(SiMe_3)_2$), 19.1 ($SrCH(SiMe_3)_2$), 6.4 ($SrCH(SiMe_3)_2$), 2.2 + 1.1 ($CCH(SiMe_3)_2$). Anal. Calcd for $C_{58}H_{94}N_2O_2Si_4Sr$ (1051.3): C, 66.26; H, 9.01; N, 2.66. Found: C, 66.33; H, 8.97; N, 2.70.

Synthesis of 6. A 50 mg portion of [$Ba\{CH(SiMe_3)_2\}_2(THF)_2$] (83.3 μ mol) and 42 mg of **I** (83.3 μ mol) were dissolved in C_6D_6 . The reaction mixture instantly turned dark green. 1H NMR data evidenced the presence of three components, among which were the heteroleptic barium alkyl complex **6** (ca. 70%) and the homoleptic barium complex **7** (ca. 20%). Upon recrystallization of the crude mixture from toluene at –30 °C for 1 week, orange crystals of compound **8** suitable for X-ray diffraction analysis were isolated from the reaction mixture. The isolated amount of **8** was, however, insufficient for the acquisition of either NMR or elemental analysis data. Although attempts to separate **6** and **7** by fractional crystallization failed, preventing the acquisition of elemental analysis data for compound **6**, compound **7** was synthesized and analyzed by independent methods (vide infra). 1H NMR ppm (C_6D_6) for **6**: 7.08–7.23 (m, 6H, Ar-H), 6.48 (d, 1H, $H-3'$, $^3J = 7.7$ Hz), 6.08 (t, 1H, $H-4'$, $^3J = 7.7$ Hz), 5.47–5.51 (m, 2H, $H-5' + H-4$), 4.88 (dd, 1H, $H-4$, $^3J_{cis} = 10.2$, $^3J = 4.4$ Hz), 4.21 (m, 1H, $H-5$), 3.69 (sept, 2H, $^iPr-CHMe_2$, $^3J = 6.8$ Hz), 3.32 (broad m, 8H, THF), 3.27 (sept, 2H, $^iPr-CHMe_2$, $^3J = 6.7$ Hz), 1.53 + 1.49 (two d, 3H each, $^iPr-CH_3$, $^3J = 6.7$ Hz), 1.37 + 1.34 (two d, 3H each, $^iPr-CH_3$, $^3J = 6.8$ Hz), 1.22–1.32 (m, 20H, THF + $^iPr-CH_3$), 0.50 (d, 1H, $CCH(SiMe_3)_2$, $^3J = 1.6$ Hz), 0.29 (s, 18H, $BaCH(SiMe_3)_2$), 0.09 + 0.04 (two s, 9H each, $CCH(SiMe_3)_2$), –1.73 (broad s, 1H, $BaCH(SiMe_3)_2$). $^{13}C\{^1H\}$ NMR ppm (C_6D_6): 179.9 (C-1'), 150.8, 149.8, 148.8, 147.0, 143.2, 142.7, 138.0, 135.7, 132.2 (C-3'), 131.1, 126.1, 125.6, 124.4 (C-5'), 124.2, 124.0, 123.9, 123.8, 123.6, 123.2, 122.9 (C-3), 122.6 (C-4), 122.0 (C-4'), 107.5 (C-2), 68.3 (THF, very broad), 38.4 (C-5), 33.8 (Ba–

Scheme 1



CH(SiMe₃)₂), 29.6 + 29.5 + 29.0 + 28.8 (¹Pr-CHMe₂), 25.6 + 25.5 + 25.2 + 25.1 + 25.0 + 24.9 + 24.5 + 24.3 (¹Pr-CH₃), 25.4 (THF), 23.8 (CCH(SiMe₃)₂), 6.1 (BaCH(SiMe₃)₂), 2.2 + 1.4 (CCH(SiMe₃)₂).

Independent Synthesis of 7. (a) A 15 mg portion of [Ba{CH(SiMe₃)₂}(THF)₂] (25.0 μmol) and 25 mg of I (50 μmol) were dissolved in *d*₈-toluene at room temperature. After 10 min the solution had turned a homogeneous dark green. Analysis by ¹H NMR spectroscopy showed the clean formation of 7 as the sole product of the reaction.

(b) [KCH(SiMe₃)₂] (300 mg, 1.50 mmol) and [Ba{CH(SiMe₃)₂}(THF)₂] (455 mg, 0.75 mmol) were stirred in THF at room temperature for 24 h prior to removal of the solvent in vacuo and extraction into toluene. Removal of the solvent yielded complex 7 as a crude dark green solid whose ¹H NMR spectrum was reminiscent of that obtained from the reaction described in (a). ¹H NMR ppm (*d*₈-tol): 6.84–7.32 (m, 12H, Ar-H), 6.46 (d, 2H, H-3', ³J = 7.7 Hz), 6.00 (t, 2H, H-4', ³J = 7.7 Hz), 5.34 (dd, 2H, H-5', ³J = 7.7 Hz, ⁴J = 4.2 Hz), 5.17 (dd, 2H, H-3, ³J_{cis} = 10.2 Hz, ⁴J = 1.5 Hz), 4.77 (dd, 2H, H-4, ³J_{cis} = 10.2, ³J = 4.3 Hz), 4.16 (broad s, 2H, H-5), 3.46–3.62 (m, 12H, THF + ¹Pr-CHMe₂), 3.09 (sept, 4H, ¹Pr-CHMe₂, ³J = 6.6 Hz), 1.45 (m, 8H, THF), 1.38 + 1.35 (two d, 6H each, ¹Pr-CH₃, ³J = 6.7 Hz), 1.24 (d, 6H, ¹Pr-CH₃, ³J = 6.8 Hz), 1.03–1.14 (m, 30H, ¹Pr-CH₃), 0.49 (broad s, 2H, CH(SiMe₃)₂, ³J = 1.6 Hz), 0.02 + -0.04 (two s, 9H each, SiMe₃). ¹³C{¹H} NMR ppm (*d*₈-tol): 180.3 (C-1'), 151.0, 150.70, 150.69, 149.9, 147.69, 147.64, 143.7, 143.25, 143.22, 138.73, 138.70, 138.61, 138.56, 138.15, 138.23, 137.47, 137.57, 132.5, 131.1, 126.7, 126.0, 124.9, 124.65, 124.71, 124.6, 123.77, 123.73, 123.5, 122.3, 122.1, 109.22, 109.19, 68.0 (THF, broad), 38.8 (C-5), 29.9 + 29.7 + 29.5 + 29.1 + 28.8 (¹Pr-CHMe₂), 26.24 + 26.18 + 26.11 + 25.9 + 25.5 + 24.4 + 23.8 (¹Pr-CH₃), 25.1 (THF), 23.4 + 23.2 (CH(SiMe₃)₂), 2.8 + 1.3 (SiMe₃). Repeated attempts at obtaining elemental analysis data for this complex failed due to its extreme air and moisture sensitivity.

Independent Synthesis of 9. Complex 1 (1.05 g, 1.50 mmol) and CaI₂ (220 mg, 0.75 mmol) were stirred in THF at room temperature for 24 h prior to removal of the solvent in vacuo and extraction into hexane. Removal of the solvent yielded complex 9 as a crude dark green solid which was recrystallized from a 10/1 toluene/THF mixture at -30 °C over several days, yielding complex 9 as a dark green solid (265 mg, 25%) in a first fraction and green single crystals in a second fraction (227 mg, 21%). Because of apparent fluxional behavior in solution ¹H NMR resonances were extremely broad and ¹³C{¹H} NMR signals were too weak for analysis. ¹H NMR (C₆D₆): 6.99–7.25 (m, 12H, Ar-H), 6.45 (broad d, 2H, H-3', ³J = 7.7 Hz), 6.05 (broad t, 2H, H-4', ³J = 7.7 Hz), 5.23–5.38 (broad m, 4H, H-3/4), 4.99 (broad d, 2H, H-5', ³J = 7.7 Hz), 4.18 (s, 2H, H-5), 3.60 (broad m, 4H, THF), 2.99–3.44 (broad m, 8H, ¹Pr-CHMe₂), 0.68–1.34 (broad m, 52H each, ¹Pr-CH₃ + THF), 0.51 (broad s, 1H, CCH(SiMe₃)₂), 0.05 + -0.04 (two broad s, 9H each, CCH(SiMe₃)₂). Anal. Calcd for C₉₀H₁₂₆CaN₄OSi₄ (1432.4): C, 75.46; H, 8.87; N, 3.91. Found: C, 75.29; H, 8.74; N, 3.84.

Notes on X-ray Crystallographic Refinements. Single-crystal X-ray diffraction data were collected at 150 K on a Nonius KappaCCD diffractometer, equipped with an Oxford Cryosystem, using graphite-monochromated Mo K α radiation ($\lambda = 0.71073$ Å). Data were processed using the Nonius software.²⁹ Crystal data and data collection and refinement parameters for 2, 8, and 9 are presented in Table S1 (Supporting Information). Structure solution, followed by full-matrix least-squares refinement, was performed using the WINGX-1.80 suite of programs throughout.³⁰ Compound 1: the dimer

cocrystallized with one solvent molecule of toluene. Compound 2: the asymmetric unit contains one-sixth of a hexameric complex. It also contains one toluene solvent molecule with 50% occupation, disordered over a 2-fold axis. Another toluene molecule is disordered about a 3-fold axis with 33% occupation. Traces of a further solvent molecule were located over a 3-fold axis but could not be properly solved. The very large crystal diffracted very well at low angle but presented a falloff in intensity above $2\theta = 35^\circ$. As a consequence, the solvent disorders could not be solved satisfactorily. The PLATON program SQUEEZE was employed to account for the solvent-containing voids. Checkcif detected a potential disorder in one of the isopropyl groups (C39–C41). Because of the lack of high-angle intensity data this could not be properly solved. Compound 2: the crystal was small and only weakly diffracting, hence the high $R(\text{int})$ value. The asymmetric unit contained the main complex and three solvent molecules: two half-occupied THF molecules (bond lengths to O3 had to be restrained) and one benzene molecule disordered over two sites in a 63:37 ratio. Atoms of the minor part of the disordered benzene were refined isotropically. ADP for C122 had been restrained. All methyl groups of the bis(trimethylsilyl)methyl substituent containing Si1 and Si2 displayed rotational disorder in a 42:58 ratio. Si–C bond lengths herein have been restrained as well as ADPs of C39 and C43.

RESULTS AND DISCUSSION

Addition of the potassium alkyl precursors [KCH(SiMe₃)₂] and [KCH(SiMe₃)₂]{2-(NMe₂)C₆H₄} to an orange suspension of I in C₆D₆ resulted in a rapid color change to form a clear, dark green solution (Scheme 1). In both cases analysis by ¹H NMR spectroscopy showed quantitative conversion of the starting materials to a single potassium complex, compounds 1 and 2, respectively, displaying six distinct multiplets in the region from 4.4 to 6.5 ppm, characteristic of selective dearomatization of the acenaphthene backbone and alkyl functionality transfer. The alkylation site of the ligand backbone was identified by a COSY experiment as the C⁵ position (Scheme 1), which was confirmed by two subsequent X-ray diffraction experiments (vide infra). Alkylation at this position generated a chiral center which caused the two magnetically inequivalent (Si(CH₃)₃) groups in both complexes to appear as two separate (9H) singlets. The asymmetric nature of the new imine-amido ligand system also caused splitting of the isopropyl methine and methyl resonances, in both the ¹H and ¹³C{¹H} NMR spectra. In the case of complex 2 the presence of a second chiral center at the methine position of the 1-trimethylsilyl-*o*-dimethylaminobenzyl substituent led to the formation of a ca. 9:1 mixture of two diastereomers, as determined by integration of the ¹H NMR spectrum.

Recrystallization at room temperature in toluene afforded large dark green crystals of compounds 1 and 2 suitable for X-ray diffraction. Although the structure of compound 1 was described in our earlier communication²⁴ and will thus not be described in detail here, selected bond length and angle data for both compounds are given in Table 1 for purposes of comparison. While compound 1 comprised a centrosymmetric dimer in the solid state, compound 2 assembles as a hexameric array in which coordination between molecules is provided by a

Table 1. Selected Bond Lengths (Å) and Angles (deg) for Compounds 1, 2, 8, and 9

	1 ^a	2 ^a	8	9 ^b
M–N1	2.694(3)	2.702(4)		2.571(4)
M–N2	2.703(3)	2.762(4)		2.372(4)
N1–C25	1.281(4)	1.267(5)	1.263(4)	1.286(7)
N2–C36	1.335(4)	1.318(5)	1.283(5)	1.355(7)
C34–C36	1.393(4)	1.415(6)	1.487(5)	1.418(8)
C30–C31	1.533(5)	1.524(6)	1.523(5)	1.519(8)
C31–C32	1.517(5)	1.514(7)	1.563(5)	1.510(9)
C32–C33	1.351(5)	1.390(6)	1.501(5)	1.345(8)
M–centroid	2.866(4)	2.884(5)		
C32–C32'			1.557(6)	
N1–M–N2	63.52(8)	61.95(11)		70.07(15)
M–N1–C25	118.2(2)	119.7(3)		109.6(3)
M–N2–C36	117.4(2)	118.2(3)		114.7(3)
N1–C25–C36	121.7(3)	119.0(4)	122.0(3)	121.6(5)
C30–C31–C32	111.6(3)	109.6(4)	114.4(3)	111.7(5)
C30–C31–C37	113.4(3)	110.5(4)	114.7(3)	113.7(5)
C32–C31–C37	111.3(3)	109.7(4)	110.1(3)	111.8(5)
C31–C32–C33			117.9(3)	126.3(6)
C33–C32–C32'			109.9(2)	
C31–C32–C32'			110.7(4)	

^aM = K. ^bM = Ca.

η interaction between a N₂-coordinated potassium atom and the C30, C34, and C35 carbon atoms of the alkylated backbone ring of the adjacent complex, with a K–centroid distance of 2.884(5) Å (Figure 1). Compound 2 crystallizes as a single *R,S/S,R* diastereomeric couple which, on the basis of the crystalline yield (~80%) and its ¹H NMR spectrum, was identified as the major diastereomer. In contrast to potassium complex 1 the K–N bond lengths in compound 2 reflect the asymmetric binding

of the alkylated imine-amido ligand (K–N_{imine} = 2.762(4) Å; K–N_{amide} = 2.702(4) Å).

In a manner similar to the potassium-based reactions, addition of [M{CH(SiMe₃)₂}₂(THF)₂] (M = Ca, Sr) to a suspension of I in C₆D₆ resulted in instant solubilization of the starting materials and a rapid color change to dark green. The analogous reaction with [Mg{CH(SiMe₃)₂}₂(THF)₂] required 24 h of heating at 60 °C to yield the same result (Scheme 2). In all three cases ¹H NMR spectra were reminiscent of that of 1, showing clean formation of the heteroleptic alkaline-earth organometallic species 3–5, with the additional 18H and 1H singlets of the metal-bound monoanionic [CH(SiMe₃)₂][–] coligand at ca. 0.3 and –1.7 ppm, respectively (Figure 2).

Although an analogous reaction with the barium dialkyl precursor [Ba{CH(SiMe₃)₂}₂(THF)₂] also led to an instant color change to dark green, ¹H NMR data at the first point of analysis showed the formation of three distinct dearomatized species, among which the heteroleptic barium alkyl complex 6 (50–70% conversion) was identified by the characteristic upfield BaCH(SiMe₃)₂ proton shift at –1.73 ppm (Scheme 3). A subsequent independent synthesis of the homoleptic barium complex 7 from the reaction between 2 equiv of I and [Ba{CH(SiMe₃)₂}₂(THF)₂] enabled the identification of the second component of the reaction mixture as complex 7, present in 25–40% yield.

A third product, present in up to 10% yield, was only identified after storage of the reaction mixture in a 1:2 toluene/hexane solution at –30 °C over several days provided orange crystals suitable for X-ray diffraction analysis. The result of this experiment, displayed in Figure 3, shows two C⁵-alkylated dearomatized BIAN ligands coupled at the C⁴ position. Selected bond length and angle data are provided in Table 1. Compound 8 crystallizes as a single 4*R*,4'*R*,5*S*,5'*S* enantiomer (absolute structure parameter 0.03(15)) in the chiral space group *P*3₁2₁. As in the ligand precursor, both N–C bond






Figure 1. ORTEP representation of compound 2: hexameric array (left); two-molecule binding motif (right). Thermal ellipsoids are drawn at the 30% probability level. Isopropyl methyl groups and hydrogen atoms are omitted except for those attached to chiral centers C31 and C37.

Scheme 2

Figure 2. Annotated ¹H NMR spectrum of the dearomatized calcium complex 4.

Scheme 3



lengths (1.263(4), 1.283(5) Å) are indicative of imine bonds. The presence of the two pseudo-tetrahedral sp^3 carbon centers C31 ($C30-C31-C32 = 114.4(3)^\circ$; $C30-C31-C37 = 114.7(3)^\circ$; $C32-C31-C37 = 110.1(3)^\circ$) and C32 ($C31-C32-C33 = 117.9(3)^\circ$; $C33-C32-C32' = 109.9(2)^\circ$; $C31-C32-C32' = 110.7(4)^\circ$) induces a significant distortion of the alkylated C_6H_3 rings of the ligand backbones, accompanied by an increase in the C–C bond lengths ($C30-C31 = 1.523(5)$ Å; $C31-C32 = 1.563(5)$ Å; $C31-C32 = 1.501(5)$ Å). While we have not fully rationalized the synthesis of this chiral, dimerized tetraimine product, we suggest its formation is possibly the result of a sterically induced radical coupling reminiscent of well-precedented reactivity observed for similarly redox inactive 4f-element complexes.³¹ In support of this hypothesis, Evans and Cowley have observed a radical anion C^5 -alkylated free

radical as an intermediate in the formation of the dialkylated dearomatized lithium complex **V**.¹⁶

Multiple attempts to obtain crystals of the alkaline-earth organometallic complexes **3–6** suitable for X-ray analyses were thwarted by the extreme solubility of these compounds even after months of storage at -30 °C in minimal amounts of hydrocarbon solvents such as hexane and pentane. NMR data and elemental analyses of the crude magnesium, calcium, and strontium products, however, left no doubt as to their formulation. Attempts to recover the alkylated amino-imine ligand by controlled hydrolysis of the organometallic complexes initially led to a color change from green to purple, followed by a gradual change toward dark red over a period of several hours. ¹H NMR data and later recrystallization of the crude product from toluene evidenced quantitative conversion to the




Figure 3. ORTEP representation of compound **8**. Thermal ellipsoids are drawn at 30% probability. Isopropyl groups and hydrogen atoms are omitted except for those attached to the chiral centers C31 and C32.

Scheme 4





Figure 4. ORTEP representation of compound **9**. Thermal ellipsoids are drawn at the 30% probability level. Isopropyl methyl groups and hydrogen atoms are omitted, except for those attached to the chiral centers C31 and C74.

Table 2. Scope of Intramolecular Hydroamination with Precatalysts 3–5

Entry	Substrate	Product	Catalyst (mol %)	Time (h)	T (°C)	NMR yield (%) ^a
1a			3 (0.5)	18	60	<5
1b			4 (0.5)	0.25	25	>99
1c			5 (0.5)	1	25	76
2			4 (1.0)	0.3	25	99
3			4 (2.0)	1	25	95
4			4 (10.0)	24	80	0
5			4 (0.25)	0.1	25	98
6			4 (2.0)	1	25	98
7			4 (5.0)	5	25	95
8			4 (10.0)	18	60	93
9			4 (10.0)	48	80	0
10			4 (12.0)	40	60	98
11			4 (20.0)	80	80	0
12a			3 (5.0)	18	60	>99
12b			4 (2.0)	1	25	85 ^b
13			4 (20.0)	48	80	0
14a			3 (20.0)	24	80	95
14b			4 (20.0)	48	80	0 ^c
15			4 (1.0)	0.5	25	98

^aNMR yields were measured against an internal TMSS standard in C₆D₆ or *d*₈-toluene. ^b11% isomerization to the internal aminoalkene. ^c21% isomerization to the internal aminoalkene.

rearomatized ligand precursor **I**, with formation of the alkane CH₂(SiMe₃)₂. Although the mechanism of this reaction remains to be elucidated, it is possible that the high stability of the aromatic BIAN ligand may be the driving force for this dealkylation process.

The kinetic stability of the heteroleptic complexes **3–5** was assessed by heating a *d*₈-toluene solution of these compounds at 80 °C for 1 week. Apart from a slight amount of protonolysis of the alkyl ligand to form the alkane CH₂(SiMe₃)₂ in the cases of **4** and **5** (<5%), possibly due to reaction with the solvent, the ¹H NMR spectra remained virtually unchanged, suggesting an absence of Schlenk-type ligand redistribution processes. Addition of 2 equiv of the ligand precursor to 1 equiv of the magnesium, calcium, or strontium dialkyl species in *d*₈-toluene did not yield the expected homoleptic complexes at room temperature. Rather, the heteroleptic complexes **3–5** were formed while the excess ligand remained unreacted. Heating of these mixtures at 90 °C over a period of several days also did not result in reaction with the second equivalent of ligand.

Stoichiometric NMR-scale reactions of calcium complex **4** with protic substrates such as hexamethyldisilazane, diphenyl-

amine, and diphenylphosphine in C₆D₆ resulted in immediate protonolysis of the alkyl coligand, as observed in the ¹H NMR spectra by the disappearance of the upfield CaCH(SiMe₃)₂ singlet at –1.72 ppm and the appearance of the characteristic alkane resonance at –0.2 ppm (Scheme 4). Given the comparable steric demands of the bis(trimethylsilyl)amide and bis(trimethylsilyl)methanide coligands, the resulting heteroleptic calcium bis(trimethylsilyl)amide complex evidenced solution stability similar to that of complex **4**. In contrast, monitoring by ¹H NMR spectroscopy of the heteroleptic calcium complexes bearing the less sterically demanding diphenylamide and diphenylphosphide coligands under the same conditions suggested slow Schlenk-type ligand redistribution toward the homoleptic calcium species **9** characterized by the broadening and shifting of the dearomatized ligand backbone and isopropyl resonances, accompanied by precipitation of insoluble colorless products tentatively attributed to homoleptic calcium amide and phosphide species (Scheme 4), which are known to be insoluble in noncoordinating hydrocarbon solvents.³² An independent synthesis of complex **9** by the reaction between

2 equiv of potassium complex **1** and calcium iodide in THF confirmed the formation of **9** in both of these reactions.

Crystals of compound **9** suitable for X-ray diffraction analysis were obtained after storage in a 4/1 toluene/THF mixture at $-30\text{ }^{\circ}\text{C}$ for several weeks. The result of this experiment is displayed in Figure 4. Selected bond length and angle data are given in Table 1. Compound **9** crystallizes as a mono-THF adduct, and the five-coordinate calcium center displays a highly distorted trigonal bipyramidal geometry with both imine nitrogen atoms in axial positions. The mean planes formed by the two almost planar ligand backbones form an angle of 82.8° . While in the case of complex **1** the KN_2C_2 ring remains planar, the presence of the large calcium center in **9** causes a slight distortion in the CaN_2C_2 rings. The $\text{N1}-\text{Ca}-\text{N2}$ and $\text{N3}-\text{Ca}-\text{N4}$ bite angles of the two alkylated BIAN ligands ($70.07(15)$ and $69.65(14)^{\circ}$, respectively) are notably smaller than the $\text{N}-\text{Ca}-\text{N}$ angles observed in calcium complexes supported by a doubly reduced BIAN ligand ($75.76(7)-78.5(2)^{\circ}$)^{7f} or the homoleptic complex of the radical anion [$\{\text{I}^{\bullet}\}_2\text{Ca}$] ($73.58(9)^{\circ}$).^{7g} The $\text{Ca}-\text{N}_{\text{imine}}$ distances of $2.571(4)$ and $2.572(5)$ Å are substantially longer than the $\text{Ca}-\text{N}_{\text{amide}}$ bonds of $2.372(4)$ and $2.354(5)$ Å, reflecting the stronger bonding between the metal center and the anionic amide moiety of the ligand. As in the potassium complex **1** the $\text{N2}-\text{C36}$ and $\text{N4}-\text{C79}$ bond lengths of the amide residues ($1.355(7)$ and $1.347(7)$ Å) are significantly longer than the $\text{N1}=\text{C25}$ and $\text{N3}=\text{C68}$ bond lengths of the imine moieties ($1.286(7)$ and $1.278(7)$ Å), while the ethanediylidene bridges are slightly shorter in **9** than in **1** (9 , $1.489(8)$ and $1.509(8)$ Å; **1**, $1.518(4)$ Å). As previously observed for compound **1**, the dearomatization of the alkylated C_6H_3 rings only induces a slight distortion from planarity, despite the presence of the sp^3 C31 and C74 carbon centers indicated by the elongated C–C bond lengths ($\text{C30}-\text{C31} = 1.519(8)$ Å; $\text{C31}-\text{C32} = 1.510(9)$ Å; $\text{C73}-\text{C74} = 1.522(8)$ Å; $\text{C74}-\text{C75} = 1.497(8)$ Å) and the pseudo-tetrahedral angles ($\text{C32}-\text{C31}-\text{C30} = 111.7(5)$; $\text{C32}-\text{C31}-\text{C37} = 111.9(5)$; $\text{C30}-\text{C31}-\text{C37} = 113.7(5)$; $\text{C75}-\text{C74}-\text{C73} = 111.5(5)$; $\text{C75}-\text{C74}-\text{C80} = 112.0(5)$; $\text{C73}-\text{C74}-\text{C80} = 114.7(5)^{\circ}$).

Catalytic Intramolecular Hydroamination of Hindered Aminoalkenes. A preliminary comparison of turnover frequencies for the intramolecular hydroamination of 1-amino-2,2-diphenyl-4-pentene showed the calcium complex **4** to be superior by far to its magnesium and strontium analogues, with the order of reactivity being $4 > 5 \gg 3$ (Table 2, entries 1a–c). The same trend has been observed for the majority of previously described alkaline-earth-catalyzed hydroamination/cyclization reactions.^{21,24,25,33} Complex **4** was then assessed for the intramolecular hydroamination of a wide range of substituted aminoalkenes on a NMR scale in C_6D_6 or d_8 -toluene. In all cases careful inspection of the ^1H NMR spectra showed no change suggestive of ligand redistribution in the characteristic shifts of the dearomatized ligand backbone throughout the reaction times up to $80\text{ }^{\circ}\text{C}$, above which signs of Schlenk equilibria started to become apparent. As already observed in the case of homoleptic alkaline-earth dialkyl precatalysts and the heteroleptic iminoanilide species **VIIa–c**, the reactive bis(trimethylsilyl)methanide coligand of **4** greatly increases reaction rates in comparison to homoleptic or heteroleptic calcium bis(trimethylsilyl)amide precatalysts.^{21,24} This is due to the large $\text{p}K_{\text{a}}$ difference between $\text{CH}_2(\text{SiMe}_3)_2$ and the aminoalkene substrates, leading to rapid and irreversible catalyst activation, as shown by the complete

disappearance of the distinctive upfield alkyl singlet at ca. -1.7 ppm within the first point of analysis in all these reactions. In contrast, $\text{HN}(\text{SiMe}_3)_2$ may compete with the substrate for neutral coordination to the calcium center and reversible σ -bond metathesis.^{33c,34} Furthermore, the presence of the stabilizing dearomatized BIAN ligand leads to increased reactivity in comparison to the simple homoleptic precursor $[\text{Ca}\{\text{CH}(\text{SiMe}_3)_2\}_2(\text{THF})_2]$.²⁴ This positive spectator ligand effect has already been noted when comparing heteroleptic and homoleptic group 2 amide precatalysts^{24,33c} and is most pronounced with the iminoanilide species **VIIa–c**, which remain the most reactive precatalysts reported for hydroamination/cyclization reactions.²¹ Complex **4** displayed the usual reactivity trends, depending on the substitution pattern of the aminoalkene substrate. Substrates providing a favorable Thorpe–Ingold effect—i.e. with larger digeminal β -substituents—required lower precatalyst concentrations and shorter reaction times (Table 2, entries 1–4). It is notable that the 2,2-diphenyl-substituted substrate provided essentially quantitative conversion to the corresponding pyrrolidine within the first point of analysis at room temperature, using only 0.5 mol % of **4** (entry 1), while the simplest substrate, 1-amino-4-pentene, did not provide any reaction under the conditions tested (entry 4). In line with previous findings, catalytic turnover was also more efficient for smaller ring sizes, in the order $5 > 6 \gg 7$ (entries 1b, 12b, and 14b) and the calcium precatalyst **4** was prone to induce alkene isomerization to internal positions for longer aminoalkene chains (entries 13b and 14b). In contrast to this latter observation, the magnesium analogue afforded complete selectivity for the cyclized product (entries 12a and 14a). As has been previously observed with heteroleptic alkaline-earth β -diketiminate species the calcium precatalyst **4** proved incapable of effecting the cyclization of 1-amino-2,2-diphenyl-6-heptene to the corresponding hexahydroazepine, even under forcing conditions (entry 15b).^{25,33a} In contrast, the magnesium precatalyst **3** (20 mol %) provided clean and near-quantitative conversion to the desired seven-membered N-heterocycle over 24 h at $80\text{ }^{\circ}\text{C}$ (entry 15a). In comparison, the *n*-butyl magnesium β -diketiminate derivative required 5.5 days at $80\text{ }^{\circ}\text{C}$ to achieve only 88% conversion.²⁵ Internal substitution of the alkene moiety only caused a small decrease in reactivity (entry 6), whereas the activated terminally substituted substrate 1-amino-2,2,5-triphenyl-4-pentene was smoothly converted to the corresponding pyrrolidine under conditions even milder than those for the parent 1-amino-2,2-diphenyl-4-pentene (entry 1b). Most notable, however, was the relative ease with which complex **4** catalyzed the intramolecular hydroamination of terminally substituted, unactivated substrates. While to date only $[\text{Ca}\{\text{CH}(\text{SiMe}_3)_2\}_2(\text{THF})_2]$ has proved successful for the cyclization of 1-amino-2,2-diphenyl-4-hexene with modest yields under forcing conditions,^{33c} complex **4** provided near-quantitative conversion to the corresponding pyrrolidine within 5 h at room temperature and only 5 mol % catalyst loading (entry 7). A marked Thorpe–Ingold effect was also apparent for these terminally substituted unactivated substrates. Although 4-methyl-2,2-diphenylpent-4-en-1-amine and (*E*)-2,2-diphenylhex-4-en-1-amine underwent smooth cyclization (entries 6 and 7), complete loss of catalytic activity was encountered in the case of 1-amino-2,2-dimethyl-4-hexene under the reaction conditions assessed (entry 9). The slightly more challenging terminal ethyl derivative 1-amino-2,2-diphenyl-4-heptene also underwent clean cyclization to 2-methyl-4,4-diphenylpyrrolidine, albeit at higher catalyst loading

(10 mol %) and temperature (60 °C) (entry 10). The isomeric 1-amino-2,2-diphenyl-5-heptene, however, showed no sign of cyclization to the desired piperidine, even at high catalyst loadings of 4 and elevated reaction temperatures (entry 13). Finally, the precatalyst also proved efficient at the hydroamination of aminoalkenes bearing secondary amine functionalities, with little loss of activity, as shown by the facile cyclization of *N*-benzyl-1-amino-2,2-diphenyl-4-pentene (entry 15). In the cases of substrates which did not undergo cyclization (entries 4, 9, 11, 13, and 14b) stoichiometric reactions between 4 and 2 or 3 equiv of the aminoalkene led to instant protonolysis of the reactive alkyl ligand, and ¹H NMR suggested the formation of stable heteroleptic amidoalkene complexes supported by the dearomatized BIAN ligand and one or two neutrally adducted aminoalkene ligands.

CONCLUSION

In conclusion, we have shown that reactions of a dipp-substituted bis(imino)acenaphthene with a range of sterically demanding potassium alkyl or heavier alkaline-earth dialkyl reagents result in facile dearomatization through alkyl transfer to the C⁵ position of the conjugated acenaphthene system. The resultant heteroleptic group 2 derivatives display enhanced kinetic stability to Schlenk-type redistribution equilibria, which we suggest is a consequence of both the steric demands of the *N*-dipp and the rigidity provided by the dearomatized fused ring acenaphthene ligand backbone. Facile solution exchange is encountered, however, when the bis(trimethylsilyl)methyl substituent is replaced by an anionic ligand with lower overall steric demands. An assessment of the heteroleptic group 2 compounds as precatalysts for the intramolecular hydroamination of aminoalkenes evidences enhanced reactivity, which we ascribe to the greater solution stability of the catalytically active species. Most notably the calcium species 4 may even be applied to the high-yielding cyclization of substrates bearing alkyl substitution at either of the alkenyl positions. We are continuing to study this chemistry and to elaborate the application of these easily accessed organometallic species to a broader spectrum of demanding stoichiometric and catalytic reactions.

ASSOCIATED CONTENT

Supporting Information

Text, figures, a table, and CIF files giving full experimental and instrument details, characterization data for products of all catalytic cyclization reactions, and details of the X-ray diffraction analyses of compounds 1, 2, 8, and 9. This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Author

*E-mail for M.S.H.: msh27@bath.ac.uk.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We thank the Engineering and Physical Sciences Research Council for funding of this work (EP/I014519/1).

REFERENCES

(1) Dvolaitzky, M. C. R. *Séances Acad. Sci. Ser. C: Sci. Chim.* **1969**, 268, 1811.

(2) For a selection of publications on the topic, see: (a) El-Ayaan, U.; Paulovicova, A.; Yamada, S.; Fukuda, Y. *J. Coord. Chem.* **2003**, 56, 373. (b) Ragaini, F.; Cenini, S.; Turra, F.; Caselli, A. *Tetrahedron* **2004**, 60, 4989. (c) Coventry, D. N.; Batsanov, A. S.; Goeta, A. E.; Howard, J. A. K.; Marder, T. B. *Polyhedron* **2004**, 23, 2789. (d) Khusniyarov, M. M.; Harms, K.; Burghaus, O.; Sundermeyer, J. *Eur. J. Inorg. Chem.* **2006**, 2985. (e) Fedushkin, I. L.; Skatova, A. A.; Ketkov, S. Y.; Eremenko, O. V.; Piskunov, A. V.; Fukin, G. K. *Angew. Chem., Int. Ed.* **2007**, 46, 4302. (f) Flapper, J.; Reek, J. N. H. *Angew. Chem., Int. Ed.* **2007**, 46, 8590. (g) Fedushkin, I. L.; Makarov, V. M.; Sokolov, V. G.; Fukin, G. K. *Dalton Trans.* **2009**, 8047. (h) Li, L.; Lopes, P. S.; Rosa, V.; Figueira, C. A.; Lemos, M. A. N. D. A.; Duarte, M. T.; Avilés, T.; Gomes, P. T. *Dalton Trans.* **2012**, 41, 5144. (i) Li, L.; Lopes, P. S.; Figueira, C. A.; Gomes, C. S. B.; Duarte, M. T.; Rosa, V.; Fliedel, C.; Avilés, T.; Gomes, P. T. *Eur. J. Inorg. Chem.* **2013**, 1404.

(3) (a) Sprengers, J. W.; de Greef, M.; Duin, M. A.; Elsevier, C. J. *Eur. J. Inorg. Chem.* **2003**, 3811. (b) Kluwer, A. M.; Koblenz, T. S.; Jonischkeit, T.; Woelk, K.; Elsevier, C. J. *J. Am. Chem. Soc.* **2005**, 127, 15470. (c) Guo, H.; Zheng, Z. L.; Yu, F.; Ma, S. M.; Holuigue, A.; Tromp, D. S.; Elsevier, C. J.; Yu, Y. H. *Angew. Chem., Int. Ed.* **2006**, 45, 4997. (d) Scarel, A.; Axet, M. R.; Amoroso, F.; Ragaini, F.; Elsevier, C. J.; Holuigue, A.; Carfagna, C.; Mosca, L.; Milani, B. *Organometallics* **2008**, 27, 1486.

(4) (a) Killian, C. M.; Tempel, D. J.; Johnson, L. K.; Brookhart, M. J. *Am. Chem. Soc.* **1996**, 118, 11664. (b) Mecking, S.; Johnson, L. K.; Wang, L.; Brookhart, M. J. *Am. Chem. Soc.* **1998**, 120, 888. (c) Tempel, D. J.; Johnson, L. K.; Huff, R. L.; White, P. S.; Brookhart, M. J. *Am. Chem. Soc.* **2000**, 122, 6686. (d) Gottfried, A. C.; Brookhart, M. *Macromolecules* **2001**, 34, 1140. (e) Gottfried, A. C.; Brookhart, M. *Macromolecules* **2003**, 36, 3085. (f) Leatherman, M. D.; Svejda, S. A.; Johnson, L. K.; Brookhart, M. J. *Am. Chem. Soc.* **2003**, 125, 3068. (g) Shiotsuki, M.; White, P. S.; Brookhart, M.; Templeton, J. L. *J. Am. Chem. Soc.* **2007**, 129, 4058.

(5) (a) Cherian, A. E.; Rose, J. M.; Lobkovsky, E. B.; Coates, G. W. *J. Am. Chem. Soc.* **2005**, 127, 13770. (b) Rose, J. M.; Cherian, A. E.; Coates, G. W. *J. Am. Chem. Soc.* **2006**, 128, 4186. (c) Rose, J. M.; Deplace, F.; Lynd, N. A.; Wang, Z. G.; Hotta, A.; Lobkovsky, E. B.; Kramer, E. J.; Coates, G. W. *Macromolecules* **2008**, 41, 9548.

(6) Fedushkin, I. L.; Skatova, A. A.; Chudakova, V. A.; Cherkasov, V. K.; Fukin, G. K.; Lopatin, M. A. *Eur. J. Inorg. Chem.* **2004**, 388.

(7) Most recent review: (a) Hill, N. J.; Vargas-Baca, I.; Cowley, A. H. *Dalton Trans.* **2009**, 2, 240. Selected recent papers: (b) Fedushkin, I. L.; Nikipelov, A. S.; Lyssenko, K. A. *J. Am. Chem. Soc.* **2010**, 132, 7874. (c) Fedushkin, I. L.; Lukoyanov, A. N.; Tishkina, A. N.; Fukin, G. K.; Lyssenko, K. A.; Hummert, M. *Chem. Eur. J.* **2010**, 16, 7563. (d) Fedushkin, I. L.; Moskalev, M. V.; Lukoyanov, A. N.; Tishkina, A. N.; Baranov, E. V.; Abakumov, G. A. *Chem. Eur. J.* **2012**, 18, 11264. (e) Fedushkin, I. L.; Markina, O. V.; Lukoyanov, A. N.; Morozov, A. G.; Baranov, E. V.; Maslov, M. O.; Ketkov, S. Y. *Dalton Trans.* **2013**, 42, 7952. (f) Fedushkin, I. L.; Skatova, A. A.; Chudakova, V. A.; Fukin, G. K.; Dechert, S.; Schumann, H. *Eur. J. Inorg. Chem.* **2003**, 3336. (g) Fedyushkin, I. L.; Skatova, A. A.; Chudakova, V. A.; Cherkasov, V. K.; Dechert, S.; Schumann, H. *Russ. Chem. Bull.* **2004**, 2142.

(8) (a) Vasudevan, K.; Cowley, A. H. *Chem. Commun.* **2007**, 3464. (b) Fedushkin, I. L.; Maslova, O. V.; Baranov, E. V.; Shavyrin, A. S. *Inorg. Chem.* **2009**, 48, 2355. (c) Fedushkin, I. L.; Maslova, O. V.; Hummert, M.; Schumann, H. *Inorg. Chem.* **2010**, 49, 2901. (d) Fedushkin, I. L.; Maslova, O. V.; Morozov, A. G.; Dechert, S.; Demeshko, S.; Meyer, F. *Angew. Chem., Int. Ed.* **2012**, 51, 10584.

(9) Fedushkin, I. L.; Skatova, A. A.; Cherkasov, V. K.; Chudakova, V. A.; Dechert, S.; Hummert, M.; Schumann, H. *Chem. Eur. J.* **2003**, 9, 5778.

(10) Fedushkin, I. L.; Skatova, A. A.; Lukoyanova, A. N.; Chudakova, V. A.; Dechert, S.; Hummert, M.; Schumann, H. *Russ. Chem. Bull.* **2004**, 53, 2751.

(11) Fedushkin, I. L.; Skatova, A. A.; Fukin, G. K.; Hummert, M.; Schumann, H. *Eur. J. Inorg. Chem.* **2005**, 2332.

(12) Fedushkin, I. L.; Khvoynova, N. M.; Skatova, A. A.; Fukin, G. K. *Angew. Chem., Int. Ed.* **2003**, 42, 5223.

- (13) Fedushkin, I. L.; Morozov, A. G.; Rassadin, O. V.; Fukin, G. K. *Chem. Eur. J.* **2005**, *11*, 5749.
- (14) Fedushkin, I. L.; Makarov, V. M.; Rosenthal, E. C. E.; Fukin, G. K. *Eur. J. Inorg. Chem.* **2006**, 827.
- (15) (a) Fedushkin, I. L.; Hummert, M.; Schumann, H. *Eur. J. Inorg. Chem.* **2006**, 3266. (b) Tishkina, A. N.; Lukoyanov, A. N.; Morozov, A. G.; Fukin, G. K.; Lyssenko, K. A.; Fedushkin, I. L. *Russ. Chem. Bull., Int. Ed.* **2009**, *58*, 2250.
- (16) (a) Evans, D. A.; Cowley, A. H. *J. Am. Chem. Soc.* **2012**, *134*, 15672. (b) Evans, D. A.; Vargas-Baca, I.; Cowley, A. H. *J. Am. Chem. Soc.* **2013**, *135*, 13939.
- (17) (a) Barrett, A. G. M.; Crimmin, M. R.; Hill, M. S.; Procopiou, P. A. *Proc. R. Soc. Ser. A: Math. Phys. Eng. Sci.* **2010**, *466*, 927. (b) Harder, S. *Chem. Rev.* **2010**, *110*, 3852. (c) Arrowsmith, M.; Hill, M. S. *Alkaline Earth Chemistry: Applications in Catalysis*. In *Comprehensive Inorganic Chemistry II*; Reedijk, J., Poepelmeier, K., Eds.; Elsevier: Oxford, U.K., 2013; Vol. 1, p 1189. Homogeneous Catalysis with Organometallic Complexes of Group 2: (d) Crimmin, M. R.; Hill, M. S. *Top. Organomet. Chem.* **2013**, *45*, 191.
- (18) (a) Schlenk, W.; Schlenk, W., Jr. *Chem. Ber.* **1929**, *62*, 920. (b) Avent, A. G.; Crimmin, M. R.; Hill, M. S.; Hitchcock, P. B. *Dalton Trans.* **2005**, 278.
- (19) (a) Harder, S. *Angew. Chem., Int. Ed.* **2003**, *42*, 3430. (b) Crimmin, M. R.; Barrett, A. G. M.; Hill, M. S.; MacDougall, D.; Mahon, M. F.; Procopiou, P. A. *Dalton Trans.* **2009**, 9715.
- (20) (a) Harder, S.; Feil, F.; Knoll, K. *Angew. Chem., Int. Ed.* **2001**, *40*, 461. (b) Feil, F.; Harder, S. *Eur. J. Inorg. Chem.* **2003**, 3401. (c) Weeber, A.; Harder, S.; Brintzinger, S. H.; Knoll, K. *Organometallics* **2000**, *19*, 1325.
- (21) Liu, B.; Roisnel, T.; Carpentier, J.-F.; Sarazin, Y. *Chem. Eur. J.* **2013**, *19*, 2784.
- (22) Arrowsmith, M.; Hill, M. S.; Kociok-Köhn, G. *Organometallics* **2010**, *29*, 4203.
- (23) (a) Jantunen, K. C.; Scott, B. L.; Hay, P. J.; Gordon, J. C.; Kiplinger, J. L. *J. Am. Chem. Soc.* **2006**, *128*, 6322. (b) Masuda, J. D.; Jantunen, K. C.; Scott, B. L.; Gordon, J. C.; Kiplinger, J. L. *Organometallics* **2008**, *27*, 803.
- (24) Arrowsmith, M.; Hill, M. S.; Kociok-Köhn, G. *Organometallics* **2011**, *30*, 1291.
- (25) Crimmin, M. R.; Arrowsmith, M.; Barrett, A. G. M.; Casely, I. J.; Hill, M. S.; Procopiou, P. A. *J. Am. Chem. Soc.* **2009**, *131*, 9670.
- (26) Maaranen, J.; Hoikka, J.; Rautio, S. Patent US2007/37939 A1, 2007.
- (27) Hitchcock, P. B.; Khvostov, A. V.; Lappert, M. F. *J. Organomet. Chem.* **2002**, *663*, 263.
- (28) Crimmin, M. R.; Barrett, A. G. M.; Hill, M. S.; MacDougall, D. M.; Mahon, M. F.; Procopiou, P. A. *Chem. Eur. J.* **2008**, *14*, 11292.
- (29) Otwinowski, Z.; Minor, W. Processing of X-ray Diffraction Data Collected in Oscillation Mode. In *Methods in Enzymology: Macromolecular Crystallography, Part A*; Carter, C. W., Jr., Sweet, R. M., Eds.; Academic Press: New York, 1997; Vol. 276 pp 1307–1326.
- (30) Farrugia, L. J. *J. Appl. Crystallogr.* **1999**, *32*, 837.
- (31) Evans, W. J. *J. Alloys Compd.* **2009**, *488*, 493.
- (32) (a) Langer, J.; Al-Shboul, T. M. A.; Younis, F. M.; Görls, H.; Westerhausen, M. *Eur. J. Inorg. Chem.* **2011**, 3022. (b) Gärtner, M.; Görls, H.; Westerhausen, M. *Z. Anorg. Allg. Chem.* **2007**, *633*, 2025. (c) Crimmin, M. R.; Barrett, A. G. M.; Hill, M. S.; Hitchcock, P. B.; Procopiou, P. A. *Inorg. Chem.* **2007**, *46*, 10410. (d) Gärtner, M.; Fischer, R.; Langer, J.; Görls, H.; Walther, G.; Westerhausen, M. *Inorg. Chem.* **2007**, *46*, 5118.
- (33) (a) Crimmin, M. R.; Casely, I. J.; Hill, M. S. *J. Am. Chem. Soc.* **2005**, *127*, 2042. (b) Datta, S.; Gamer, M. T.; Roesky, P. W. *Organometallics* **2008**, *27*, 1207. (c) Arrowsmith, M.; Crimmin, M. R.; Barrett, A. G. M.; Hill, M. S.; Kociok-Köhn, G.; Procopiou, P. A. *Organometallics* **2011**, *30*, 1493.
- (34) Barrett, A. G. M.; Crimmin, M. R.; Hill, M. S.; Kociok-Köhn, G.; Lachs, J. R.; Procopiou, P. A. *Dalton Trans.* **2008**, 1292.