This article was downloaded by: [Kungliga Tekniska Hogskola] On: 06 October 2014, At: 11:31 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/gpss20

Novel and Highly Efficient Protection of Aliphatic Alcohols and Phenols with Hexamethyldisilazane in the Presence of La(NO₃)₃. 6 H₂O

Batool Akhlaghinia ^{a b}

^a Department of Chemistry, Faculty of Sciences , Ferdowsi University of Mashhad , Mashhad, Iran

^b School of Chemistry, Damghan University of Basic Sciences, Damghan, Iran Published online: 21 Sep 2009.

To cite this article: Batool Akhlaghinia (2009) Novel and Highly Efficient Protection of Aliphatic Alcohols and Phenols with Hexamethyldisilazane in the Presence of $La(NO_3)_3$. 6 H_2O , Phosphorus, Sulfur, and Silicon and the Related Elements, 184:10, 2530-2535, DOI: <u>10.1080/10426500802509616</u>

To link to this article: http://dx.doi.org/10.1080/10426500802509616

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with

primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <u>http://www.tandfonline.com/page/terms-and-conditions</u>



Novel and Highly Efficient Protection of Aliphatic Alcohols and Phenols with Hexamethyldisilazane in the Presence of $La(NO_3)_3 \cdot 6 H_2O$

Batool Akhlaghinia

Department of Chemistry, Faculty of Sciences, Ferdowsi University of Mashhad, Mashhad, Iran and School of Chemistry, Damghan University of Basic Sciences, Damghan, Iran

Aliphatic alcohols and phenols are protected with hexamethyldisilazane in the presence of lanthanum nitrate hexahydrate $(La(NO_3)_3 \cdot 6H_2O)$ in excellent yields at room temperature.

Keywords Alcohol; hexamethyldisilazane (HMDS); lanthanum nitrate hexahydrate; phenol; trimethylsilyl ether

INTRODUCTION

The role of silvl groups has already been recognized as an important part of organic chemistry from both analytical and synthetic points of view, especially as a protecting group in many syntheses of reasonable complexity.¹ Generally the formation of silvl ethers is achieved by treatment of the corresponding alcohols with silvl chlorides or silvl triflates in the presence of a base,² Li₂S,³ and sometimes a nonionic super base catalyst.⁴ However, some of these methods frequently suffer from drawbacks such as the lack of reactivity or the difficulty in removing the ammonium salts derived from the reaction of the by-produced acid and the co-bases during the silvlation reaction. 1,1,1,3,3,3-Hexamethyldisilazane (HMDS) is a stable, commercially available, and cheap reagent for trimethylsilvlation of hydroxy compounds, giving ammonia as the only byproduct. On the other hand,

Received 28 August 2008; accepted 24 September 2008.

We gratefully acknowledge the partial support of this study by Damghan University Research Council.

Address correspondence to Batool Akhlaghinia, Department of Chemistry, Faculty of Sciences, Ferdowsi University of Mashhad, Mashhad, Iran. E-mail: akhlaghinia@um.ac.ir or Department of Chemistry, Damghan University of Basic Sciences, Damghan, Iran. E-mail: b_akhlaghinia@dubs.ac.ir

because of easiness in handling and neutrality of this silylating reagent using HMDS does not need special precautions. However, the low silylating power of 1,1,1,3,3,3-hexamethyldisilazane is a main drawback for its application, which needs forceful conditions and long reaction times in many instances. A variety of catalysts such as $(CH_3)_3SiCl,^5$ $ZnCl_2,^6$ complexes of metal chlorides⁷ with nitrogen ligands, zirconium sulfophenyl phosphonate,⁸ K-10 montmorillonite,⁹ and a special type of catalyst with the general formula X¹-NH-X², in which at least one of the groups X¹ and X² is electron-withdrawing such as an ester or amide group, has also been reported to promote the silylation of a wide range of functional groups using HMDS.¹⁰

Here we report the use of an efficient and cheap system for the trimethylsilylation of aliphatic alcohols and phenols.

RESULTS AND DISCUSSION

Recently we have reported the trimethylsilylation of alcohols and phenols with HMDS in the presence of $Cu(2,3-tmtppa)^{4+11}$, $CuSO_4 \cdot 5 H_2O$, and anhydrous FeCl₃.¹² In continuation of this study we are especially interested in exploring the potential use of a neutral catalyst. Here according to the previous catalytic effect of $La(NO_3)_3 \cdot 6 H_2O$,¹³ we describe a new protocol for the mild and rapid trimethylsilylation of a wide variety of aliphatic alcohols and phenols using HMDS and a catalytic amount of $La(NO_3)_3 \cdot 6 H_2O$. First, the protection of benzyl alcohol under the catalytic effect of $La(NO_3)_3 \cdot 6 H_2O$ was studied. Benzyl trimethylsilyl ether was produced immediately in quantitative yield. Employing a 1:1 ratio in acetonitrile at room temperature gave the best results and produced the respective trimethylsilyl ethers in quantitative yields (Table I). We then applied these conditions for the protection of structurally different alcohols and phenols (Scheme 1).

As shown in Table I, this method is very suitable for the conversion of primary, secondary, tertiary, and benzylic alcohols as well as aryl alcohols to the corresponding trimethylsilyl ethers.

ROH + HMDS
$$\xrightarrow{\text{La (NO_3)_3.6 H_2O}}$$
 ROSiMe₃

R: primary, secondary, tertiary alkyl and phenyl

SCHEME 1

Substrate	$\operatorname{Conversion} \%^a$	Isolated yield%
	100	98
	100	97
ОН		
ОН	100	97
	100	98
OH	100	95
	100	96
OH OH	100	50
ОН	100	98
<i>С</i>	100	98
ОН	100	98
	100	07
ОН	100	97
	100	00
ОН	100	90
CI CI	100	98
OH OH		
O ₂ N	100	96
ОН 	100	95
 ОН		
	100	95
U OH	100	<u> </u>
	100	98
Óн	100	07
< >-он	100	51

TABLE I Trimethylsilylation of Aliphatic Alcohols and Phenols Catalyzed by La $(NO_3)_3\cdot\, 6H_2O$ in Dry CH_3CN

(Continued on next page)

Substrate	$Conversion\%^a$	Isolated yield%
	100	97
ОН		
∨ он∣	85	80
	100	90
ОН	80	73
ОН	90	85
ОН	80	78
ОН	95	90
ОН	90	88
ОН	85	80
СІ	60	58
O ₂ N OH	70	67
ОН	50	46

TABLE I Trimethylsilylation of Aliphatic Alcohols and Phenols Catalyzed by La (NO₃)₃. 6H₂O in Dry CH₃CN (*Continued*)

^aGC yield using *n*-octane as internal standard.

To estimate the selectivity of this new method, we studied the reaction of some binary mixtures of substrates. Excellent selectivity was observed in the conversion of primary alcohols to the corresponding trimethylsilyl ethers in the presence of secondary and tertiary alcohols

Binary Mixture	Products	Yield $(\%)^a$
ОН	OSiMe ₃	100
		98
ОН	ОН	
ОН	OSiMe ₃	100
\sim		97
OH OH	OH OSiMe ₃	100
ОН	ОН	100
		100
ОН	OSiMe ₃	
ОН	ОН	100
ОН	OSiMe ₃	100
OH	ОН	100

TABLE II Selective Trimethylsilyl Ether Formationfrom Different Binary Mixtures

 a GC yield using *n*-octane as internal standard.

(Table II). Benzyl alcohol is selectively protected with this reagent in the presence of phenol (Table II).

In this article, we have introduced La $(NO_3)_3$ · 6 H₂O as a novel, efficient, and convenient solid acid catalyst for the trimethylsilylation of hydroxyl groups. The new catalyst is reusable and is easily separated from the reaction mixture. Workup and isolation of the product by this protocol are also straight forward.

EXPERIMENTAL

Chemicals were obtained from Merck and Fluka. FT-IR spectra were recorded with a Perkin Elmer RXI spectrophotometer. ¹H NMR spectra were recorded with a Bruker Avance DPX 250 MHz instrument. The products were purified by column chromatography and the purity of the products was determined by GC on a Shimadzu GC 10-A instrument using *n*-octane as internal standard or by thin layer chromatography using silica gel polygram on SIL G/UV 254 plates.

General Procedure for Protection of Alcohols

To a flask containing $La(NO_3)_3 \cdot 6 H_2O$ (0.004 g, 0.01 mmol) in dry acetonitrile (3–5 mL), HMDS (0.576 g, 1 mmol) was added under stirring at room temperature. Subsequently, the alcohol (1 mmol) was added. GC analysis showed that the reaction was completed immediately. The solvent was evaporated under reduced pressure. Column chromatography of the crude product on silica gel using *n*-hexane as eluent gave the corresponding trimethylsilyl ether in quantitative yield.

REFERENCES

- (a) M. Lalonde and T. H. Chan, Synthesis, 817 (1985); (b) T. W. Greene and P. G. M. Wuts, Protective Groups in Organic Synthesis, 2nd ed. (Wiley, New York, 1991); (c) P. J. Kocienski, In Protective Groups, R. Enders, R. Noyori, and B. M. Trost, eds. (Thieme Stuttgart, 1994).
- [2] S. K. Ghaudhary and O. Hernandez, Tetrahedron Lett., 20, 99 (1979).
- [3] G. A. Olah, B. G. B. Gupta, S. C. Narang, and R. Malhotra, J. Org. Chem., 44, 4272 (1979).
- [4] (a) B. A. D. Sa, D. Mc Leod, and J. G. Verkade, J. Org. Chem., 62, 5057 (1997); (b) B.
 A. D. Sa and J. G. Verkade, J. Am. Chem. Soc., 118, 12832 (1996).
- [5] (a) S. H. Langer, S. Connell, and J. Wender, J. Org. Chem., 23, 50 (1958); (b) P. Gauttret, S. El-Ghamarti, A. Legrand, D. Coutrier, and B. Rigo, Synth. Commun., 26, 707 (1996).
- [6] H. Firouzabadi and B. Karimi, Synth. Commun., 23, 1633 (1993).
- [7] (a) H. Firouzabadi, A. R. Sardarian, Z. Khayat, and B. Karimi, Synth. Commun., 27, 2709 (1997); (b) A. Bongini, G. Cardillo, M. Orena, and S. Sandri, Synthesis, 618 (1979).
- [8] M. Curini, F. Epifano, M. C. Marcotullio, O. Rosati, and U. Constantino, Synth. Commun., 29, 541 (1999).
- [9] Z. H. Zhang, T. S. Li, F. Yang, and C. G. Fu, Synth. Commun., 28, 3105 (1998).
- [10] C. A. Bruynes and T. K. Jurriens, J. Org. Chem., 47, 3966 (1982).
- [11] B. Akhlaghinia, M. Asadi, E. Safaei, and M. Heydarpoor, *Phosphorus, Sulfur, and Silicon*, 179, 2099 (2004).
- [12] (a) B. Akhlaghinia and S. Tavakoli, Synthesis, 1755 (2005); (b) B. Akhlaghinia, Phosphorus, Sulfur, and Silicon, 182, 687 (2007).
- [13] T. S. Reddy, M. Narasimhulu, N. Suryakiran, K. C. Mahesh, K. Ashalatha, and Y. Venkateswarlu, *Tetrahedron Lett.*, 47, 6825 (2006).