

Mercuration of Schiff Bases of Substituted Benzylideneanilines


Kulling Ding,[†] Yangjie Wu,^{*†} Hongwen Hu,[‡] Lianfang Shen,[§] and Xin Wang^{||}

Department of Chemistry, Zhengzhou University, Zhengzhou 450052, PRC, Department of Chemistry, Nanjing University, Nanjing 210008, PRC, Laboratory of Magnetic Resonance and Atomic and Molecular Physics, Wuhan Institute of Physics, Academia Sinica, Wuhan 430071, PRC, and National Applied Laboratory of Organic Chemistry, Lanzhou University, Lanzhou 730000, PRC

Received June 18, 1992

As part of an effort to investigate substituent effects on the intramolecular coordination between N and Hg, the mercuration of 36 substituted benzylideneanilines was studied. The structure characterization of the products by IR, ¹H NMR, ¹³C NMR, and MS indicates that for all of the reactions examined, the mercury is directed to the ortho position of the N-phenyl ring or the para position of the N-phenyl ring when these sites are not occupied by a substituent. The position of the HgCl group in the mercurated product of N-(4-nitrobenzylidene)-β-naphthylamine has been confirmed by single-crystal X-ray determination, which also provided circumstantial evidence for the existence of the N→Hg intramolecular coordination with a four-membered ring. The possible mechanism of the reaction was proposed, in which the mercuration at the ortho position of the N-phenyl ring was facilitated by the imino moiety upon formation of a coordination complex with Hg(OAc)₂ in the first step, followed by a subsequent electrophilic substitution at the ortho position of the N-phenyl ring. This reaction is different from the metalation of benzylideneanilines by transition metals, in which the metal atom is usually directed to the ortho position of the C-phenyl ring, and provides a new example of the so-called "cyclometalation" reaction.

Over the last two decades, there has been a great deal of interest in the ortho-metalation reaction of substituted aromatic compounds in which nitrogen-containing groups act as electron donor ligands.¹ Numerous relevant organometallic intramolecular coordination compounds have been synthesized and some have been found to be useful in organic syntheses.² Until now, attention has been focused on the reactions of transition metals (e.g. Pd, Pt etc.). Compared to the transition metals, the coordinating ability of the nontransition metals (e.g. Hg, Sn, etc.) is weak. The direct orthomercuration reactions are primarily restricted to the reaction of substituted azobenzenes³ and phenylhydrazones,⁴ in which nitrogen donor ligands facilitate the formation of the C-Hg σ-bond. Work on mercuration of Schiff bases of the substituted benzylideneanilines was first published by Singh et al.,⁵ where two mercurated derivatives were reported.



In their molecules, mercury is considered to be bonded to the ortho position of the C-phenyl ring. Because of the weak coordinating ability of mercury and specific electron population distribution of benzylideneaniline,⁶ the mercury is very likely directed to the ortho position of the N-phenyl ring and, as a result, may form an intramolecular coordination via a four-membered ring. In our opinion, the structure of the mercurated Schiff bases reported by Singh et al. may be incorrect. It is known that in arylmercury compounds ortho substituents with p- or π-electrons have substantial effects on reaction rates and spectral properties.⁷ As part of an effort to investigate substituent effects on the intramolecular coordination between N and Hg, mercuration of some substituted benzylideneanilines has been studied. The present paper reports the results of the direct mercuration of substituted benzylideneanilines,

effects of substituents, solvents, and electrophiles on this reaction, and the possible mechanism of the reaction.

Results and Discussion

We began our investigation with the Schiff bases (1a-7a) since the product structure can be easily identified by ¹H NMR spectroscopy due to the presence of the strong electron-withdrawing NO₂ group in the C-phenyl ring. The mercuric chlorides of the corresponding Schiff bases (1b-7b) were synthesized by in situ reaction of Schiff bases with mercuric acetate (1:1 molar ratio) in refluxing dry methanol for 4 h, followed by the addition of lithium chloride. Reactions were carried out on a 5-mmol scale, and the initial concentration of each substrate was kept the same (~0.1 M). For all the new compounds prepared, compositions and structures were determined and confirmed by elemental analysis, IR, NMR, and MS.

Characterization of Products. ¹H NMR. The elemental analyses and IR spectra indicated that all the mercurated derivatives are composed of one Schiff base unit and one HgCl group except for 3b and 10b where two

- (1) (a) Dehand, J.; Pfeffer, M. *Coord. Chem. Rev.* 1976, 18, 327-357. (b) Bruce, M. I. *Angew. Chem., Int. Ed. Engl.* 1977, 16, 73-86. (c) Omae, I. *Chem. Rev.* 1979, 79, 287-321.
- (2) Ryabov, A. D. *Synthesis* 1985, 233-252.
- (3) (a) Roling, P. V.; Dill, J. L. *J. Organomet. Chem.* 1974, 69, C33-C35. (b) Roling, P. V.; Kirt, D. D.; Dill, J. L.; Hall, S.; Hollstrom, C. J. *Organomet. Chem.* 1976, 116, 39-53. (c) Roling, P. V.; Picking, R. A. *J. Organomet. Chem.* 1976, 122, 289-293.
- (4) (a) Butler, R. N.; King, W. B. *Chem. Ind.* 1975, 647-648. (b) Butler, R. N.; King, W. B. *J. Chem. Soc., Perkin Trans. 1* 1976, 986-989. (c) Butler, R. N.; O'Donohue, A. M. *Tetrahedron Lett.* 1978, 275-278. (d) Butler, R. N.; O'Donohue, A. M. *J. Chem. Soc., Perkin Trans. 2* 1979, 1387-1390.
- (5) Singh, H. B.; McWhinnie, W. R. *J. Chem. Soc., Dalton Trans.* 1985, 821-825.
- (6) Smith, W. F. *Tetrahedron* 1963, 19, 445-454.
- (7) (a) Wu, Y. J.; Chen, Z. D.; Zhang, X. P.; Hu, H. W. *Youji Huaxue (Organic Chemistry)* 1982, 5, 331-339; *Chem. Abstr.* 1983, 98, 72315c. (b) Wu, Y. J.; Chen, Z. D.; Zhang, X. P.; Zhang, Z. W.; Liu, Y. A.; Ni, J. X. *Chem. J. Chin. Univ.* 1983, 4, 335-341; *Chem. Abstr.* 1983, 99, 52914q. (c) Chen, Z. D.; Wu, Y. J.; Zhang, X. P.; Chen, R. F.; Hu, H. W. *Acta Chim. Sin. (Chin. Ed.)* 1983, 41, 57-65; *Chem. Abstr.* 1983, 98, 160863x. (d) Chen, Z. D.; Wu, Y. J.; Zhang, X. P.; Su, Q. Y.; Yi, M. P.; Hu, H. W. *Acta Chim. Sin. (Chin. Ed.)* 1984, 42, 792-795; *Chem. Abstr.* 1984, 101, 192108q. (e) Pan, Z. H.; Wei, X. C.; Wu, Y. J.; Chen, Z. D.; Wang, Y. L.; Hu, H. W. *Acta Chim. Sin. (Chin. Ed.)* 1985, 43, 801-805; *Chem. Abstr.* 1986, 104, 43559x.


* To whom correspondence should be addressed.

[†] Zhengzhou University.

[‡] Nanjing University.

[§] Wuhan Institute of Physics.

^{||} Lanzhou University.



HgCl groups are present. The position of the mercury atom was determined by ¹H NMR. As expected, the aromatic region of the spectrum for each compound contains a group of left-right symmetrical multiplet peaks (4 H) characterizing the four-spin AA'BB' system at lower field due to the electron-withdrawing effect of the nitro group. Furthermore, it was found that the substituents X have no effect on the chemical shift values of these protons. So, these signals can only be assigned to the C-phenyl protons (H2, H3). The remaining two doublets (2 H) and a doublet of doublets (1 H) which belong to the three-spin ABX system at higher field correspond to the three N-phenyl protons, for which the δ values vary considerably with different substituents X. The assignment of ¹H NMR spectra (Table I) for the Schiff base type organomercury compounds 1b–7b showed that the mercury is directed to the ortho position of the N-phenyl ring rather than the ortho position of the C-phenyl ring. Further investigation of the ¹H NMR spectra of mercurated products 8b–13b gave the same conclusion. In the cases of 3a and 10a, the mercury atom is also directed to the para position of the N-phenyl ring, in addition to the ortho position. This can be observed from the spectral data of 3b and 10b.

The NOE spectrum of compound 1b was obtained by irradiation of the methyl proton at 2.42 ppm and the methine proton at 9.00 ppm, respectively. The significant enhancements of the signals at 7.28 (10%) and 7.53 ppm (10%) identify these as protons H5 and H6, respectively, closest to the methyl being irradiated, and correspondingly, the enhancement of the signals at 8.25 ppm (12.5%) identified these peaks as due to proton H2, closest to the

Table I. ¹H NMR Data for Compounds 1b–13b^a

compd	1b–7b				8b–13b		
	H-1	H-2	H-3	H-4	H-5	H-6	H-7
1b	9.00	8.25	8.46	7.50	7.28	7.53	
2b	8.99	8.22	8.45	7.64	7.03	7.35	
3b	8.95	8.22	8.45	7.50	7.50	7.68	
4b ^b							
5b	8.98	8.25	8.47	7.53	7.60	7.89	
6b	8.99	8.26	8.49	7.30	7.75	8.05	
7b	9.30	8.34	8.50				
8b	8.84	8.00	7.60	7.61	7.44	7.24	7.49
9b	8.84	8.03	7.63	7.64	7.57	7.02	7.42
10b	8.83	8.00	7.63	7.64	7.41	7.49	7.60
1b	8.84	8.04	7.64	7.64	7.49	7.48	7.74
12b	8.84	8.04	7.62	7.63	7.45	7.61	7.87
13b	8.82	8.03	7.64	7.64	7.31	7.76	8.05

^a All data recorded in DMSO-*d*₆ at 20 °C. ^b This compound cannot be dissolved in DMSO-*d*₆.




Figure 1. Drawing of structure of 7b.

methine proton being irradiated. So, the results of the NOE experiment support the assignment by ¹H NMR.

¹³C NMR. It was observed from the ¹³C{¹H} NMR spectra of the aromatic carbons of organomercurials 8b–13b, that the chemical shift values of five tertiary and one quaternary carbons were virtually not affected by different substituents, whereas the δ values of the other three tertiary and three quaternary carbons varied markedly with different substituents. Table II lists the assigned ¹³C NMR spectra for 8b–13b. An examination of the variations of the chemical shifts of C6, C7, and C11 with different substituents shows that electron-withdrawing groups cause a downfield shift and that electron-donating groups cause an upfield shift, respectively. The chemical shifts of C6 and C7 exhibit a good linear correlation with the Brown σ_p⁺ and σ_m⁺ constants, respectively. Thus by regression analyses, we have the following equations:


$$\delta(^{13}\text{C}6) = 153.1 + 7.78\sigma_p^+ \quad r = 0.989 \quad (1)$$

$$\delta(^{13}\text{C}7) = 117.2 + 5.40\sigma_m^+ \quad r = 0.940 \quad (2)$$

There is no linear correlation between δ(¹³C11) and σ_m⁺, probably due to the N→Hg intramolecular coordination which affects the electron distribution on the C11 atom. The δ values of the methine carbon (C1) fit the Hammett constants reasonably well. The following relationship was obtained by linear regression analysis:

$$\delta(^{13}\text{C}1) = 159.0 + 6.27\sigma_p \quad r = 0.971 \quad (3)$$


MS. For the mercuration of 1a–13a, if the mercury atom was directed either to the ortho positions of the C-phenyl ring (C3) or of the N-phenyl ring (C7), the cor-

Table II. $^{13}\text{C}\{^1\text{H}\}$ NMR Data for Compounds 8b-13b^a



compd	C-1	C-2	C-3	C-4	C-5	C-6	C-7	C-8	C-9	C-10	C-11
8b	158.5	135.5	128.8	128.4	131.3	151.5	116.8	129.7	136.7	137.1	151.4
9b	156.9	136.3	128.8	128.2	131.1	146.7	117.4	121.6	157.8	114.3	154.0
10b	159.6	136.0	128.8	128.5	131.5	153.5	117.1	137.0	151.0	144.1	149.8
11b	160.3	135.9	128.9	128.7	131.7	153.7	118.8	128.7	130.6	135.9	152.9
12b	160.4	135.9	128.9	128.6	131.7	154.1	119.3	131.5	129.5	138.7	153.6
13b	160.3	135.9	128.9	128.7	131.7	154.3	119.8	137.3	92.5	144.7	153.7

^a All data recorded in DMSO-*d*₆ at 20 °C.

responding different fragment ions should be observed in the mass spectra.




It was found that the ions *m/e* M - 149 and M - 104 are present, but the ions *m/e* 385 and 340 are absent in the spectra of 1b-7b and 8b-13b, respectively. Therefore, mercury attack on the ortho position of the C-phenyl ring can be eliminated. In the case of 6b, the cation *m/e* M - 149 is not formed, probably due to the high melting point and the weak C-I bond strength of 6b. The fragmentation behavior observed for 1b-13b can be expressed as follows:



As expected, the mercury is directed into the ortho position of the *N*-phenyl ring.

In order to reexamine the results reported by Singh et al., the mass spectra of mercurated derivatives (14b, 15b) of (3-methoxybenzylidene)aniline and (2,4-dinitrobenzylidene)aniline were recorded. The appearance of an *m/e* 313 peak indicated that the mercury atom was directed into the *N*-phenyl ring. Further examination of the $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum of 14b showed 13 peaks in the aromatic region. This also led us to believe that the mercury was directed to the ortho position of the *N*-phenyl ring.




X-ray Crystallographic Results for 7b. In order to confirm the position of HgCl on the naphthalene ring for compound 7b and to provide at least circumstantial evidence for the existence of N→Hg intramolecular coordination in the mercurated Schiff bases, a single-crystal X-ray structure determination of 7b was carried out. Crystals of 7b suitable for X-ray analysis were obtained by slow recrystallization from 1,4-dioxane. Elemental analysis indicated that the crystals contain 0.5 molecule of crystalline 1,4-dioxane, which was also supported by thermoanalysis. The loss of weight for TG at 141 °C on the DTA curve is about 7.9%, which is equivalent to 0.5 mol of 1,4-dioxane. Another endothermal peak at 273 °C represents the melting of 7b.


The crystal structure of 7b is shown in Figure 1. The structure is thus confirmed as α-(chloromercurio)-*N*-(4-nitrobenzylidene)-β-naphthylamine. In other words, the HgCl is situated at the α-position of the β-naphthylamine unit. The oxygen atoms in 1,4-dioxane do not coordinate with the mercury atom and exist in the unit cell relying on van der Waals forces. The lengths of Hg-C10 [2.099 (9) Å] and Hg-Cl [2.303 (2) Å] are similar to those found in other organomercuric chlorides.⁸ The bond angles of N1-C1-C10 [114.3 (8)°] and C1-C10-Hg [117.2 (2)°] are less than 120°, which may arise from a N→Hg interaction.

Structural Effect on the Reaction. Further investigation of the structural effects of the substrates (16a-26a) on mercuration revealed that, whether the C-phenyl ring bears an electron-donating group or an electron-withdrawing group, the mercury in the product molecules (16b-26b) is attached to the ortho position of the *N*-phenyl ring, without exception.


(8) Furmanova, N. G.; Kuz'Mina, L. G.; Struchkov, Yu. T. In *Organometallic Chemistry Review*; Seyferth, D., Davies, A. G., Fischer, E. O., Normant, J. F., Reutov, O. A., Eds.; Journal of Organometallic Chemistry Library; Elsevier Scientific Publishing Co.: New York, 1980; Vol. 9, pp 153-187.




16a, R = *p*-NMe₂; 17a, R = *p*-MeO; 18a, R = 3,4-(MeO)₂;
19a, R = *o*-MeO; 20a, R = *p*-Cl; 21a, R = *m*-Cl; 22a, R = *o*-Cl;
23a, R = *p*-Br; 24a, R = *m*-Br; 25a, R = *m*-NO₂; 26a, R = *o*-NO₂




For the purpose of comparison with 7a, mercuration of 27a bearing a strong electron-donating NMe₂ group on the C-phenyl ring was carried out. It was found that the reaction took place at the same position as in 7a.



We have tried to prepare 4'-(ClHg)-benzo-15-crown-5 from 4'-(H₂N)-benzo-15-crown-5 according to the literature method.⁹ Instead, we isolated an unexpected novel complex with the composition [Na(B15C5)]⁺₂[Hg₂Cl₈]²⁻·H₂O.¹⁰ Nonetheless, mercuration of the derivatives of benzo-15-crown-5 (28a-31a) revealed that the mercury was also






28a, X = CH₃; 29a, X = CH₃O; 30a, X = Cl; 31a, X = Br




directed into the ortho position of the *N*-phenyl ring as usual, without the formation of a coordination complex of Hg(OAc)₂ with the benzo-15-crown-5 unit.

As mentioned previously, in the mercuration of 3a and 10a, dimercurials (3b and 10b) were obtained as the main products. In order to understand the ortho substituent effect on this reaction, five ortho-substituted Schiff bases (32a-36a) were used in the mercuration reaction. When the ortho position of the C-phenyl ring was not occupied by a substituent and while the ortho position of the *N*-phenyl ring was substituted by a methyl group, the Schiff bases (32a, 33a) were mercured in the para position of the *N*-phenyl ring; and if the ortho position of the *N*-phenyl ring was substituted by a methoxy group, the substrate (34a) was mercured in both the ortho position and the para position of the *N*-phenyl ring. When the ortho position of C-phenyl ring was occupied by the nitro




group (35a, 36a), the reaction took place in both the ortho position and the para position of the *N*-phenyl ring, no matter whether the ortho position of the *N*-phenyl was occupied by the methyl group or not.

The regioselectivity of the reaction can be interpreted in terms of the electron distribution and conformation of the substrates. In the cases of 3a, 10a, and 35a, mercuric acetate can attack both the ortho and para positions of the *N*-phenyl ring, because of the higher electron densities at those sites.⁶



In the cases of 32a and 33a, however, the stable conformation is probably I rather than II, because electron repulsion from the shorter distance between the methine




proton and the *o*-methyl protons disfavors II. Thus the *o*-methyl group may block coordination of the imino nitrogen atom with mercuric acetate and eliminate the preference for attack at the ortho position of the *N*-phenyl



(9) Hu, H. W.; Ni, J. S.; Kao, T. Y. *Acta Chim. Sin. (Chin. Ed.)* 1979, 37, 9-13; *Chem. Abstr.* 1979, 91, 20645w.

(10) Tao, J. C.; Ding, K. L.; Wu, Y. J.; Zhu, Y. *Chin. J. Chem.* 1992, 10, 507-512.

ring. If the *o*-methyl group in **33a** is replaced by a methoxy group, the preferred conformation of molecule **34a** is probably the same as in **III**, due to the longer distance




between methine proton and methoxy oxygen. When the ortho position of the *C*-phenyl ring is substituted by a nitro group as in **36a**, the molecule will prefer to adopt conformation **IV** that relieves the possible repulsion between



the nitro and methyl groups in the case of **V**. Therefore, the conformations of **34a** and **36a** have no significant effect on the coordination between N and Hg, and as a result, the mercury can be directed to the para position as well as to the ortho position of the *N*-phenyl ring.

Effects of Changing the Electrophile. When **1a** was treated with HgCl_2 in methanol and refluxed for 24 h, no reaction was observed. Perchloric acid, which can be used as a catalyst to accelerate the rate of aromatic mercuration,¹¹ has been successfully applied to the mercuration of azobenzene and its derivatives.^{3b} Thus, an attempt was made to increase the rate of mercuration of Schiff base by adding a small amount of perchloric acid to the reaction system. Unfortunately, the color of the reaction mixture was changed from yellow to dark brown immediately upon the addition of perchloric acid, which was probably caused by oxidation of the Schiff bases by perchloric acid. In a further investigation of the effect of the electrophilic species on the reaction, $\text{Hg}(\text{OAc})_2 \cdot \text{BTMAC}$ (benzyltrimethylammonium chloride) was used as an electrophile instead of $\text{Hg}(\text{OAc})_2$. The reaction proceeded rather slowly. When **1a** was treated with $\text{Hg}(\text{OAc})_2 \cdot \text{BTMAC}$ in methanol and refluxed for 24 h, the yield of reaction (without the addition of LiCl) was 75.8%, lower than that when $\text{Hg}(\text{OAc})_2$ was used as the mercurating agent (refluxed in methanol for 4 h, yield 84.2%). In this reaction,




the real electrophile is probably ClHgOAc , and the lower yield of the reaction may arise from its reduced coordinating ability to nitrogen and weakened electrophilicity for attacking at the ortho position of the *N*-phenyl ring of **1a**. Therefore, this mercurating agent may be used for the preparation of para-mercurated Schiff bases. For ex-

Table III. Solvent Effect on the Mercuration of **1a**

	solvents					
	CH_3OH	$\text{C}_2\text{H}_5\text{OH}$	$\text{C}_4\text{H}_9\text{O}_2$	$\text{CH}_3\text{OC}_2\text{H}_5$ H_2OH	CH_2Cl_2	CH_3COOH
time (h)	4	4	4	4	4	24
temp ($^\circ\text{C}$)	65	78	101	125	40	118
yield (%)	84.2	88.4	84.2	96.8	trace	0

ample, it reacts with 4'- $\text{O}_2\text{NC}_6\text{H}_4\text{CH}=\text{NC}_6\text{H}_5$ (**3a**) to give, as expected, the para-mercurated derivative **37b** as the major product.



Solvent Effects on the Reaction. An examination of solvent effects on the mercuration reaction of **1a** (Table III) revealed that those which solvate HOAc better favor the reaction whereas those with weaker solvation ability inhibit the reaction. In HOAc alone, no reaction occurs. This suggests that the last step of the reaction, the conversion from σ -complex to the product, may be the rate-determining step, consistent with the general mechanism of aromatic electrophilic mercuration.¹¹


Mechanism of the Reaction. Roling and co-workers reported a mercuration mechanism for azobenzene, in which the mercury is directed to the ortho position by coordinating with the azo nitrogen, followed by a subsequent electrophilic substitution.^{3b} The mercuration of substituted phenylhydrazones was facilitated by a favorable attack of the *n*-electrons of the hydrazone imino moiety.^{4d} The results obtained in this study of the mercuration reaction of Schiff bases of substituted benzyldenenanilines lead us to conclude that in this reaction the mercury is directed to the ortho position of the *N*-phenyl ring by coordinating with the nitrogen atom followed by a subsequent electrophilic substitution (Scheme I). The mechanism embodies four separate steps: (I) the coordination complex formation between $\text{Hg}(\text{OAc})_2$ and the imino moiety, (II) the formation of a π -complex with the *N*-phenyl ring, (III) the conversion of the π -complex to a σ -complex, and (IV) the loss of HOAc from the σ -complex and the formation of the product. The coordination of imino nitrogen with mercury is present in the whole process of the ortho-mercuration.

Support of mercury coordination (step I) is found in the 1:1 complex formed between mercuric chloride and either 2-(phenylimino)ethyl butyrate¹² or *trans*- $\text{PtCl}(\text{PEt}_3)_2 \cdot (\text{CH}=\text{NC}_6\text{H}_4\text{-}p\text{-CH}_3)$.¹³ It is interesting to note that the mercuration of the second ortho position was not observed, either when 2 mol equiv of mercuric acetate was used with **1a** or when **1b** was treated separately with an additional 1 mol equiv of $\text{Hg}(\text{OAc})_2$. This would be expected for involvement of the *n*-electrons in the formation of the ortho-mercurated derivative since they occupy the frontier orbital of the imino moiety and thus are likely to coordinate with the mercury atom in $\text{Hg}(\text{OAc})_2$. This can also be illustrated on the basis of regioselectivity of mercuration for **32a**–**36a**. The evidence of the π -complex formation in the mercuration reaction has not been reported, but the intramolecular mercury–arene coordination is conceivable.


(11) Bamford, C. H.; Tipper, C. F. H. *Comprehensive Chemical Kinetics*; Elsevier Publishing Co.: New York, 1972; Vol. 13, pp 186–194.

(12) Thakob, V. M.; Shau, R. C. *J. Indian Chem. Soc.* 1949, 26, 251–253.

(13) Bonati, F.; Clark, H. C. *Can. J. Chem.* 1979, 57, 483–486.

Scheme I. Probable Mechanism for Mercuration of Schiff Bases

For example, ^{199}Hg NMR studies on γ -(arylpropyl)mercury compounds have disclosed the presence of mercury–arene coordination.¹⁴ It is well-known that the ground-state molecule of azobenzene is planar. In contrast, the molecular conformation of benzylideneaniline is different from that of azobenzene: the twist angles of the *N*-phenyl ring and the *C*-phenyl out of the $\text{C}=\text{N}=\text{C}-\text{C}$ plane are 55.2° and 10.3° , respectively.¹⁵



Therefore, benzylideneaniline is more favorable to the formation of π -complex than azobenzene in terms of the ground-state conformation of the molecules.

The last step of the reaction (i.e. the loss of HOAc and the formation of cyclomercurated products) is probably the rate-determining step, based on solvent effects. Further studies on the mechanism are still in progress.¹⁶ In the cases of **3a**, **10a**, and **32a–36a**, formation of the para-mercuration products is possible through an uncomplexed mechanism.

Conclusions

The mercuration of Schiff bases of 36 substituted benzylideneanilines studied to date gives different results than those reported by Singh et al.⁵ Characterization of the structures of the products by IR, ^1H NMR, ^{13}C NMR, and

MS indicates that for all of the reactions, the mercury is directed to the ortho position of the *N*-phenyl ring or the para position of the *N*-phenyl ring when it is not occupied by a substituent.

The position of the HgCl group in compound **7b** is confirmed by a single-crystal molecular structure determination, which also provided circumstantial evidence for the existence of an $\text{N}\rightarrow\text{Hg}$ intramolecular coordination bond with a four-membered ring in the Schiff base type arylmercury compound. On the basis of this structural information and the relevant literature a probable mechanism is proposed, in which the mercuration at the ortho position of the *N*-phenyl ring is facilitated by the imino moiety which forms a coordination complex with $\text{Hg}(\text{OAc})_2$ in the first step, followed by a subsequent electrophilic substitution at the ortho position of the *N*-phenyl ring. This reaction is different from the metalation of benzylideneaniline by transition metals, in which the metal atom is usually directed to the ortho position of a *C*-phenyl ring,¹ and provides a new example for the so-called “cyclo-metalation” reaction.

Experimental Section

All reactions were performed in reagent grade solvents. The substituted benzylideneanilines were prepared following a standard procedure as described in the literature.¹⁷ Melting points were measured on a WC-1 apparatus and are uncorrected. Elemental analyses were determined with a Carlo-Erba 1106. ^1H NMR and ^{13}C NMR were recorded on a Varian XL-200 or Bruker AC-80 and are referenced to the residual solvent resonance ($\text{DMSO}-d_6$, ^1H NMR at 2.60 ppm and ^{13}C NMR at 39.7 ppm). Infrared spectra were recorded on a Perkin-Elmer FT-1750 spectrometer. EI mass spectra were obtained on a VG-ZAB-MS spectrometer, operated at 20 or 70 eV with a source temperature of ca. 230°C . Samples were introduced by means of a direct insertion probe. DAT and TG were measured on a RIGAKU thermal analyzer.

General Procedure for the Workup of the Mercuration Reaction. In a round-bottom flask equipped with a condenser, was placed equal amounts (moles) of the Schiff base and mercuric acetate (enough reagent grade methanol was added to make the concentration of the Schiff base $\sim 0.1\text{ M}$). A typical run is 1.25 g (5 mmol) of *N*-(4-nitrobenzylidene)-4-toluidine (**1a**), 1.6 g (5 mmol) of mercuric acetate, and 50 mL of methanol. This mixture was stirred and refluxed for 4 h. At the end of the reaction, the mixture was allowed to cool down to room temperature, the lithium chloride (typically 0.50 g or 12 mmol dissolved in hot methanol) was added, and the resulting thick mixture was stirred for about 10 min. The resulting precipitate was filtered off, washed with ether, and dried in vacuo. Crystallization from chloroform or other solvents gave crystals of the corresponding mercuration derivatives. The compounds prepared by this procedure are as follows.

2-(Chloromercurio)-*N*-(4-nitrobenzylidene)-4-toluidine (1b). The filtered precipitate was crystallized from chloroform to give 2.0 g (84%) of **1b** as yellow needles: mp $265.5\text{--}266.5^\circ\text{C}$; ^1H NMR (200 MHz, $\text{DMSO}-d_6$) δ 9.00 (s, 1 H), 8.46 (d, 2 H, $J = 8.6\text{ Hz}$), 8.25 (d, 2 H, $J = 8.2\text{ Hz}$), 7.53 (s, 1 H), 7.50 (d, 1 H, $J = 8.8\text{ Hz}$), 7.28 (d, 1 H, $J = 8.0\text{ Hz}$), 2.41 (s, 3 H); ^{13}C NMR (50.3 MHz, $\text{DMSO}-d_6$) δ 156.7, 152.3, 150.7, 148.7, 141.7, 137.3, 136.5, 129.6, 129.2, 124.0, 116.9, 20.0; IR (KBr) 3046, 2913, 1625, 1597, 1574, 1515, 1467, 1339, 888, 856, 813, 750 cm^{-1} ; EIMS (70 eV) [m/e (relative intensity)] 476 (M^+ , 31.46), 327 ($M - 149$, 4.50), 239 ($M - \text{HgCl}$, 12.58). Anal. Calcd for $\text{C}_{14}\text{H}_{11}\text{ClHgN}_2\text{O}_2$: C, 35.37; H, 2.32; N, 5.89. Found: C, 35.17; H, 2.31; N, 5.64.

The selected physical and analytical data for the following compounds (**2b–37b**) are summarized in Table IV.

2-(Chloromercurio)-4-methoxy-*N*-(4-nitrobenzylidene)-aniline (2b). Crystallization of the filtered solid from chloroform gave **2b** as orange needles.

(14) Kiefer, E. F.; Waters, W. L.; Calson, D. A. *J. Am. Chem. Soc.* 1968, 90, 5127–5131.

(15) Burgi, H. B.; Dunitz, J. D. *Helv. Chim. Acta* 1970, 53, 1747–1762.

(16) In order to further investigate the reaction mechanism we intend to carry out the kinetic measurements including the substituent effects, quantitative solvent effects, and deuterium isotope effects. These will be reported elsewhere.

(17) Vogel, A. I. *Practical Organic Chemistry*; Longmans: London, 1970; p 635.

Table IV. Selected Physical and Analytical Data for Compounds 2b–37b

compd	yield (%)	mp (°C)	anal. calcd (found)			MS [<i>m/e</i> (<i>M</i> ⁺)]	
			C	H	N	calcd	obsd
2b	63	264–266	34.22 (34.22)	2.24 (2.30)	5.70 (5.21)	492	492
3b	87	346–348	22.41 (22.50)	1.15 (1.41)	4.02 (3.79)	698	698
4b	85	338.5–339.5	31.48 (31.74)	1.62 (1.93)	5.65 (5.26)	496	496
5b ^a	93	343–344	30.82 (30.95)	2.05 (1.91)	4.79 (4.81)	540	540
6b	78	>350	26.57 (26.54)	1.36 (1.43)	4.77 (4.15)	588	588
7b ^a	90	272–273	41.06 (41.02)	2.70 (2.75)	5.04 (4.93)	512	512
8b	70	165–166	39.07 (38.96)	2.79 (2.80)	3.26 (3.08)	431	431
9b	54	187–188	37.67 (37.76)	2.69 (2.72)	3.14 (3.18)	447	447
10b	50	276.5–277.5	23.96 (23.60)	1.38 (1.79)	2.15 (1.91)	651	651
11b	44	210–211	34.63 (34.45)	2.00 (2.24)	3.11 (2.78)	451	451
12b	43	210–212	31.52 (31.19)	1.82 (1.60)	2.83 (2.53)	495	495
13b	74	219–221	28.78 (28.77)	1.66 (1.65)	2.58 (2.47)	543	543
14b		158–159	37.67 (37.76)	2.69 (2.98)	3.14 (2.85)	447	447
15b		174–176				507	507
16b	95	245–246.5	40.59 (40.50)	3.59 (3.58)	5.92 (5.81)	474	474
17b	87	184–186	39.13 (38.86)	3.04 (3.02)	3.04 (2.98)	461	461
18b	81	216–217	39.18 (38.93)	3.27 (3.27)	2.86 (2.73)	491	491
19b	91	226.5–228	39.13 (38.98)	3.04 (3.04)	3.04 (2.87)	461	461
20b	85	200–202	36.17 (35.88)	2.37 (2.39)	3.01 (3.21)	465	465
21b	93	>350	36.17 (36.16)	2.37 (2.29)	3.01 (2.98)	465	465
22b	94	220–222	36.17 (35.74)	2.37 (2.35)	3.01 (3.20)	465	465
23b	95	224–226	33.01 (32.72)	2.16 (2.15)	2.75 (2.78)	509	509
24b	80	208–210	33.01 (32.78)	2.16 (2.15)	2.75 (2.78)	509	509
25b	90	278–280	35.37 (35.17)	2.32 (2.29)	5.89 (5.86)	476	476
26b	86	240–241	35.37 (35.21)	2.32 (2.27)	5.89 (5.81)	476	476
27b	83	209–211	44.79 (44.61)	3.34 (3.30)	5.50 (5.35)	510	510
28b	75	170–172	44.76 (44.78)	4.51 (4.53)	2.18 (1.91)	645	645
29b	61	172–174	43.67 (43.63)	4.40 (4.43)	2.12 (2.18)	661	661
30b	66	189–191	41.57 (40.92)	3.92 (3.91)	2.11 (2.00)	665	665
31b	54	186–187	38.95 (38.81)	3.67 (3.73)	1.98 (1.94)	709	709
32b	90	202–204	39.07 (38.73)	2.79 (2.73)	3.26 (3.28)	431	431
33b	88	290–291	35.37 (35.34)	2.32 (2.31)	5.89 (5.88)	476	476
34b	85	219–220	27.94 (28.03)	2.07 (2.08)	3.62 (3.73)		b
35b ^a	86	297–299	24.32 (24.37)	1.62 (1.60)	3.78 (3.47)	698	698
36b	90	311–312	23.66 (23.94)	1.41 (1.55)	3.94 (4.10)		b
37b	52	254–256	33.84 (33.47)	1.95 (1.95)	6.07 (5.63)	462	462

^a Contains 0.5 molecule of crystalline 1,4-dioxane. ^b No *M*⁺ was found due to the weakness of the C–Hg bond, but *M* – HgOAc and *M* – HgCl were observed for 34b and 36b, respectively.

2,4-Bis(chloromercurio)-*N*-(4-nitrobenzylidene)aniline (3b). The crude product was crystallized from DMSO to give 3b as yellow needles.

2-(Chloromercurio)-4-chloro-*N*-(4-nitrobenzylidene)aniline (4b). The filtered solid was crystallized from 1,4-dioxane to give 4b as yellow needles.

2-(Chloromercurio)-4-bromo-*N*-(4-nitrobenzylidene)aniline (5b). The crude product was crystallized from 1,4-dioxane to yield yellow needles of 5b·1/2C₄H₈O₂.

2-(Chloromercurio)-4-iodo-*N*-(4-nitrobenzylidene)aniline (6b). Crystallization from chloroform gave yellow crystals of 6b.

α-(Chloromercurio)-*N*-(4-nitrobenzylidene)-β-naphthylamine (7b). The filtered solid was crystallized from 1,4-dioxane to give 7b·1/2C₄H₈O₂ as yellow needles.

2-(Chloromercurio)-*N*-(benzylidene)-4-toluidine (8b). Crystallization from chloroform gave 8b as white needles.

2-(Chloromercurio)-4-methoxy-*N*-(benzylidene)aniline (9b). The filtered solid was dissolved in hot chloroform and then the methanol was added dropwise until the white precipitate appeared. The mixture was heated again to make the precipitate dissolve. After cooling, white crystals of 9b were obtained.

2,4-Bis(chloromercurio)-*N*-(benzylidene)aniline (10b). The filtered solid was crystallized from the mixture solvent of DMSO and methanol to give a white solid of 10b.

2-(Chloromercurio)-4-chloro-*N*-(benzylidene)aniline (11b). Crystallization from chloroform gave white crystals of 11b.

2-(Chloromercurio)-4-bromo-*N*-(benzylidene)aniline (12b). Crystallization from chloroform yielded white crystals of 12b.

2-(Chloromercurio)-4-iodo-*N*-(benzylidene)aniline (13b). Crystallization from chloroform gave white needles of 13b.

14b and 15b were synthesized according to the literature method.⁵

2-(Chloromercurio)-*N*-(4-(dimethylamino)benzylidene)-4-toluidine (16b). Crystallization from dichloromethane gave

yellow crystals of 16b.

2-(Chloromercurio)-*N*-(4-methoxybenzylidene)-4-toluidine (17b). Crystallization from chloroform gave white crystals of 17b.

2-(Chloromercurio)-*N*-(3,4-dimethoxybenzylidene)-4-toluidine (18b). Crystallization from 1,4-dioxane yielded white crystals of 18b.

2-(Chloromercurio)-*N*-(2-methoxybenzylidene)-4-toluidine (19b). Crystallization from methanol gave white crystals of 19b.

2-(Chloromercurio)-*N*-(4-chlorobenzylidene)-4-toluidine (20b). Crystallization from chloroform gave white crystals of 20b.

2-(Chloromercurio)-*N*-(3-chlorobenzylidene)-4-toluidine (21b). Crystallization from 1,4-dioxane gave white crystals of 21b.

2-(Chloromercurio)-*N*-(2-chlorobenzylidene)-4-toluidine (22b). Crystallization from methanol gave white crystals of 22b.

2-(Chloromercurio)-*N*-(4-bromobenzylidene)-4-toluidine (23b). Crystallization from chloroform gave white crystals of 23b.

2-(Chloromercurio)-*N*-(3-bromobenzylidene)-4-toluidine (24b). Crystallization from methanol gave white crystals of 24b.

2-(Chloromercurio)-*N*-(3-nitrobenzylidene)-4-toluidine (25b). Crystallization from 1:1 dioxane/methanol gave pale yellow crystals of 25b.

2-(Chloromercurio)-*N*-(2-nitrobenzylidene)-4-toluidine (26b). Crystallization from chloroform gave yellow crystals of 26b.

α-(Chloromercurio)-(*N*-dimethylaminobenzylidene)-β-naphthylamine (27b). Crystallization from chloroform gave yellow crystals of 27b.

Mercuration Reaction of 28a–31a. 4'-((Arylimino)methylbenzo-15-crown-5 (28a–31a) were prepared by treating 4'-((formylbenzo)-15-crown-5 with corresponding substituted anilines (1:1 molar ratio) in refluxing ethanol. Mp: for 28a 104.5–105.5 °C; 29a 102–103 °C; 30a 104–105 °C; 31a 123–125 °C. 4'-((Formylbenzo)-15-crown-5 was prepared according to the literature method.¹⁸ Mp: 80–81 °C.

Mercuriation of 28a. To a solution of 28a (1.00 g, 2.6 mmol) in methanol was added 0.80 g of mercuric acetate (2.5 mmol); the mixture was refluxed for 4 h and allowed to cool to room temperature. The resulting white crystals were filtered off, washed with methanol, and dried in vacuo to give 28b.

Compounds 29b-31b were prepared by using the above method.

4-(Chloromercurio)-*N*-benzylidene-2-toluidine (32b). Crystallization of the filtered solid from methanol yielded white crystals of 32b.

4-(Chloromercurio)-*N*-(nitrobenzylidene)-2-toluidine (33b). Crystallization of the filtered solid from 1,4-dioxane gave 33b as yellow needles.

2,4-Bis(acetomercurio)-6-methoxy-*N*-(4-nitrobenzylidene)aniline (34b). Crystallization from methanol yielded orange crystals of 34b.

2,4-Bis(chloromercurio)-*N*-(2-nitrobenzylidene)aniline (35b). Crystallization of the filtered solid from 1,4-dioxane gave yellow crystals of 35b.^{1/2}C₄H₈O₂.

2,4-Bis(chloromercurio)-6-methyl-*N*-(2-nitrobenzylidene)aniline (36b). The filtered solid was recrystallized from 1,4-dioxane to give yellow needles of 36b.

4-(Chloromercurio)-*N*-(4-nitrobenzylidene)aniline (37b). To a solution of 1a (1.13 g, 5 mmol) and BTMAC (benzyltrimethylammonium chloride) (0.95 g, 5 mmol) in dry methanol (50 mL) was added 1.60 g (5 mmol) of mercuric acetate. The mixture was stirred and refluxed for 6 h, during which a yellow solid deposited. This solid was filtered off and dried in vacuo and then extracted with hot chloroform until the extracts were nearly colorless. Concentration of the extracts and subsequent crystallization gave yellow needles of 37b as the main product. The remaining insoluble solid was recrystallized from DMSO to give yellow needles of 3b.

X-ray Structure Determination of 7b. Single crystals of 7b suitable for X-ray diffraction study were obtained by slow evaporation of the solution of 7b in 1,4-dioxane at room tem-

perature. A single crystal of approximate dimensions 0.2 × 0.2 × 0.25 mm was mounted on the end of a glass fiber. The X-ray diffraction intensity data of 3196 independent reflections, of which 2666 with $I > 3\sigma(I)$ were observable, were collected with an Enraf-Nonius CAD-4 diffractometer by using graphite monochromated Cu K α radiation with the ω -2 θ scan mode within the range of $2 \leq 2\theta \leq 120^\circ$. The structure was solved by direct methods using the MULTAN82 program. The non-hydrogen atoms were found by difference Fourier (ΔF) synthesis and blocked diagonal matrix least-squares procedures. Hydrogen atoms were fixed by the theoretical method. After three cycles of refinement by full-matrix least-squares procedure with anisotropic thermal parameters for all non-hydrogen atoms, the final values of $R = [\sum \|F_o\| - |F_c|]/\sum \|F_o\| = 0.073$ and $R_w = \{[\sum w(F_o - |F_c|)^2]/\sum wF_o^2\}^{1/2} = 0.075$ were obtained.¹⁹ An ORTEP drawing of 7b is shown in Figure 1.

Acknowledgment. We are grateful to the National Natural Science Foundation of PRC and the Natural Science Foundation of Henan Province for the support of this research. We thank Professor Wang Hongfei and Professor Cheng Jinpei for valuable discussion of this paper. We also thank Dr. Kraus for comments on this paper.

Supplementary Material Available: A listing of spectral data for compounds 2b-37b, figures showing the thermal analysis curve and molecular packing in the unit cell for 7b, and tables of positional parameters and their estimated deviations for all non-hydrogen atoms, bond distances, bond angles, deviations of atoms from the mean plane, and crystal data and experimental details for the structure determination of 7b (14 pages). Ordering information is given on any current masthead page.

OM920366N

(18) Hyde, E. M.; Shaw, B. L.; Shepherd, I. J. *Chem. Soc., Dalton Trans.* 1978, 1696-1705.

(19) All crystallographic calculations were carried out on a PDP 11/44 computer with the CAD4-SDP program package (Enraf-Nonius, Delft, The Netherlands).

Synthesis and Photochemistry of the Aqueous-Soluble $(\eta^5\text{-C}_5\text{H}_4\text{CH}_2\text{CH}_2\text{NH}_3^+)_2\text{Mo}_2(\text{CO})_6$ Complex. Generation of 19-Electron Complexes in Aqueous Solution

Alfred Avey and David R. Tyler*

Department of Chemistry, University of Oregon, Eugene, Oregon 97403

Received May 28, 1992

The synthesis and photochemistry of the $(\eta^5\text{-C}_5\text{H}_4\text{CH}_2\text{CH}_2\text{NH}_3^+)_2\text{Mo}_2(\text{CO})_6$ complex are described. The NO_3^- salt is soluble in aqueous solutions at pH 8 or lower; the PF_6^- salt was used in organic solvents. The photochemistry of the molecule is similar to the photochemistry of the $\text{Cp}_2\text{Mo}_2(\text{CO})_6$ ($\text{Cp} = \eta^5\text{-C}_5\text{H}_5$) complex: irradiation into the low-energy tail of the lowest energy absorption band ($\lambda_{\text{max}} \approx 503 \text{ nm}$; $d\pi \rightarrow \sigma^*$) results in homolytic cleavage of the Mo-Mo bond to form $(\eta^5\text{-C}_5\text{H}_4\text{CH}_2\text{CH}_2\text{NH}_3^+)\text{Mo}(\text{CO})_3$ radicals. These 17-electron radicals can be trapped with chlorine atom donors (e.g., CCl_4 in nonaqueous solution or $\text{CCl}_3\text{CH}_2\text{OH}$ in aqueous solution) to form $(\eta^5\text{-C}_5\text{H}_4\text{CH}_2\text{CH}_2\text{NH}_3^+)\text{Mo}(\text{CO})_3\text{Cl}$. Irradiation of $(\eta^5\text{-C}_5\text{H}_4\text{CH}_2\text{CH}_2\text{NH}_3^+)_2\text{Mo}_2(\text{CO})_6$ in the presence of ligands [e.g., phosphines in nonaqueous solution or water-soluble phosphines such as PTA (phosphatriazaadamantane) in aqueous solution] resulted in disproportionation of the dimer complex. The disproportionation reactivity is consistent with the intermediate formation of 19-electron adducts. Like their counterparts in nonaqueous solvents, the putative 19-electron adducts generated in aqueous solution are good reductants. They were used to reduce methylviologen and cytochrome *c* in aqueous solution. Reduction of protons to H_2 did not occur, and this led us to reexamine our earlier results in which we suggested that 19-electron complexes formed by irradiation of $(\text{CpCOOH})_2\text{W}_2(\text{CO})_6$ in basic aqueous solution reduced water to H_2 . CO_2 is formed in a 1:1 ratio with H_2 in this reaction, and we propose the water-gas shift reaction is the source of the H_2 .

We recently began a research program to explore the reactivity of 19-electron organometallic complexes in

aqueous solution.¹⁻⁴ Nineteen-electron complexes have several features that make them ideal reducing agents,⁸