

# Metallacyclophanes formed by a tetrapyrazolyl ligand and copper(II) cation

Abdelaziz Jouaiti, Marielle Lœi, Mir Wais Hosseini\* and André De Cian

Laboratoire de Chimie de Coordination Organique, Université Louis Pasteur, F-67000 Strasbourg, France, UMR CNRS 7513. E-mail: hasseini@chimie.u-strasbg.fr

Received (in Cambridge, UK) 15th August 2000, Accepted 18th September 2000

First published as an Advance Article on the web 19th October 2000

Using a tetrakis(pyrazolyl)benzene ligand and  $\text{CuCl}_2$  or  $\text{Cu}(\text{CF}_3\text{SO}_3)_2$  salts, binuclear metallamacrocycles of cyclophane type have been exclusively obtained; structural characterisation by X-ray on single-crystal reveals that in both cases, chloride or triflate anions are coordinated to Cu metal centres adopting a square pyramidal coordination geometry.


Metallamacrocycles, macrocyclic frameworks based on interconnection of organic ligands by metal cations, have been attracting much attention over the past decade. Although the first examples were reported by Saalfrank *et al.*<sup>1</sup> and Fujita,<sup>2</sup> many other groups are now active in the design and synthesis of metallamacrocycles.<sup>3</sup> However, as connecting metallic centres the majority of examples reported so far are based on the use of diamagnetic transition metals such as Pd(II) and Pt(II) requiring square coordination geometry. Thus, using these metals, a variety of metallamacrocycles has been reported using ligands such as 4,4'-bipyridine, pyrimidine, bipyrazine, 4,7-phenanthroline or bis-pyridines interconnected by rigid or flexible spacers.<sup>4</sup> Using pyrazolyl based ligands, we have previously reported the formation of metallamacrocycles and metalatubulanes.<sup>5</sup>

Here, we report the design of new ligands based on pyrazolyl units as coordination sites and its self-assembly into binuclear metallamacrocycles using pentacoordinated paramagnetic Cu(II) cation.

Ligands **1** and **2** (Scheme 1), both based on aromatic cores (benzene or biphenyl) and bearing four pyrazolyl units, are designed to act as tetradentate ligands. For ligand **1**, two different arrangements of the coordination sites may be envisaged leading to either a seven (1A) or an eight membered ring cyclic system (1B) upon chelating to metal cations (Scheme 1). In both cases, owing to the disposition of the coordination sites ligand **1** may form metallamacrocycles in the presence of metals requiring square, octahedral or square pyramidal coordination geometry. For the latter case, the two possibilities (4A and 4B) are shown (Scheme 1). Ligand **2** was designed as an extended analogue of ligand **1** thus allowing an increase in the size of the metallamacrocycle.

Ligand **1** was first prepared in 8% yield by reacting 1,2,4,5-tetrachlorobenzene with sodium pyrazolate in DMF at 95 °C for 3 d. The yield could be increased to 68% by using 1,2,4,5-tetrabromobenzene and under reflux for 72 h. Pure compound **1** was obtained as a colourless solid after chromatography ( $\text{SiO}_2$ ,  $\text{CH}_2\text{Cl}_2$ -MeOH) and was characterised by X-ray crystallography† (Fig. 1) which revealed the following relevant bond distances and angles ( $d_{\text{NN}} = 1.37 \text{ \AA}$ ,  $d_{\text{C(Ph)N}} = 1.42 \text{ \AA}$ ; NNCC dihedral angles of  $-25.8$  and  $-77.2^\circ$ ). Ligand **2** was prepared by a nickel coupling reaction using  $\text{NiBr}_2(\text{PPh}_3)_2$ <sup>6</sup> in the presence of Zn and  $\text{NEt}_4\text{I}$  in THF using compound **3** as the starting material. The latter was obtained upon treatment of 1,3,5-benzene by sodium pyrazolate in DMF at 95 °C for 4 d in 22% yield after chromatography [ $\text{Al}_2\text{O}_3$ ,  $\text{CH}_2\text{Cl}_2$ -hexane (8/2)].

Cu(II) was selected as the metal ion because of its paramagnetic nature and the fact that it forms pentacoordinated complexes. Both the chloride and triflate Cu(II) complexes have been obtained as single-crystals upon slow diffusion of a MeOH



Scheme 1

solution of  $\text{CuCl}_2$  or  $\text{Cu}(\text{OTf})_2$  (5 mg) into a  $\text{CHCl}_3$  solution of ligand **1** (2 mg). For both complexes, their structure was investigated in the crystalline state by X-ray crystallography.

The reaction of **1** and  $\text{CuCl}_2$  in  $\text{CHCl}_3$ -MeOH afforded a green crystalline material which was structurally characterised by X-ray diffraction (Fig. 2).† The solid contains the complex unit  $[\text{1}_2\text{Cu}_2(\text{Cl}_2(\text{CuCl}_4)_2)]$  and two MeOH and  $\text{CHCl}_3$  solvent molecules with no specific interactions with the complex. The cationic part of the complex is indeed a metallamacrocycle composed of two ligands **1**, two Cu(II) cations and two  $\text{Cl}^-$





Fig. 1 The X-ray structure of free ligand **1**. H atoms are omitted for clarity, for bond length and distances see text.




**Fig. 2** The X-ray structure of the metalloamacycle  $[1_2Cu_2Cl_2]^{2+}$  obtained between **1** and  $CuCl_2$ . H atoms, anions and solvent molecules are omitted for clarity.


anions. Owing to the parallel orientation of the two phenyl rings, the binuclear metallamacrocycle is of the cyclophane type with an internal cavity of  $4.29 \times 7.42$  Å (Scheme 1). The coordination sphere around the pentacoordinated  $Cu(II)$  cations is composed of four nitrogen atoms belonging to two ligands **1** and one chloride anion with a slightly distorted square pyramidal geometry. The two poles of the ligand **1**, composed each of two pyrazolyl moieties ( $d_{NN} = 1.37$  Å,  $d_{C(Ph)N} = 1.42$  Å; NNCC dihedral angle of  $57.8^\circ$ ) and occupying the square base in the coordination sphere around the metal, act as bidentate units and form two seven membered metallarings of type **4A** (Scheme 1) with Cu–N distances of ca. 2.01 Å and NCuN angles of  $86.9$ – $91.4^\circ$  (Scheme 1). The axial position is occupied by a Cl anion with Cu–Cl distance of 2.44 Å. The two positive charges on the  $[1_2Cu_2Cl_2]^{2+}$  complex are neutralised by a  $[CuCl_4]^{2-}$  anion with  $d_{CuCl} = 2.283$  Å and ClCuCl angle of  $90^\circ$  (Fig. 3).

The  $[1_2Cu_2(CF_3SO_3)_2(CF_3SO_3)_2]$  complex was obtained as a blue-violet crystalline solid upon reaction of **1** and  $Cu(CF_3SO_3)_2$  in  $CHCl_3$ –MeOH. The latter complex was also structurally characterised by X-ray diffraction (Fig. 4).<sup>†</sup> The solid contains the complex unit  $[1_2Cu_2(CF_3SO_3)_2(CF_3SO_3)_2]$  and two  $CH_2Cl_2$  solvent molecules with no specific interactions with the complex. The triflate anions are disordered. The cationic part of the complex of type **4A** (Scheme 1) shows almost identical structural features as the above chloride complex (Scheme 1). Again a metallamacrocycle (internal cavity of  $4.16$  Å  $\times$   $7.29$  Å) composed of two ligands **1**, two  $Cu(II)$  cations and two triflate anions is observed. The coordination sphere around the pentacoordinated  $Cu(II)$  cations is, as in the chloride case, comprised of four nitrogen atoms belonging to two ligands **1** and one triflate anion ( $d_{NN} = 1.38$  Å  $d_{C(Ph)N} = 1.42$  Å; NNCC dihedral angles of  $-59.6$  and  $58.9^\circ$ , Cu–N distances of ca. 1.99 Å and NCuN angles of  $87.4$ – $92.6^\circ$ . Interestingly, the axial position is occupied by a triflate anion with Cu–O distance of 2.20 Å. The two positive charges on the  $[1_2Cu_2(CF_3SO_3)_2]^{2+}$  complex are neutralised by two external triflate anions (Fig. 4).

We have previously reported examples of metallatubulanes based on interconnection of metallamacrocycles into infinite 1-D



**Fig. 3** The packing of the dicationic metallamacrocycle  $[1_2Cu_2Cl_2]^{2+}$  and  $CuCl_4^{2-}$  anions. H atoms and solvent molecules are not presented for clarity.



**Fig. 4** The X-ray structure of the metallamacrocycle  $[1_2Cu_2(CF_3SO_3)_2]^{2+}$  obtained between **1** and  $Cu(CF_3SO_3)_2$ . Then triflate anions are disordered. H atoms, anions and solvent molecules are omitted for clarity.

coordination networks with tubular topology.<sup>5,7,8</sup> In relation to this, it may be of interest that, since the two anions occupying the summit of the square pyramid on each Cu centre are oriented in a divergent fashion, by interconnecting the metalloacyclophanes new types of coordination polymers may be obtained using bridging bidentate anionic ligands such as azido or isocyanato anions.

In conclusion, the synthesis of paramagnetic binuclear copper metallacyclophanes has been achieved and their solid state structural elucidation revealed that the  $Cu(II)$  cation adopts a slightly distorted square pyramidal geometry with the axial position occupied either by chloride or triflate anion. The formation of extended metallacyclophanes using ligand **2** is being currently pursued. Furthermore, the interconnection of the obtained metallamacrocycles leading to paramagnetic coordination networks through the substitution of monodentate coordinated anions by bidentate anions is currently under investigation.

## Notes and references

<sup>†</sup> Crystal data: (colourless, 294 K),  $C_{18}H_{14}N_8$ ,  $M = 342.37$ , triclinic,  $a = 6.384(1)$ ,  $b = 7.007(1)$ ,  $c = 9.941(1)$  Å,  $\alpha = \beta = 76.00$ ,  $\gamma = 80.00$ ,  $U = 416.5(2)$  Å<sup>3</sup>,  $Z = 1$ , space group  $P1$ ,  $D_c = 1.36$  g cm<sup>-3</sup>,  $\mu = 0.089$  mm<sup>-1</sup>, 1508 data with  $I > 3\sigma(I)$ ,  $R = 0.040$ ,  $R_w = 0.0649$ .  $[(1_2Cu_2Cl_2)(CuCl_4)]$ , (green, 294 K),  $2(CuClC_{18}H_{14}N_8) \cdot CuCl_4 \cdot 2CHCl_3 \cdot 2MeOH$ ,  $M = 1390.91$ , orthorhombic,  $a = 10.0302(5)$ ,  $b = 11.3568(6)$ ,  $c = 23.5388(7)$  Å,  $U = 2681.3(4)$  Å<sup>3</sup>,  $Z = 2$ ,  $Immm$ ,  $D_c = 1.72$  g cm<sup>-3</sup>,  $\mu = 1.830$  mm<sup>-1</sup>, 1486 data with  $I > 3\sigma(I)$ ,  $R = 0.042$ ,  $R_w = 0.052$ .  $[1_2Cu_2(CF_3SO_3)_2(CF_3SO_3)_2]$ , (blue-violet, 173 K),  $C_{38}H_{28}Cu_2F_6N_{16}O_6S_2 \cdot 2CF_3SO_3 \cdot 2CH_2Cl_2$ ,  $M = 1577.95$ , orthorhombic,  $a = 19.7081(7)$ ,  $b = 17.9293(6)$ ,  $c = 17.1966(6)$  Å,  $U = 6076.5(6)$  Å<sup>3</sup>,  $Z = 4$ , space group  $Cmca$ ,  $D_c = 1.72$  g cm<sup>-3</sup>,  $\mu = 1.121$  mm<sup>-1</sup>, 2127 data with  $I > 3\sigma(I)$ ,  $R = 0.077$ ,  $R_w = 0.096$ . Data for all 3 structures have been obtained on a Nonius Kappa CCD (Mo–K $\alpha$ ) diffractometer. Structural determinations were performed using the Nonius OpenMolenN package.<sup>9</sup> CCDC 182/1781. See <http://www.rsc.org/supp-data/cc/b0/b006665f/> for crystallographic data in .cif format.

- 1 R. W. Saalfrank, A. Stark, K. Peters and H.-G. von Schnering, *Angew. Chem., Int. Ed.*, 1988, **27**, 851.
- 2 M. Fujita, in *Comprehensive Supramolecular Chemistry*, ed. J. L. Atwood, J. E. D. Davies, D. D. MacNicol and F. Vögtle, Vol. 9 (ed. J. P. Sauvage and M. W. Hosseini, Elsevier, 1996, p. 253).
- 3 J. K. M. Sanders, in *Comprehensive Supramolecular Chemistry*, ed. J. L. Atwood, J. E. D. Davies, D. D. MacNicol and F. Vögtle, Vol. 9 (ed. J. P. Sauvage and M. W. Hosseini), Elsevier, 1996, p. 131; P. J. Stang and B. Olenyuk, *Acc. Chem. Res.*, 1997, **30**, 502; C. M. Drain and J.-M. Lehn, *J. Chem. Soc., Chem. Commun.*, 1994, 2313; T. Beissel, R. E. Powers and K. N. Raymond, *Angew. Chem., Int. Ed. Engl.*, 1996, **35**, 1084.
- 4 C. V. Krishnamohan Sarma, S. T. Griffin and R. D. Rogers, *Chem. Commun.*, 1998, 215; R.-D. Schnebeck, L. Radaccio, E. Zangrando and B. Lippert, *Angew. Chem., Int. Ed.*, 1988, **37**, 119; J. R. Hall, S. J. Loeb, G. K. H. Shimizu and G. P. A. Yap, *Angew. Chem., Int. Ed.*, 1998, **37**, 121; R. Schneider, M. W. Hosseini, J.-M. Planeix, A. De Cian and J. Fischer, *J. Chem. Soc., Chem. Commun.*, 1988, 1625.
- 5 M. Loï, M. W. Hosseini, A. Jouaiti, A. De Cian and J. Fischer, *Eur. J. Inorg. Chem.*, 1999, 1981.
- 6 M. Iyoda, H. Otsuka, K. Sato, N. Nisato and M. Oda, *Bull. Chem. Soc. Jpn.*, 1990, **63**, 80; V. K. Yamamoto, *Bull. Chem. Soc. Jpn.*, 1954, **27**, 501.
- 7 G. Mislin, E. Graf, M. W. Hosseini, A. De Cian, N. Kyritsakas and J. Fischer, *Chem. Commun.*, 1998, 2545.
- 8 C. Klein, E. Graf, M. W. Hosseini, A. De Cian and J. Fischer, *Chem. Commun.*, 2000, 239.
- 9 OpenMolenN, Interactive Structure Solution, Nonius B. V., Delft, The Netherlands, 1997.