

5-Amino-6-phenyl-1,6-dihydro-
pyridazin-3(2H)-oneHéctor Novoa de Armas,^{a,*†} Norbert M. Blaton,^a
Oswald M. Peeters,^a Camiel J. De Ranter,^a Beatriz Pita,^b
Eddy Sotelo,^b Enrique Ravina^b and Margarita Suárez^c^aLaboratorium voor Analytische Chemie en Medicinale Fysicochemie, Faculteit Farmaceutische Wetenschappen, Katholieke Universiteit Leuven, Van Evenstraat 4, B-3000 Leuven, Belgium, ^bDepartamento de Química Orgánica, Laboratorio de Química Medicinal, Facultad de Farmacia, Universidad de Santiago de Compostela, Santiago de Compostela, 15706 Galicia, Spain, and ^cDepartamento de Química Orgánica, Laboratorio de Síntesis Orgánica, Universidad de La Habana, Apartado 10400, La Habana, Cuba

Correspondence e-mail: hector.novoa@farm.kuleuven.ac.be


Received 13 September 1999

Accepted 24 November 1999

In the title compound, C₁₀H₉N₃O, the pyridazinone moiety is essentially planar and forms a dihedral angle of 49.5 (1)° with the phenyl substituent. The molecular packing is stabilized by van der Waals interactions and hydrogen bonds.

Comment

It is known that 6-aryl-3(2H)-pyridazinones and their 4,5-dihydro derivatives display several pharmacological activities, all of them related to cardiotonics, such as reduction of blood pressure, inhibition of platelet aggregation, positive inotropic activity, and others (Robertson *et al.*, 1986). Likewise, 6-aryl-pyridazinones with nitro and acyl substituents at the 4- and 5-positions show good antiaggregating properties (Schudt *et al.*, 1991). We have previously reported the synthesis of 5-aminomethyl-6-aryl-4,5-dihydropyridazinones and 6-aryl-5-aminomethyl-3(2H)-pyridazinones (Ravina *et al.*, 1990). Some of these compounds show a good *in vitro* inhibitory activity on ADP-induced rat platelet aggregation. As a continuation of this previous report on the chemistry and pharmacology of this class of compounds, we carried out the crystal structure



determination of 5-amino-6-phenyl-1,6-dihydropyridazin-3(2H)-one, (I). This enamine-like compound can be employed in the synthesis of hetero-condensed pyridazinones. Recently,

† On leave from: Centro de Química Farmacéutica, Departamento de Análisis, Apartado 16042, La Habana, Cuba.




Figure 1

A plot of (I) showing the atomic numbering scheme. Displacement ellipsoids are drawn at 50% probability level, and H atoms are shown as spheres of arbitrary radii.

we used this compound as an intermediate in the synthesis of pyrido[2,3-*d*]pyridazines (Pita *et al.*, 2000).

There are no unusual bond distances and angles in (I), and they are in the range of calculated values using the *AM1* method in related structures (Estevez *et al.*, 1998). The bond lengths in the pyridazinone ring range from 1.304 (2) to 1.450 (3) Å. The torsion angle between the pyridazinone and the phenyl ring, found using the quantum chemical *AM1* method in *MOPAC* (Stewart, 1990) for the lower energy conformations, is in the range 40–140°, with a heat of formation of 38.30 kcal mol⁻¹ (1 kcal = 4.184 kJ). In the crystal structure this angle is -51.3 (3)°, which corresponds to the minimum in the energy calculations. The calculated favoured conformation of the enol form corresponds to torsion angles in the same range (40–140°) and a heat of formation of 34.20 kcal mol⁻¹, which shows that this enol is the predominant form at equilibrium. This is contrary to the fact that in the crystal the molecule is present in the amide form, which corresponds to a higher heat of formation. The dihedral angle between the respective least-squares planes of the pyridazinone ring and the phenyl ring is 49.5 (1)°. The mean *Csp*²—*Csp*² bond length within the phenyl ring is 1.382 (1) Å. A view of (I) can be seen in Fig. 1.

The N3 atom of the amine and the N1 atom of the amide group in the pyridazinone ring are involved in two intermolecular hydrogen bonds with a neighbouring O1, forming an infinite two-dimensional network in the plane [001].

Experimental

A suspension of 5-bromo-6-phenyl-3(2H)-pyridazinone (0.5 g, 1.9 mmol), ammonium chloride (0.3 g, 5.6 mmol) and ammonium hydroxide (50 ml) was heated at 458 K at pressure of 374 psi (1 psi ≈ 6.895 kPa) for 3 h in a Parr reactor. The mixture was evaporated *in vacuo* and washed with ammonium hydroxide, and the solid obtained, (I), was recrystallized from ethanol (yield 70%; m.p. 517 K). Spectroscopic analysis: IR (KBr), cm⁻¹: 3480–3425 (NH), 1670 (CO); ¹H NMR (DMSO-*d*₆, p.p.m.): δ 12.12 (*s*, 1H, NH, deuterium oxide exchangeable), 7.50–7.43 (*m*, 5H, aromatics), 5.71 (*s*, 1H, CH—CO), 5.96 (*s*, 2H, NH₂); ¹³C NMR (p.p.m.): δ 162.5 (C3), 99.1 (C4), 149.2 (C5), 140.1 (C6), 134.4 (C1'), 129.1 (C2', C6'), 128.8 (C3', C5'), 129.1 (C4'); analysis, calculated: C 64.16, H 4.85, N 22.45%; found: C 64.20, H 4.78, N 22.43%.

Table 1

Selected geometric parameters (Å, °).

O1—C1	1.265 (2)	N3—C3	1.346 (3)
N1—N2	1.344 (2)	C1—C2	1.395 (3)
N1—C1	1.367 (2)	C2—C3	1.376 (3)
N2—C4	1.304 (2)	C3—C4	1.450 (3)
N2—N1—C1	125.86 (15)	N3—C3—C2	122.12 (17)
N1—N2—C4	118.06 (15)	N3—C3—C4	120.82 (16)
O1—C1—C2	126.04 (17)	N2—C4—C5	114.71 (16)
N1—C1—C2	115.88 (16)	N2—C4—C3	121.97 (16)
O1—C1—N1	118.07 (16)		

Crystal data

C₁₀H₉N₃O
M_r = 187.20
 Orthorhombic, *Pbca*
a = 8.752 (2) Å
b = 10.525 (5) Å
c = 20.619 (5) Å
V = 1899.3 (11) Å³
Z = 8
D_x = 1.309 Mg m⁻³

Cu *Kα* radiation
 Cell parameters from 25 reflections
 θ = 10.78–28.12°
 μ = 0.73 mm⁻¹
T = 293 (2) K
 Prism, light green
 0.48 × 0.20 × 0.14 mm

Data collection

Siemens *P4* four-circle diffractometer
 2 θ / ω scans
 Absorption correction: ψ -scan (North *et al.*, 1968)
T_{min} = 0.626, *T_{max}* = 0.903
 1733 measured reflections
 1266 independent reflections
 1102 reflections with *I* > 2 σ (*I*)

R_{int} = 0.024
 θ_{max} = 57.19°
h = -1 → 9
k = -1 → 11
l = -1 → 22
 3 standard reflections every 100 reflections
 intensity decay: <1.0%

Refinement

Refinement on *F*²
R(*F*) = 0.038
wR(*F*²) = 0.104
S = 1.100
 1266 reflections
 128 parameters
 H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0532P)^2 + 0.757P]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.002$
 $\Delta\rho_{\text{max}} = 0.14 \text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.14 \text{ e \AA}^{-3}$
 Extinction correction: *SHELXL97* (Sheldrick, 1997)
 Extinction coefficient: 0.0110 (8)

Data collection: *XSCANS* (Siemens, 1996); cell refinement: *XSCANS*; data reduction: *XSCANS*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97*; molecular graphics: *DIAMOND* (Bergerhoff, 1996); software used to prepare material for publication: *PLATON*

Table 2

Hydrogen-bonding geometry (Å, °).

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
N1—H1...O1 ⁱ	0.86	1.96	2.818 (2)	180
N3—H3A...O1 ⁱⁱ	0.86	2.06	2.880 (3)	159

Symmetry codes: (i) $-x, -y, -z$; (ii) $\frac{1}{2} + x, \frac{1}{2} - y, -z$.

(Spek, 1990), *PARST* (Nardelli, 1983, 1995) and *PARSTCIF* (Nardelli, 1992).

This work was supported in part by the Administration Générale de la Coopération au Développement, AGCD (Matricule: 911264) from the Belgian Government (ABOS-AGCD) and the Katholieke Universiteit, Leuven. BP and MS are indebted to Proyectos Alma Mater-U. H. and the Spanish CYTED for financial support.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: SK1336). Services for accessing these data are described at the back of the journal.

References

- Bergerhoff, G. (1996). *DIAMOND*. Gerhad-Domagk-Strasse 1, 53121 Bonn, Germany.
 Estevez, I., Raviña, E. & Sotelo, E. (1998). *J. Heterocycl. Chem.* **35**, 1421–1428.
 Nardelli, M. (1983). *Comput. Chem.* **7**, 95–98.
 Nardelli, M. (1992). *PARSTCIF*. University of Parma, Italy.
 Nardelli, M. (1995). *J. Appl. Cryst.* **28**, 659.
 North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
 Pita, B., Sotelo, E., Raviña, E., Suárez, M., Ochoa, E., Verdecia, Y., Novoa de Armas, H., Blaton, N., Peeters, O. M. & De Ranter, C. (2000). In preparation.
 Raviña, E., Teran, C., Santana, L., García-Domínguez, N. & Estevez, I. (1990). *Heterocycles*, **31**, 1967–1971.
 Robertson, D. W., Krushinski, J. H., Beedle, E. E., Wyss, W., Pollosk, G. D., Wilson, H., Kaufmann, R. F. & Hayes, J. S. (1986). *J. Med. Chem.* **29**, 1832–1836.
 Schudt, C., Winder, S., Muller, B. & Ukena, D. (1991). *Biochem. Pharmacol.* **42**, 153–158.
 Sheldrick, G. M. (1997). *SHELXL97* and *SHELXS97*. University of Göttingen, Germany.
 Siemens (1996). *XSCANS*. Version 2.2. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
 Spek, A. L. (1990). *Acta Cryst.* **A46**, C-34.
 Stewart, J. J. P. (1990). *MOPAC6.0*. QCPE No. 455. Frank J. Seiler Research Laboratory, US Air Force Academy, CO 80840, USA.