

## Unexpected Dipolarophilic Activity of Diphenylvinylphosphine towards Nitrones

Alberto Brandi,<sup>a\*</sup> Andrea Goti,<sup>b</sup> and K. Michał Pietrusiewicz<sup>\*c</sup>

<sup>a</sup> Istituto di Chimica, Università della Basilicata, 85100 Potenza, Italy

<sup>b</sup> Centro di Studio sulla Chimica e la Struttura dei Composti Eterociclici e loro Applicazioni, CNR, Via G. Capponi 9, 50121 Firenze, Italy

<sup>c</sup> The Polish Academy of Sciences, Centre of Molecular and Macromolecular Studies, Boczna 5, 90-362 Łódź, Poland

The reactivity of diphenylvinylphosphine towards nitrones is dominated by the dipolarophilic nature of its double bond rather than by the ability of the P<sup>III</sup> to act as an oxygen acceptor.


Phosphines were early found to act as efficient deoxygenating agents towards a variety of amine oxides including nitrones.<sup>1–3</sup> Both acyclic<sup>1</sup> and cyclic nitrones [5,5-dimethylpyrroline *N*-oxide (DMPO)<sup>4</sup> (**1a**) and 3,4-dihydroisoquinoline *N*-oxide<sup>5</sup> (**1b**)] were reported to undergo smooth deoxygenation to the parent imine upon treatment with triphenylphosphine under thermal conditions, the driving force for the process being the formation of a strong P=O bond. We now demonstrate that nitrones can undergo facile 1,3-dipolar cycloaddition to diphenylvinylphosphine (**2**) with almost complete exclusion of the oxygen transfer process (Scheme 1).

Thus, the reaction of 1.3 equiv. of DMPO (**1a**) with 1 equiv. of diphenylvinylphosphine (**2**) (benzene, room temperature, 6 days) led to the formation of the corresponding phosphinoisoxazolidines (**3a**) in 88% yield (<sup>31</sup>P n.m.r., entry 1, Table 1). Analogous reactions of two other nitrones (**1b**) and (**1c**) (entries 2 and 3) were similarly successful even though prolonged heating was applied in these cases in order to attain more practical conversion rates. Oxidation of the phosphine function in all these cases was practically negligible. The products were conveniently isolated by simple crystallisation of the crude reaction mixtures from ligroin or by a quick passage through a short pad of silica gel. All products were assigned the structures displayed on the basis of careful <sup>1</sup>H, <sup>13</sup>C, and <sup>31</sup>P n.m.r. analyses of the isolated materials as well as crude reaction mixtures. In all three cases a strong preference for the formation of the 5-substituted isoxazolidine was manifested, as expected on the basis of perturbation theory.<sup>6</sup>

In the reaction of *C,N*-diphenyl nitrone (**1d**) with (**2**) (1:1 stoichiometric ratio) in boiling benzene the deoxygenation of the nitrone turned out to be the preferred reaction pathway as

evidenced by the isolation of diphenylvinylphosphine oxide (**5**) (32% yield) as well as by detection of benzyldeneaniline (**6**) and azoxybenzene (**7**) in the crude reaction mixture (g.c.-m.s.) (Scheme 2). A mixture of isomeric 5- and 4-substituted phosphinylisoxazolidines, (**8**) and (**9**) respectively (Scheme 2), was also formed in 32% yield† with their ratio being exactly the same as found earlier in the reaction of the same nitrone with diphenylvinylphosphine oxide (**5**).<sup>7</sup> It can therefore be concluded that oxidation of phosphorus precedes the cycloaddition in the studied case since with the reversed sequence of events much more of the 5-substituted isomer would have been expected, due to the electron donating character of the phosphino group. With two equivalents of the nitrone (**1d**) the adducts (**8**) and (**9**)<sup>7</sup> can be isolated in 76% overall yield.

Whereas the absence of deoxygenation process, even in the presence of an excess of nitrone, in entries 1 and 2 can be explained by the high dipole reactivity of the cyclic nitrones (even though both were reported to be deoxygenated by triphenylphosphine)<sup>4,5</sup> the divergent behaviour of the two acyclic nitrones (**1c**) and (**1d**) is less apparent, since *C,N*-diphenylnitrone (**1d**) is known to be at least five times more reactive as a dipole than *C*-phenyl-*N*-methyl nitrone (**1c**).<sup>8</sup> The observed difference in reactivity is probably due to the higher polarity of the N–O bond in the *N*-phenyl nitrone which makes the oxygen more readily accessible for phosphorus.‡



**Table 1.** Reaction of nitrones with diphenylvinylphosphine (**2**).

Entry	Nitron	Reaction conditions	Products (regioisomeric ratio)	% Yield <sup>a</sup>	<sup>31</sup> P N.m.r. of adducts		
					5-Subst. (diast. ratio)	4-Subst.	
1	( <b>1a</b> )	Room temp., 6 days	( <b>3a</b> )	62 (88)	-8.77 (6:1)	-7.76	—
2	( <b>1b</b> )	80 °C, 24 h	( <b>3b</b> )	45 (79)	-7.02 (2:1)	-3.02	—
3	( <b>1c</b> )	80 °C, 48 h	( <b>3c</b> ) ( <b>4c</b> ) (87:13)	63 (75)	-6.36 (2.3:1)	-9.62	-7.13
4	( <b>1d</b> ) <sup>b</sup>	80 °C, 48 h	( <b>8</b> ) ( <b>9</b> ) (40:60)	76	28.09 (1.7:1)	27.76	27.27

<sup>a</sup> Yields of isolated phosphorus containing products; in parentheses are <sup>31</sup>P n.m.r. yields. <sup>b</sup> 2 equiv.

In summary, it has been demonstrated that 1,3-dipolar cycloaddition of nitrones to vinylphosphine is a viable process with the possible exclusion of *N*-aryl nitrones. Phosphinoisoxazolidines are of great potential utility as ligands.

We thank the Italian National Research Council (CNR) and the Polish Academy of Sciences (CPBP-01.13) for financial support, and Prof. F. De Sarlo and Prof. J. Michalski for their interest.

Received, 10th October 1988; Com. 8/03691H

## References

- 1 L. Horner and H. Hoffmann, *Angew. Chem.*, 1956, **68**, 473.
- 2 L. Horner and E. Jurgens, *Chem. Ber.*, 1957, **90**, 2184.
- 3 E. Howard and W. F. Olszewski, *J. Am. Chem. Soc.*, 1959, **81**, 1483.
- 4 F. Angolini and R. Bonnet, *Can. J. Chem.*, 1962, **40**, 181.
- 5 E. Schmitz, *Chem. Ber.*, 1958, **91**, 1488.
- 6 K. N. Houk, J. Sims, C. R. Watts, and L. J. Luskus, *J. Am. Chem. Soc.*, 1973, **95**, 7301.
- 7 K. M. Pietrusiewicz and A. Brandi, *Phosphorus Sulfur*, 1988, **37**, 4000.
- 8 R. Huisgen, H. Seidl, and I. Bruning, *Chem. Ber.*, 1969, **102**, 1102.