



A mild and efficient method for preparation of azides from alcohols using acidic ionic liquid [H-NMP]HSO₄

Abdol R. Hajipour^{a,b,*}, Asiyeh Rajaei^b, Arnold E. Ruoho^a

^a Department of Pharmacology, University of Wisconsin, Medical School, 1300 University Avenue, Madison, 53706-1532 WI, USA

^b Pharmaceutical Research Laboratory, Department of Chemistry, Isfahan University of Technology, Isfahan 84156, Islamic Republic of Iran

ARTICLE INFO

Article history:

Received 31 October 2008

Revised 26 November 2008

Accepted 26 November 2008

Available online 3 December 2008

Keywords:

Acidic ionic liquid

Azide

Alcohol

Chemoselective

ABSTRACT

We report here an efficient method for the synthesis and characterization of [H-NMP]HSO₄ and its application as an efficient catalyst and solvent for preparation of azides from corresponding alcohols under mild conditions. This processor showed high chemoselectivity for conversion of various alcohols to their corresponding azides.

© 2008 Elsevier Ltd. All rights reserved.

1. Introduction


Conversions of alcohols to their corresponding azides are very important and essential in functional group transformation. Alkyl azides have frequently been used for the synthesis of amino groups and for the construction of N-heterocycles.¹ There are a few reported methods for preparation of alkyl azide including using Mitsunobu reactions using hydrazoic acid as the azide source,² oxidation–reduction condensation using a quinone derivative and trimethylsilylazide (TMSN₃),³ *N*-(*p*-toluenesulfonyl) imidazole (Tslm)/tetrabutylammonium iodide (TBAI)/triethylamine (TEA),⁴ diphenylphosphorazidate (DPPA),⁵ zinc azide/bis pyridine complex,⁶ (DPPA)/1, 8-diaza bicyclo[5.4.0]undec-7-en,⁷ NaN₃/BF₃·ET₂O,⁸ and other useful methods.^{9,2f} Azides have been used for preparation of medicines such as azido pyrimidines and purines due to their anti HIV-1 biological activity,¹⁰ also 5-azido-5-deoxyxylofuranose has been employed for synthesis of sugar triazole derivatives as anti tuberculosis agents.¹¹ Protic acidic ionic liquids (ILs) have frequently been used as green solvents and catalysts in place of conventional organic solvents and acid catalysts.¹² In continuation of our ongoing projects to develop efficient methods for organic functional groups transformation,^{13–15} here we wish to report the synthesis and characterization of pyrrolidinium bisulfate [H-NMP]HSO₄, which is a new acidic ionic liquid,¹⁶ as a highly chemoselective, inexpensive, and efficient catalyst and also as a solvent

for conversion of alcohols to the corresponding azides using NaN₃ (Scheme 1).

2. Results and discussion

Initially to optimize the conversion of benzyl alcohol to benzyl azide, we tried the reaction under solvent-free conditions and in different solvents such as water, acetonitrile, and dichloromethane and under different temperatures including 60, 80, 100, and 120 °C for evaluating the acidity and applications of this Brønsted acid. Different amounts of acidic ionic liquid and sodium azide were also tried for optimization of the reaction conditions. We found that reaction at 120 °C and using 3 mmol of ionic liquid, 1 mmol of benzyl alcohol, and 1.5 mmol of sodium azide gave the best yield in shorter reaction time. We applied these conditions for synthesis of various azides from corresponding alcohols (Table 1).¹⁷

The procedure showed high chemoselectivity for azidation of secondary benzyl alcohols in the presence of primary, tertiary benzylic, and also aliphatic alcohols. As demonstrated in Table 1 in comparison to secondary alcohols with withdrawing groups the secondary alcohols with donor groups were reacted faster and in

























Scheme 1.

* Corresponding author. Tel.: +98 311 391 3262; fax: +98 311 391 2350.

E-mail address: haji@cc.iut.ac.ir (A.R. Hajipour).

Table 1

Conversion of various alcohols to corresponding azides with optimized procedure

Entry	Substrate	Product	Time (h)	Conversion (%) ^a
1			6	80
2			12	5
3			8	40
4			24	0
5			2	100
6			2	100
7			1	100
8			1	100
9			3	80
10			2	95
11			1.5	100

(continued on next page)

Table 1 (continued)

Entry	Substrate	Product	Time (h)	Conversion (%) ^a
12			1	100
13			1	100
14			1	100
15			1	100
16			4	40
17			2	100
18			1	90
19			8	0
20			1	100

^a Conversions were determined with TLC.

higher yields. Aliphatic alcohols were not converted to their azides under these conditions (entries 4 and 19) and using alcohols were recovered without any conversion even in longer reaction times. We believed that the proton of bisulfate is the proton donor due to its higher acidity.

3. Conclusion

In conclusion, this method is a novel and convenient method for conversion of various alcohols to the corresponding azides using novel and inexpensive acid ionic liquid [H-NMP]HSO₄. The high

yield, green chemistry, using inexpensive and straightforward isolation of the products are the advantages of this method over reported procedures. The method is chemoselective for conversion of secondary benzylic alcohols in the presence of primary benzylic and aliphatic alcohols and also for conversion of primary benzylic alcohols in the presence of aliphatic alcohols. In this method, the halogen groups in benzene rings were intact during the reaction.

Acknowledgments

We gratefully acknowledge the funding support received for this project from the Isfahan University of Technology (IUT), IR Iran (A.R.H.) and Grant GM 33138 (A.E.R.) from the National Institutes of Health, USA. Further financial support from Center of Sensor and green chemistry Research (IUT) is gratefully acknowledged.

References and notes

- (a) Brase, S.; Gil, C.; Knepper, K.; Zimmermann, V. *Angew. Chem., Int. Ed.* **2005**, *44*, 5188–5240; (b) Scriven, E. F. V.; Turnbull, K. *Chem. Rev.* **1988**, *88*, 297; (c) Boyer, J. H.; Canter, F. C. *Chem. Rev.* **1954**, *54*, 1; (d) Smith, P. A. S. In *Open Chain Nitrogen Compounds*; W.A. Benjamin: New York, 1966; Vol. II, p 211.
- (a) Mitsunobu, O.; Wada, M.; Sano, T. *J. Am. Chem. Soc.* **1972**, *94*, 679; (b) Hughes, D. L. *Org. React.* **1992**, *42*, 358–359; (c) Loibner, H.; Zbiral, E. *Helv. Chim. Acta* **1977**, *60*, 417; (d) Mitsunobu, O. *Synthesis* **1981**, *1*; (e) Hughes, D. L. *Org. Prep. Proced. Int.* **1996**, *28*, 127; (f) Saito, A.; Saito, K.; Tanaka, A.; Oritani, T. *Tetrahedron Lett.* **1997**, *38*, 3955; (g) Mizuno, M.; Shior, T. *Chem. Commun.* **1997**, 2165; (h) Fabiano, E.; Golding, B. T.; Sadeghi, M. M. *Synthesis* **1987**, 190; (i) Bessodes, M.; Abushanab, E.; Antonakis, K. *Tetrahedron Lett.* **1984**, *25*, 5899; (j) Mitsunobu, O. *Bull. Chem. Soc. Jpn.* **1967**, *40*, 4235; (k) Lee, S.-H.; Yoon, J.; Chung, S.-H.; Lee, Y.-S. *Tetrahedron* **2001**, *57*, 2139.
- Kurada, K.; Hayashi, Y.; Mukaiyama, T. *Tetrahedron* **2007**, *63*, 6358.
- Soltani Rad, M. N.; Behrouz, S.; KHalafi Nejad, A. *Tetrahedron Lett.* **2007**, *48*, 3445.
- Lal, B.; Pramanik, B. N.; Manhas, M. S.; Bose, A. K. *Tetrahedron Lett.* **1977**, 1977.
- Viaud, M. C.; Rollin, P. *Synthesis* **1990**, 130.
- Thompson, A. S.; Humphrey, G. R.; Demarco, A. M.; Mathre, D. J.; Grabowski, J. J. *J. Org. Chem.* **1993**, *58*, 5886.
- Sampath Kumar, H. M.; Subba Reddy, B. V.; Anjaneyulu, S.; Yadav, J. S. *Tetrahedron Lett.* **1998**, *39*, 7385.
- For example, see: (a) Mizuno, M.; Shioiri, T. *Chem. Commun.* **1997**, 2165; (b) Yu, C.; Liu, B.; Hu, L. *Org. Lett.* **2000**, *2*, 1959; (c) Iranpoor, N.; Firouzabadi, H.; Akhlaghinia, B.; Nowrouzi, N. *Tetrahedron Lett.* **2004**, *45*, 3291.
- Kim, H. O.; Park, Y. H.; Moon, H. R.; Jeong, L. S. *Bioorg. Med. Chem. Lett.* **2002**, 2403.
- Singh, B. K.; Yadav, A. K.; Kumar, B.; Gaikwad, A.; Sinha, S. K.; Chaturvedi, V.; Tripathi, R. P. *Carbohydr. Res.* **2008**, *343*, 1153.
- (a) Tamar, L. G.; Calum, J. *Chem. Rev.* **2008**, *108*, 206; (b) Johnson, K. E.; Richard, R. M.; Pagni, M.; Bartmess, J. *Monatsh. Chem.* **2007**, *138*, 1077.
- Hajipour, A. R.; Zarei, A.; Khazdooz, L.; Mirjalili, B. B. F.; Sheikhan, N.; Zahmatkesh, S.; Ruoho, A. E. *Synthesis* **2005**, *20*, 3644.
- (a) Hajipour, A. R.; Arbabian, M.; Ruoho, A. E. *J. Org. Chem.* **2002**, *67*, 8622; (b) Hajipour, A. R.; Ruoho, A. E. *Org. Prep. Proced. Int.* **2002**, *34*, 647; (c) Hajipour, A. R.; Mazloumi, G. *Synth. Commun.* **2002**, *32*, 23; (d) Hajipour, A. R.; Ruoho, A. E. *J. Chem. Res. (S)* **2002**, 547; (e) Hajipour, A. R.; Adibi, H.; Ruoho, A. E. *J. Org. Chem.* **2003**, *68*, 4553; (f) Hajipour, A. R.; Bageri, H.; Ruoho, A. E. *Bull. Korean Chem. Soc.* **2004**, *25*, 1238; (g) Hajipour, A. R.; Malakutikhah, M. *Org. Prep. Proced. Int.* **2004**, *364*, 647; (h) Hajipour, A. R.; Ruoho, A. E. *Sul. Lett.* **2002**, *25*, 151; (i) Hajipour, A. R.; Mirjalili, B. F.; Zarei, A.; Khazdooz, L.; Ruoho, A. E. *Tetrahedron Lett.* **2004**, *45*, 6607; (a) Hajipour, A. R.; Mahboubkhah, N. *J. Chem. Res. (S)* **1998**, 122; (b) Hajipour, A. R.; Mallakpour, E.; Adibi, H. *Chem. Lett.* **2000**, 460; (c) Hajipour, A. R.; Mallakpour, E.; Khoei, S. *Chem. Lett.* **2000**, 120; (d) Hajipour, A. R.; Mallakpour, E.; Khoei, S. *Synlett* **2000**, 740; (e) Hajipour, A. R.; Ruoho, A. E. *Org. Prep. Proced. Int.* **2005**, *37*, 298; (f) Hajipour, A. R.; Ruoho, A. E. *Org. Prep. Proced. Int.* **2005**, *37*, 279.
- Hajipour, A. R.; Kooshki, B.; Ruoho, A. E. *Tetrahedron Lett.* **2005**, *46*, 5503.
- Synthesis of acidic ionic liquid [H-NMP]HSO₄*: In a round-bottomed flask, to a cold mixture of 1-methyl-2-pyrrolidone, (0.99 g) in 15 ml of dichloromethane on an ice bath was added. One gram of concentrated sulfuric acid (98%, $d = 1.84$) was added dropwise. The reaction mixture was stirred at room temperature for 3 h, then the solvent was evaporated under reduce pressure to produce 1.97 g of [H-NMP]HSO₄ (100%). FT-IR (KBr, cm^{-1}) 2300–3600 (s, Br), 1660 (s), 1509 (m), 1302 (m), 1114 (w), 962 (w), 610 (w) cm^{-1} . ¹H NMR (500 MHz, D₂O): δ 1.4 (m, 2H, CH₂), 1.8 (t, 2H, CH₂), 2.2 (s, 3H, CH₃), 2.9 (t, 2H, CH₂). ¹³C NMR (500 MHz, D₂O): δ 22.22 (CH₂), 35.09 (CH₂), 42.10–43.51 (CH₂, CH₃), 180 (C=O) ppm.
- General procedure for the one-pot conversion of alcohol to azide*: In a 25 ml round-bottomed flask, a mixture of alcohol (1 mmol), NaN₃ (1.5 mmol), and ionic liquid [H-NMP]HSO₄ (3 mmol) was stirred on an oil bath at 120 °C. The progress of the reaction was checked with TLC (cyclohexane/ethyl acetate 70:30). When the reaction was completed, the mixture was extracted with cyclohexane (2 × 10 ml), filtered off the solid. The organic layer was washed with NaOH 5% (10 ml). The organic phase was dried with MgSO₄, and the solvent was evaporated under reduced pressure. The crude product was purified by column chromatography using silica gel (cyclohexane/ethyl acetate 70:30).