

Facile Synthesis of Optically Active anti- α,β -Dihydroxy Ester Derivatives

Teruaki MUKAIYAMA, Hiromi UCHIRO, Isamu SHIINA, and Shu KOBAYASHI

Department of Applied Chemistry, Faculty of Science,
Science University of Tokyo, Kagurazaka, Shinjuku-ku, Tokyo 162

Anti- α,β -dihydroxy thioesters are prepared with excellent enantioselectivities by the asymmetric aldol reaction between both achiral aldehydes and silyl enol ethers derived from α -benzyloxy thioesters in the presence of a chiral promoter consisted of chiral diamine coordinated tin(II) triflate and dibutyltindiacetate.

Optically active 1,2-diol units are widely distributed in natural products such as macrolides, polyethers and carbohydrates, etc. Recently several asymmetric oxidation reactions of olefins by using osmium tetroxide with a chiral ligand has been developed in high optical yields,^{1,2)} however, there still remain some severe problems such as the preparation of optically active anti-1,2-diols.

In the previous papers,^{3,4)} we have demonstrated that the asymmetric aldol reaction between both achiral silyl enol ethers of thioesters and aldehydes is performed with almost perfect diastereo- and enantioselectivities by using a new chiral promoter, combination of chiral diamine coordinated tin(II) triflate and tributyltinfluoride (dibutyltindiacetate). Then this asymmetric reaction was intended to apply to the reaction of silyl enol ether derived from α -benzyloxy thioester with aldehydes in order to introduce two hydroxy groups simultaneously accompanied with stereoselective carbon-carbon bond formation.⁵⁾

In the first place, the reaction of benzaldehyde with trimethylsilyl enol ether of S-ethyl 2-benzyloxyethanethioate⁶⁾ was carried out in dichloromethane by using a chiral promoter consisted of tin(II) triflate, (S)-1-methyl-2-[(N-1-naphthylamino)methyl]pyrrolidine and tributyltinfluoride. The reaction smoothly proceeded at -78 °C to afford the corresponding aldol-type adduct in 69% yield with anti preference (syn/anti=26/74). The enantiomeric excesses of syn and anti aldols proved to be 30% and 97%, respectively. It is noteworthy to point out that the anti aldol-type adduct was predominantly obtained with excellent ee in this case, while in the case of asymmetric aldol reaction of silyl enol ether of S-ethyl propanethioate with aldehydes, syn aldol-type adducts were obtained (syn/anti=100/0. syn aldol=>98%ee).⁴⁾

Next, several chiral diamines were examined in order to improve the diastereoselectivity. When (S)-1-methyl-2-[(1-piperidin-1-yl)methyl]pyrrolidine was employed, the aldol-type adduct was obtained in 54% chemical yield with excellent diastereo- and enantioselectivity (syn/anti=1/99, anti aldol=97%ee). Further, the chemical yield was improved without losing the stereoselectivities when combination of (S)-1-ethyl-2-[(1-piperidin-1-yl)methyl]pyrrolidine and dibutyltindiacetate was employed as a promoter (Table 1).








Table 1. Effect of Chiral Diamine

Chiral diamine	Additive	Yield / %	syn : anti	ee / % (anti)
	$n\text{Bu}_3\text{SnF}$	69	26 : 74	97
	R = Me, $n\text{Bu}_3\text{SnF}$	70	1 : 99	97
	Et, $n\text{Bu}_3\text{SnF}$	54	1 : 99	>98
	$n\text{Pr}$, $n\text{Bu}_3\text{SnF}$	54	1 : 99	>98
	$n\text{Pen}$, $n\text{Bu}_3\text{SnF}$	38	1 : 99	97
	R = Me, $n\text{Bu}_2\text{Sn}(\text{OAc})_2$	74	1 : 99	96
	Et, $n\text{Bu}_2\text{Sn}(\text{OAc})_2$	83	1 : 99	96

The results of the present asymmetric aldol reaction by using several kinds of aldehydes such as aromatic, aliphatic, α,β -unsaturated aldehydes and a dienal, are shown in Table 2. In every case, anti- α,β -dihydroxy thioesters are obtained in high yields with excellent diastereo- and enantioselectivities.

A typical experimental procedure is described for the reaction of silyl enol ether derived from S-ethyl 2-benzyloxyethanethioate with benzaldehyde; to a solution of tin(II) triflate (0.4 mmol) and (S)-1-ethyl-2-[(piperidin-1-yl)methyl]pyrrolidine (0.48 mmol) in dichloromethane (1 ml) was added dibutyltin diacetate (0.44 mmol) at room temperature. The mixture was stirred for 30 min and then cooled to $-78\text{ }^\circ\text{C}$. Dichloromethane solutions (0.5 ml each) of silyl enol ether of S-ethyl 2-benzyloxyethanethioate (0.4 mmol) and benzaldehyde (0.27 mmol) were successively added. The reaction mixture was further stirred for 20 h, and quenched with aqueous sodium hydrogen carbonate. After usual work up, S-ethyl 2,3-dihydroxy-3-phenylpropanethioate was obtained in 83% yield (syn/anti=1/99, anti aldol=96%ee).⁷⁾

Thus, the present asymmetric aldol-type reaction makes it possible to introduce two hydroxy groups in 1,2-trans position stereoselectively during new carbon-carbon bond formation. It is noteworthy to mention that thus obtained aldol-type adducts, optically active anti- α,β -dihydroxy ester derivatives, are generally difficult to prepare by the conventional methods of asymmetric oxidation because it is hard to obtain starting materials, cis- α,β -unsaturated ester equivalents.⁸⁾ Further investigation concerning application of this asymmetric aldol reaction to synthesis of natural products are in progress.

Table 2. Synthesis of anti- α,β -Dihydroxy Thioesters

Aldehyde	Yield / %	syn : anti	ee / % ^{a)} (config.)
PhCHO	83	1 : 99	96 (2R, 3R)
CH ₃ CH ₂ CHO	72	2 : 98	97
c-C ₆ H ₁₁ CHO	59	9 : 91	96
(E)-PhCH=CHCHO	88	2 : 98	98
(E)-CH ₃ CH=CHCHO	85	2 : 98	97
(E,E)-CH ₃ CH=CHCH=CHCHO	83	2 : 98	95


a) Determined by HPLC analysis of the corresponding acetyl derivatives.

The present research was partially supported by Grant-in-Aids for Scientific Research No. 01649008 from Ministry of Education, Science and Culture.

References

- 1) E. N. Jacobsen, I. Marco, W. S. Mungall, G. Schroder, and K. B. Sharpless, *J. Am. Chem. Soc.*, **110**, 1968 (1988); J. S. M. Wai, I. Marco, S. Svendsen, M. G. Finn, E. N. Jacobsen, and K. B. Sharpless, *ibid.*, **111**, 1123 (1989); K. Tomioka, M. Nakajima, and K. Koga, *ibid.*, **109**, 6213 (1987); T. Yamada and K. Narasaka, *Chem. Lett.*, **1986**, 131; M. Tokles and J. K. Snyder, *Tetrahedron Lett.*, **27**, 3951 (1986); M. Hiramata, T. Oishi, and S. Ito, *J. Chem. Soc., Chem. Commun.*, **1989**, 665.
- 2) E. J. Corey, P. D. Jardine, S. Virgil, P-W Yuen, and R. D. Connell, *J. Am. Chem. Soc.*, **111**, 9243 (1989).
- 3) S. Kobayashi and T. Mukaiyama, *Chem. Lett.*, **1989**, 297; S. Kobayashi, T. Sano, and T. Mukaiyama, *ibid.*, **1989**, 1319; S. Kobayashi, Y. Fujishita, and T. Mukaiyama, *ibid.*, **1989**, 2069; T. Mukaiyama and S. Kobayashi, *J. Organomet. Chem.*, **382**, 39 (1990).
- 4) T. Mukaiyama, H. Uchiro, and S. Kobayashi, *Chem. Lett.*, **1989**, 1001; T. Mukaiyama, H. Uchiro, and S. Kobayashi, *ibid.*, **1989**, 1757; T. Mukaiyama, S. Kobayashi, H. Uchiro, and I. Shiina, *ibid.*, **1990**, 129.

- 5) It was reported from this laboratory that chiral diamine coordinated tin(II) enolate of 3-(2-benzyloxyacetyl)-thiazolidine-2-thione reacted with aldehydes to afford the corresponding anti aldol-type adducts in good diastereo- and enantioselectivities; T. Mukaiyama and N. Iwasawa, *Chem. Lett.*, **1984**, 753.
- 6) This silyl enol ether was prepared from S-ethyl 2-benzyloxyethanethioate by the usual procedure (1. LDA/THF, -78 °C, 2. TMSCl; 80%yield, E/Z=72/28). In this paper, the stereochemical description E and Z of silyl enol ethers of thioesters are employed as they correspond to ketene silyl acetals.
- 7) The absolute configuration was determined by the comparison of the authentic tribenzoate **1** after the following derivations.



- 8) Recently, Corey et al. reported that anti- α,β -diols were difficult to prepare in high ees via the asymmetric dioxyosmylation, according to the consideration of their mechanistic model; ref. 2).
- 9) Y. Ohgo, J. Yoshimura, M. Kono, and T. Sano, *Bull. Chem. Soc. Jpn.*, **42**, 2957 (1969).

(Received April 5, 1990)