

Enantioselective synthesis of the C14–C25 portion of the cytotoxic natural product, amphidinolide B1

Duck-Hyung Lee ^{*} and Man-Dong Rho


Department of Chemistry, Sogang University, Seoul 121-742, South Korea

Received 21 October 1999; revised 5 January 2000; accepted 28 January 2000

Abstract


The C14–C25 portion of the cytotoxic natural product, amphidinolide B1 (**1**), was synthesized enantioselectively using Myers asymmetric alkylation protocol, epoxide opening with higher order cuprate, Sharpless asymmetric epoxidation of allylic alcohol **17**, and Sharpless asymmetric dihydroxylation of **24** as key steps. © 2000 Elsevier Science Ltd. All rights reserved.

Amphidinolides A–S have recently been isolated from dinoflagellates, genus *Amphidinium*,¹ which is symbiotic with Okinawan marine flatworms of the genus *Amphiscolops* sp. and generally exhibited potent toxicities against cancer tumor cell lines.² This family has been a challenging target for synthetic chemists in recent years³ and some synthesis of the segments has been reported, including amphidinolide B1 (**1**).⁴ In our program toward the synthesis of the C14–C25 portion **2** of amphidinolide B1 (**1**), C1–C28 and C20–C25 fragments have already been synthesized in our laboratory⁵ and we report herein the enantioselective synthesis of segments **3a**, **3b** and the coupling of these two components via a lithium–halogen exchange for the C14–C25 portion **2** of amphidinolide B1 (**1**). The features of the synthesis are: Myers asymmetric alkylation protocol for the chirality at C23; epoxide opening with higher order cuprate for the C18 asymmetric center; Sharpless asymmetric epoxidation for the chirality at C16; and Sharpless asymmetric dihydroxylation for the C21 and C22 asymmetric center.



^{*} Corresponding author.


First, pseudoephedrin amide **4** was treated with alkyl iodide **5**, using the Myers protocol⁶ to give the alkylation product **6** in 68% yield as a 12.8:1 ratio of diastereomers (determined by 500 MHz ¹H NMR) (Scheme 1). Alkyl iodide **5** was prepared in three steps from ethyl (*S*)-lactate via protection of a secondary alcohol with *p*-methoxybenzyl-2,2,2-trichloroacetimidate in the presence of TsOH, reduction of ester functionality with LiAlH₄ and iodination of the corresponding primary alcohol with triphenylphosphine–iodine in a 56% overall yield. Amide **6** was hydrolyzed with a 3.2N aqueous NaOH solution at reflux for 24 h to the carboxylic acid **7** and the acid **7** was reduced with BH₃–DMS in THF at 0°C for 40 min to the primary alcohol **8** in a 58% two-step isolation yield. Swern oxidation of primary alcohol **8** afforded aldehyde **9** and then the aldehyde **9** was treated with CrCl₂ and CHI₃ in THF at rt for 7 h using the Takai protocol⁷ to introduce the C21,C22 *trans*-vinyl iodide functionality of the C21–C25 fragment **3a** of amphidinolide B1 (**1**) in 21% overall yield from pseudoephedrine amide **4**. The synthetic route presented herein has several advantages over the previous one published in our laboratory,^{5b} such as shorter reaction steps, more reliable results in multi-gram scale, and the purity of the product (almost pure versus 6:1 diastereomeric mixture).



Scheme 1. Synthesis of C21–C25 portion of amphidinolide B1 (**1**). (a) LDA, LiCl, THF, –78°C, 1 h, 0°C, 15 min, rt, 5 min; **5**, 0°C to rt, 3 days, 68%; (b) NaOH, MeOH, *t*-BuOH, reflux, 24 h, 71%; (c) BH₃–DMS, THF, 0°C, 40 min, 82%; (d) DMSO, (COCl)₂, CH₂Cl₂, –78°C; DIPEA, –50°C, 1 h, quant.; (e) CrCl₂, CHI₃, THF, rt, 7 h, 53%


The other fragment **3b** was synthesized as described below (Scheme 2). Bromination of the crotyl alcohol **10** with Br₂ in CCl₄ at –5°C for 2 h and elimination of dibromide **11** with LDA and HMPA in THF at –78°C for 40 min were followed by protection of the corresponding primary alcohol **12** with TBSCl and TEA in CH₂Cl₂ in the presence of a catalytic amount of DMAP to give (*E*)-vinyl bromide **13** in a 57% three-step overall yield⁸. Treatment of vinyl bromide **13** with *t*-BuLi in ether and reaction of the corresponding vinyl lithium reagent with CuCN provided the higher order cuprate, which reacted smoothly with epoxide **14**⁹ in diethyl ether at rt for 12 h to afford secondary alcohol **15** regioselectively in 91% yield. After protection of secondary alcohol **15** with TIPSOTf and TEA in CH₂Cl₂ at 0°C for 10 min, the compound **16** was subjected to selective desilylation with HF–Pyr leading to the corresponding allylic alcohol **17** in a 87% two-step yield. Sharpless asymmetric epoxidation¹⁰ of allylic alcohol **17** using D-(–)-DET, Ti(O-*i*Pr)₄ and TBHP in CH₂Cl₂ at –20°C for 7 h afforded the epoxy alcohol **18** in 99% yield as a 11:1 ratio of diastereomers (determined by 500 MHz ¹H NMR). The epoxy alcohol **18** was treated with TBSCl and TEA in CH₂Cl₂ in the presence of DMAP at rt for 4 h, and the epoxide **19** was opened with the higher order cuprate derived from methyl lithium to give tertiary alcohol **20** regioselectively in a 73% two-step isolation yield. Protection of tertiary alcohol **20** with TBSOTf and TEA in CH₂Cl₂ at rt for 2 h (85% yield) was followed by hydrogenolysis of the benzyl ether **21** by the

use of hydrogen (1 atm) and 10% Pd/C in ethanol to afford alcohol **22** in 90% yield. Swern oxidation of primary alcohol **22** provided the C14–C20 portion **3b** of amphidinolide B1 (**1**) in quantitative yield.



Scheme 2. Synthesis of C14–C20 portion of amphidinolide B1 (**1**). (a) Br_2 , CCl_4 , $-5^\circ C$, 2 h, 92%; (b) LDA, HMPA, THF, $-78^\circ C$, 40 min, 67%; (c) TBSCl, DMAP, TEA, CH_2Cl_2 , rt, 40 min, 92%; (d) $t-BuLi$, ether, $-78^\circ C$, 1 h; CuCN, ether, 20 min; **14**, THF, $-78^\circ C$ to rt, 12 h, 91%; (e) TIPSOtF, TEA, CH_2Cl_2 , $0^\circ C$, 10 min, quant.; (f) HF–Pyr, pyr, THF, $0^\circ C$ to rt, 2 h, 87%; (g) $Ti(O^i-Pr)_4$, D-(–)-DET, 4 Å MS, TBHP, CH_2Cl_2 , $-20^\circ C$, 7 h, 99%; (h) TBSCl, DMAP, TEA, CH_2Cl_2 , $0^\circ C$ to rt, 4 h, 73%; (i) $(CH_3)_2Cu(CN)Li_2$, ether, $-78^\circ C$ to rt, 12 h, quant.; (j) TBSOTf, TEA, CH_2Cl_2 , $0^\circ C$ to rt, 2 h, 85%; (k) H_2 , Pd/C, EtOH, rt, 20 min, 90% (l) DMSO, $(COCl)_2$, CH_2Cl_2 , $-78^\circ C$, 15 min; TEA, rt, 20 min, quant.

With two fragments **3a** and **3b** in hand, coupling reactions were followed (Scheme 3). Lithium–halogen exchange of vinyl iodide **3a** with $t-BuLi$ at $-78^\circ C$ and addition of aldehyde **3b** to the resulting vinyl lithium reagent at $-78^\circ C$ for 15 min provided the allylic alcohol **23** in a 73% yield as a mixture of diastereomers at C20. Subsequent oxidation of allylic alcohol **23** with Dess–Martin periodinane¹¹ in Pyr: CH_2Cl_2 (ca. 1:4) at rt for 2 days gave a conjugated ketone **24** in a 86% isolation yield. Sharpless asymmetric dihydroxylation of conjugated ketone **24** with AD-mix- α in the presence of methanesulfonamide¹² in $t-BuOH:H_2O$ (1:1) at $0^\circ C$ afforded the diol **2**¹³ in 35% yield as a 6:1 ratio of diastereomers (determined by 500 MHz 1H NMR) along with the recovered starting material **24** (ca. 35%) after extensive investigations.



Scheme 3. Coupling of **3a** and **3b**. (a) **3a**, $t-BuLi$, $-78^\circ C$, 15 min; **3b**, THF, $-78^\circ C$, 20 min, 73%; (b) Dess–Martin periodinane, Pyr, CH_2Cl_2 , rt, 2d, 86%; (c) AD-mix- α , $CH_3SO_2NH_2$, $t-BuOH:H_2O$ (1:1), $0^\circ C$, 1.5 days, 35%

Although further optimization might still be necessary, enantioselective synthesis of C14–C25 portion **2** of amphidinolide B1 (**1**) was completed in 5.5% overall yield via a 15-step sequence starting from (*E*)-crotyl alcohol **10**.

Acknowledgements

This research was assisted financially by KOSEF (Korea Science and Engineering Foundation, 971-0302-012-2 and 98-05-01-04-01-3).

References

1. Kobayashi, J.; Ishibashi, M. *Chem. Rev.* **1993**, 93, 1753.
2. For the isolation of amphidinolides, see: (a) Kobayashi, J.; Takahashi, M.; Ishibashi, M. *Tetrahedron Lett.* **1996**, 37, 1449 (Amphidinolide Q). (b) Kobatyssshi, J.; Ishibashi, M.; Takahashi, M. *Tetrahedron* **1997**, 53, 7827 (Amphidinolide R & S) and references cited therein.
3. (a) Maleczka Jr., R. E.; Terrell, L. R.; Ward, J. S.; Joseph, S. W. *Tetrahedron Lett.* **1999**, 40, 3097. (b) Chakraborty, T. K.; Suresh, V. R. *Tetrahedron Lett.* **1998**, 39, 7775. (c) Chakraborty, T. K.; Suresh, V. R. *Tetrahedron Lett.* **1998**, 39, 9109. (d) Williams, D. R.; Kissel, W. S. *J. Am. Chem. Soc.* **1998**, 120, 11 198. (e) Kobatyssshi, J.; Tsuda, M.; Hatakeyama, A. *J. Chem. Soc., Perkin Trans. 1* **1998**, 149. (f) Gerald, P.; Gregory, J. H. *Tetrahedron Lett.* **1998**, 39, 703. (g) Kobatyssshi, J.; Hatakeyama, A.; Tsuda, M. *Tetrahedron* **1998**, 54, 697. (h) Kobatyssshi, J.; Ishiyama, I.; Ishibashi, M. *Chem. Pharm. Bull.* **1996**, 44, 1819.
4. (a) Chakraborty, T. K.; Thippeswamy, D.; Suresh, V. R.; Jayaprakash, S. *Chem. Lett.* **1997**, 563; Chakraborty, T. K.; Suresh, V. R. *Chem. Lett.* **1997**, 56. (b) Lee, D. H.; Lee, S. W. *Tetrahedron Lett.* **1997**, 38, 7909. (c) Lee, D. H.; Rho, M. D. *Bull. Korean Chem. Soc.* **1998**, 19, 386. (d) Pattenden, G.; Cid, M. B. *Synlett.* **1998**, 540. (e) Nishiyama, S.; Ohi, K.; Shima, K.; Hamada, K.; Saito, Y.; Yamada, N.; Ohba, S. *Bull. Chem. Soc. Jpn.* **1998**, 71, 2433. (f) Chakraborty, T. K.; Thippeswamy, D. *Synlett.* **1999**, 150. (g) Myles, D. C.; Eng, H. M. *Tetrahedron Lett.* **1999**, 40, 2275; Myles, D. C.; Eng, H. M. *Tetrahedron Lett.* **1999**, 40, 2279. (h) Kobatyssshi, J.; Ishiyama, H.; Takemura, T.; Tsuda, M. *Tetrahedron* **1999**, 55, 4583.
5. (a) Lee, D. H.; Lee, S. W. *Tetrahedron Lett.* **1997**, 38, 7909. (b) Lee, D. H.; Rho, M. D. *Bull. Korean Chem. Soc.* **1998**, 19, 386.
6. (a) Myers, A. G.; Yang, B. H.; Chen, H.; Mckinstry, L.; Kopecky, D. J.; Gleason, J. L. *J. Am. Chem. Soc.* **1997**, 119, 6496. (b) Myers, A. G.; Mckinstry, L. *J. Org. Chem.* **1996**, 61, 2428.
7. Takai, K.; Nitta, K.; Utimoto, K. *J. Am. Chem. Soc.* **1986**, 108, 7408.
8. Corey, E. J.; Bock, M. G.; Kozikowski, A. P.; Rama Rao, A. V. *Tetrahedron Lett.* **1978**, 19, 1051.
9. Epoxide **14** was prepared in a 48% three-step overall yield from L-aspartic acid (**23**) following the literature procedure (see: Rapoport, H.; Frick, J. A.; Klassen, J. B.; Bathe, A.; Abramson, J. M. *Synthesis* **1992**, 621).
10. Sharpless, K. B.; Gao, Y.; Hanson, R. M.; Klunder, J. M.; Ko, S. Y.; Masamune, H. *J. Am. Chem. Soc.* **1987**, 109, 5765.
11. Martin, J. C.; Dess, D. B. *J. Org. Chem.* **1983**, 48, 4156.
12. Sharpless, K. B.; Amberg, W.; Bennani, Y. L.; Crispino, G. A.; Hartung, J.; Jeong, K. S.; Kwong, H. L.; Morikawa, K.; Wang, Z. M.; Xu, D.; Zhang, X. L. *J. Org. Chem.* **1992**, 57, 2768.
13. Spectral data for **2**. $R_f=0.49$ (EtOAc:n-hexane, 1:20); $[\alpha]_D^{24} -7.692$ (c 0.26, CHCl₃); FT-IR (CDCl₃) 3443.15 (OH), 2926.15, 2855.81, 1717.09; ¹H NMR (500 MHz) δ 7.260 (d, J=8.5 Hz, 2H), 6.865 (d, J=8.5 Hz, 2H), 4.614 (m, 1H), 4.553 (d, J=11 Hz, 1H), 4.337 (d, J=11.5 Hz, 1H), 4.157 (m, 1H), 3.798 (s, 3H), 3.762 (dd, J=9.5, 4 Hz, 1H), 3.729 (d, J=4.5 Hz, 1H), 3.389 (dd, J=9.5, 8 Hz, 1H), 2.950 (dd, J=16, 6 Hz 1H), 2.811 (dd, J=16, 5.5 Hz, 1H), 2.030 (m, 1H), 1.861–1.732 (m, 4H), 1.662–1.158 (m, 2H), 1.256 (s, 3H), 1.208 (d, J=6 Hz, 3H), 1.055 (s, 21H), 0.968 (d, J=7 Hz, 3H), 0.954 (d, J=7 Hz, 3H), 0.887 (s, 9H), 0.878 (s, 9H), 0.088 (s, 6H), 0.034, 0.030 (s, s, 6H); ¹³C NMR (125 MHz) δ 209.506, 130.756, 129.325, 113.743, 77.890, 74.984, 72.143, 69.837, 66.707, 64.516, 55.249, 47.961, 46.803, 44.737, 40.826, 34.204, 27.052, 26.167, 20.037, 18.244, 15.633, 12.999, 12.715, –1.610, –1.659, –5.429; anal. calcd for C₄₅H₈₈O₈Si₃: C, 64.23; H, 10.54. Found: C, 64.27; H, 10.53.