CHEMISTRY LETTERS, pp. 585-588, 1985.

SYNTHESIS OF α -AMINO KETONES FROM α , β -EPOXY SULFOXIDES

Tsuyoshi SATOH, Youhei KANEKO, Kiichi SAKATA, and Koji YAMAKAWA^{*} Faculty of Pharmaceutical Sciences, Science University of Tokyo, Ichigaya-funagawara-machi, Shinjuku-ku, Tokyo 162

Various kinds of α -amino ketones and α -arylamino ketones are synthesized in good yields by aminolysis of α , β -epoxy sulfoxides which are easily prepared from alkylated chloromethyl phenyl sulfoxide and carbonyl compounds.

α-Amino ketones (<u>2</u>) are very important and versatile compounds in synthetic organic chemistry. They are useful intermediates for drugs^{1,8}) or as building blocks for nitrogen containing heterocyclic compounds.²) In spite of the impressive development of the methods for the synthesis of β-amino ketones, such as Mannich reaction³) or Michael addition of amines to α,β-unsaturated carbonyl compounds,⁴) only limited numbers of methods for the construction of α-amino ketones (<u>2</u>) are reported. Aminolysis of α-halo ketones⁵) usually gives poor yields. Reduction of acyl cyanides⁶ gives only one type (RCOCH₂NH₂) of α-amino ketones. The Neber rearrangement⁷) gives only α-amino ketones having no substituents on the nitrogen atom. Aminolysis of enediol silyl ether¹ was reported to give about 45% yield of α-amino ketones. Aminolysis of epoxy-ether is good method, and practically used in a preparation of some drugs but only syntheses of phenyl α-amino ketones were reported.⁸)

Recently we reported a method for the synthesis of dialkyl ketones and α -sulfenylated carbonyl compounds from α,β -epoxy sulfoxides (<u>1</u>).⁹⁾ In the course of the studies on the synthesis of α -substituted carbonyl compounds from α,β -epoxy sulfoxides (<u>1</u>), here a new and efficient method for the synthesis of α -amino ketones (<u>2</u>) is reported according to Eq. 1.



 α,β -Epoxy sulfoxides (<u>1</u>) are very easily prepared in good overall yields from alkyl halides (R¹), carbonyl compounds (R², R³), and chloromethyl phenyl sulfoxide.¹⁰) We have found that the β -carbon of the α,β -epoxy sulfoxides is

| Run | Epoxy sulfoxide | Amine ^{a)} | Solvent | Temp | Time | α-Amino ketones | Yield ^{b)} % |
|-----|--------------------|---------------------|---------|--------|---------|---------------------|--------------------------|
| 1 | <u>3</u> L | А | - | r.t. | 3 h | • ^ | 97 |
| | Р | Α | - | r.t. | 21 h | | 100 |
| 2 | <u>3</u> L | в | - | r.t. | 4 h | Pn | 100 |
| | P | В | - | r.t. | 23 h | | 99 |
| | P | В | - | 80 °C | 90 min | ~~~Y | 90 |
| | Р | В | DMSO | 80 °C | 100 min | Pfi | 100 |
| | Р | в | HMPA | 80 °C | 150 min | | 94 |
| 3 | <u>3</u> L | с | HMPA | r.t. | 27 h | | 89 |
| | P | с | HMPA | 110 °C | 40 min | Ph Ph | 100 |
| 4 | <u>3</u> L | D | - | r.t. | 8 d | | 96 |
| | P | D ^{c)} | DMSO | 50 °C | 31 h | | 44 |
| 5 | <u>3</u> L | Ed) | DMSO | 60 °C | 3 h | O N ^{CH} 3 | 53 |
| | Р | E ^{d)} | DMSO | 60 °C | 5 h | | 52 |
| 6 | <u>3</u> L | F ^c) | DMSO | 55 °C | l h | | 72 |
| | Р | F ^{c)} | DMSO | 55 °C | l h | NHCh2Ph | 73 |
| 7 | <u>4</u> L | A | - | 100 °C | 2 h | | 97 |
| | Р | A | - | 100 °C | 12 h | | 85. |
| 8 | <u>4</u> L | G | - | 80 °C | l h | | 100 |
| | Р | G | - | 80 °C | 6 h | 。~ ∟ | 85 |
| 9 | <u>4</u> L | c ^{e)} | DMSO | 100 °C | 7 h | | 94 |
| | Ρ | c ^{e)} | DMSO | 100 °C | 20 h | | 80 |
| 10 | <u>5</u> | A | - | 100 °C | 6 h | | 76 |
| 11 | <u>5</u> | G | - | 80 °C | 4 h | | 100 |
| 12 | <u>5</u> | c ^{e)} | DMSO | 100 °C | 30 h | | 48 |
| | | | | | | | |

Table 1. α -Amino ketones from α . β -epoxy sulfoxides with amines

a) A: piperidine, B: morpholine, C: 1,2,3,4-tetrahydroisoquinoline, D: diethylamine, E: N-(3-hydroxy-4-methoxy)benzyl-N-methylamine, F: benzylamine, G: pyrrolidine.
b) Isolated yields after silica gel column chromatography.
c) 45 equiv. of amine was used.
d) 4 equiv. of amine was used.
e) 40 equiv. of amine was used.
e) 40 equiv.

very reactive to various kinds of nucleophiles⁹⁾ including amines or aromatic amines to afford α -substituted ketones in good yields under mild conditions.

The results of the reaction of the α,β -epoxy sulfoxides <u>3</u> (R¹ = n-hexyl, R², R³ = phenyl, H), <u>4</u> (R¹ = cyclohexyl, R², R³ = p-chlorophenyl, H), and <u>5</u> (R¹ = benzyl, R², R³ = -(CH₂)₅-) with various kinds of amines are summarized in Table 1. The epoxides <u>3</u> and <u>4</u> have two isomers (L and P), respectively, but the stereochemistries have not yet been determined. The reaction were usually conducted in DMSO or HMPA (0.5 M concentration of epoxy sulfoxide and 10 equiv. of amines are used unless otherwise noted) or in amine without solvent in the case of volatile amines under nitrogen atmosphere.

As shown in Table 1, many kinds of α -amino ketones are synthesized from the epoxy sulfoxides in good yields. The amine having a complex structure (Run 5) can be used to afford α -amino ketone, though the yield is moderate, which implies that this method would have practical value. Isomers L and P showed some differences in reactivity toward the amines. The epoxides having cyclohexane ring in R¹ (<u>4</u>) showed less reactivity than those of normal alkyl group in R¹ (<u>3</u>) due to steric hinderance on the β carbon of the epoxide by cyclohexyl group (see Run 1 and 7). The dipolar aprotic solvent known to be effective for nucleophilic bimolecular reaction¹¹) was used in this reaction. Run 3 shows that DMSO is slightly more effective than HMPA. Monoalkylamines usually gave a complex mixture but only benzylamine afforded an α -amino ketone in about 70% yield (Run 6).

| Epoxy sulfoxide | Aromatic amine (equiv | .) ^{a) Temp} | Time | α-Anilino ketones | Yield ^b) | | | | | |
|--------------------|--------------------------|-----------------------|------|-------------------|----------------------|--|--|--|--|--|
| <u>3</u> L | н (50) | 100 °C | l h | 0 | 97 | | | | | |
| Р | н (50) | 100 °C | 2 h | Ph | 81 | | | | | |
| <u>3</u> L | I (90) | 100 °C | 2 h | O H H NDb | 73 | | | | | |
| P | I (90) | 100 °C | 2 h | | 56 | | | | | |
| <u>3</u> L | J (5) | 100 °C | 3 h | | 63 | | | | | |
| P | J (5) | 100 °C | 5 h | | 47 | | | | | |
| | | | | | | | | | | |

Table 2. α -Arylamino ketones from α , β -epoxy sulfoxides with aromatic amines

a) H: aniline, I: N-methylaniline, J: p-aminoacetanilide; The reaction was conducted in HMPA and the concentration of $\underline{3}$ was 0.4 M. b) Isolated yields after silica gel column chromatography.

The results of the reaction of <u>3</u> with aromatic amines are summarized in Table 2. As the nucleophilicity of aromatic amines is decreased by the delocalization of the nitrogen lone pair of electrons, the reaction of aromatic amines with the α,β -epoxy sulfoxide required a higher temperature than those of amines and the reaction gave moderate yields.

The other methods reported to be effective for opening of epoxides, such as aminosilanes 12 or reactions at alumina surfaces 13 were not effective in this reactions.

The authors are grateful to Dr. Hiroshi Hara of this department for a generous gift of amines.

- References
 - H. M. Fischler, H. G. Heine, and W. Hartmann, Ger. Offen. 2145419; Chem. Abstr., <u>78</u>, 159051z (1973).
 - 2) D. Mayer, "Houben-Weyl, Methoden der Organischen Chemie," 4 th ed, Vol. 7/2c, p. 2251 (Thieme, Stuttgart 1977).
 - 3) M. Tramontini, Synthesis, 1973, 703.
 - 4) Y. Ogata, A. Kawasaki, and I. Kishi, J. Chem. Soc., B, <u>1968</u>, 703.
 - 5) S. L. Friess and H. D. Baldridge, J. Am. Chem. Soc., <u>78</u>, 2482 (1956);
 H. Berbalk, K. Eichinger, and E. Deeker, Monatsh. Chem., <u>107</u>, 401 (1976).
 - 6) C. G. Stuckwisch, J. Org. Chem., <u>37</u>, 318 (1972); A. Pfaltz and S. Anwar, Tetrahedron Lett., <u>25</u>, 2977 (1984).
 - 7) H. E. Baumgarten and J. M. Petersen, J. Am. Chem. Soc., <u>82</u>, 459 (1960);
 C. Obrien, Chem. Rev., <u>64</u>, 81 (1964).
- 8) Y. J. L'Italien, U. S. 3171858; Chem. Abstr., <u>62</u>, 14575c (1965); A. Wander, Swiss 401054; Chem. Abstr., <u>65</u>, 3835e (1966); C. L. Stevens and R. W. Fleming, Fr. 1447116; Chem. Abstr., <u>66</u>, 104826c (1967); H. Schulz, E. Jassmann, and R. Kowarsch, Ger. (East) 50619; Chem. Abstr., <u>67</u>, 21614b (1967); C. L. Stevens and C. H. Chang, J. Org. Chem., <u>27</u>, 4392 (1962); C. L. Stevens, R. D. Elliot, and B. L. Winch, J. Am. Chem. Soc., <u>85</u>, 1464 (1963); C. L. Stevens, M. E. Munk, and C. H. Chang, J. Org. Chem., <u>29</u>, 3146 (1964); C. L. Stevens, A. Thuillier, and F. A. Daniher, ibid., <u>30</u>, 2962 (1965); C. L. Stevens, A. B. Ash, A. Thuillier, J. H. Amin, A. Balys, W. E. Dennis, J. P. Dickerson, R. P. Glinski, H. T. Hanson, M. D. Pillai, and J. W. Stoddard, ibid., <u>31</u>, 2593 (1966); C. L. Stevens, A. Thuillier, K. G. Taylor, F. A. Daniher, J. P. Dickerson, H. T. Hanson, N. A. Nielsen, N. A. Tikotkar, and R. M. Weier, ibid., <u>31</u>, 2601 (1966).
- 9) T. Satoh, Y. Kaneko, T. Kumagawa, T. Izawa, K. Sakata, and K. Yamakawa, Chem. Lett., <u>1984</u>, 1957.
- 10) T. Durst, J. Am. Chem. Soc., <u>91</u>, 1034 (1969); T. Durst, K-C Tin, F. de Reinach-Hirtzbach, J. M. Decesare, and M. D. Ryan, Can. J. Chem., <u>57</u>, 258 (1978).
- 11) A. J. Parker, Chem. Rev., <u>69</u>, 1 (1969); E. Juaristl and J. D. Reyna, Tetrahedron Lett., <u>25</u>, 3521 (1984).
- 12) A. Papini, A. Ricci, M. Taddei, G. Seconi, and P. Dembech, J. Chem. Soc., Perkin Trans. 1, 1984, 2261.
- 13) G. H. Posner, D. Z. Rogers, C. M. Kinzig, and G. M. Gurria, Tetrahedron Lett., <u>1975</u>, 3597; G. H. Posner and D. Z. Rogers, J. Am. Chem. Soc., <u>99</u>, 8208, 8214 (1977).

(Received February 13, 1985)