

Accelerated synthesis of 3,5-disubstituted 4-amino-1,2,4-triazoles under microwave irradiation

Fouad Bentiss,^a Michel Lagrenée^a and Didier Barbry^{b,*}

^aLaboratoire de Cristallochimie et Physicochimie du Solide, CNRS UPRESA 8012, ENSCL, B.P. 108, F-59652 Villeneuve d'Ascq Cedex, France

^bLaboratoire d'Ingénierie Moléculaire, UPRES EA 2699, Université des Sciences et Technologies de Lille, F-59655 Villeneuve d'Ascq Cedex, France


Received 1 October 1999; accepted 20 December 1999

Abstract

A number of symmetrically 3,5-disubstituted 4-amino-1,2,4-triazoles are quickly prepared by the reaction of aromatic nitriles on hydrazine dihydrochloride in the presence of an excess of hydrazine hydrate in ethylene glycol under microwave irradiation. © 2000 Elsevier Science Ltd. All rights reserved.

Keywords: 4-amino-1,2,4-triazoles; microwave irradiation.

4-Amino-1,2,4-triazoles **3** are potentially good corrosion inhibitors.^{1,2} These compounds can be prepared by the nucleophilic attack of hydrazine derivatives on nitriles.^{3–9} The initial product, in each case, is the 1,2-dihydro-1,2,4,5-tetrazine **2** which rearranges at elevated temperatures or on treatment with acid to the corresponding 4-amino-1,2,4-triazole **3**. One of these methods⁸ involves the reaction between 2-cyanopyridine **1h** and hydrazine hydrate. The reaction product was 3,6-bis(2-pyridyl)-1,2-dihydro-1,2,4,5-tetrazine **2h**. This compound was rearranged by further treatment with hydrochloric acid to 3,5-bis(2-pyridyl)-4-amino-1,2,4-triazole **3h**. Recently, a one-step synthesis of 3,5-disubstituted 4-amino-1,2,4-triazoles by direct reaction of hydrazine derivatives on nitriles bearing a variety of substituted aromatic rings or heteroaromatic has been reported.¹⁰



* Corresponding author. Fax: (33) 03 20 43 49 74; e-mail: didier.barbry@univ-lille1.fr (D. Barbry)

Table 1
Data for the synthesis of 3,5-diaryl-4-amino-1,2,4-triazoles **3a–i** under microwave irradiation

Compound	Ar	Microwave Irradiation		Classical Heating		Mp (°C)
		Time (min)	Yield (%)	Time (min)	Yield (%)	
3a	C ₆ H ₅	4	92	60	97	263
3b	4-CH ₃ C ₆ H ₄	6	66	60	67	294
3c	4-H ₂ NC ₆ H ₄	4	70	45	67	226
3d	4-HOC ₆ H ₄	10	96	45	87	295
3e	3-CH ₃ O, 4-OHC ₆ H ₃	6	79	60	86	228
3f	4-CH ₃ OC ₆ H ₄	6	73	45	83	277
3g	4-ClC ₆ H ₄	5	63	45	61	290
3h	2-pyridyl	5	95	45	85	186
3i	4-pyridyl	6	58	45	75	340

This method allows the synthesis of the solely symmetrically 3,5-disubstituted 4-amino-1,2,4-triazoles. We report that this reaction takes place in good yields and rapidly under microwave irradiation (Table 1). This technique has been applied with success to a number of reactions proceeding with or without solvent.^{11,12}




Fig. 1. Yield versus time for 3,5-diphenyl-4-amino-triazole at 130°C

The initial product, in each synthesis of aminotriazoles, is the orange-coloured, dihydro-1,2,4,5-tetrazine **2** which rearranges on treatment under acidic conditions into the corresponding 4-amino-1,2,4-triazole **3**. Addition of hydrazine dihydrochloride or sulfate was used to generate protons which promoted the rearrangement of **2** into the triazole **3**. Without using hydrazine dihydrochloride, the main reaction product was **2**. For example, using the same conditions, a mixture of benzonitrile **1a** and hydrazine hydrate, heated in ethylene glycol, leads to a mixture of **2** and **3** with about 70% of **2** (estimated by integration of the nitrogen protons in ¹H NMR). Compound **2** can be differentiated from **3** by oxidation of the former to the red-coloured tetrazine. However, a further reliable basis of differentiation is their ¹H

NMR spectra. The nitrogen protons of dihydrotetrazine appeared at a much lower field ($\delta=9.1$ ppm) than did those of 4-amino-1,2,4-triazoles ($\delta=6.4$ ppm).

Under classical heating in a high boiling polar solvent such as ethylene or diethylene glycol obtention of compounds **3** in good yield required much longer reaction times (45–60 min) at 130°C depending on starting compounds **1** of the reaction.¹⁰ With microwave irradiation some reactions were complete after just 4 min (compounds **3a** and **3c**) and did not exceed 10 min (compound **3d**). The rates of formation of 3,5-diphenyl-4-aminotriazole at 130°C with classical heating and under microwave irradiation have been compared (Fig. 1). Superheating of the solvent under microwave irradiation may lead to the observed rate enhancement.¹⁴

References

1. Memari, B.; Elattari, H.; Traisnel, M.; Bentiss, F.; Lagrenée, M. *Corros. Sci.* **1998**, *40*, 391–399.
2. Bentiss, F.; Lagrenée, M.; Traisnel, M.; Hornez, J. C. *Corros. Sci.* **1999**, *41*, 789–803.
3. Muller, E.; Herrdegen, L. *J. Prakt. Chem.* **1921**, *102*, 113–155.
4. Dallacker, F. *Monatsh. Chem.* **1960**, *91*, 294–304.
5. Zajack, W. W.; Siuda, J. F.; Nolan, M. J.; Santossusso, T. M. *J. Org. Chem.* **1971**, *36*, 3539–3541.
6. Abdel-Rahman, M. O.; Kira, M. A.; Tolba, M. N. *Tetrahedron Lett.* **1968**, 3871–3872.
7. Bowie, R. A.; Gardner, M. D.; Neilson, D. G.; Watson, K. M.; Mahmood, S.; Ridd, V. J. *Chem. Soc., Perkin Trans. 1* **1972**, 2395–2399.
8. Geldard, J. F.; Lions, F. *J. Org. Chem.* **1965**, *30*, 318–320.
9. Matsukawa, K.; Setsune, J.; Takagi, K.; Kitao, T. *Dyes and Pigment* **1982**, *3*, 307–315.
10. Bentiss, F.; Lagrenée, M.; Traisnel, M.; Mernari, B.; Elattari, H. *J. Heterocyclic Chem.* **1999**, *36*, 149–152.
11. Caddick, S. *Tetrahedron* **1995**, *51*, 10403–10432.
12. Loupy, A.; Petit, A.; Hamelin, J.; Texier-Boullet, F.; Jacquault, P.; Mathe, D. *Synthesis* **1998**, 1213–1234.
13. *Typical procedure*: A mixture of the corresponding aromatic nitrile (0.01 mol), hydrazine dihydrochloride (0.01 mol) and hydrazine hydrate (0.03 mol) in ethylene glycol (5 ml) was introduced into a quartz cylindrical flask placed in a Synthwave 402 monomode microwave oven under a reflux condenser and irradiated for 4–10 min (60 W) at 130°C. After cooling, the reaction mixture was diluted with water (10 ml). The precipitate thus obtained was filtered, washed with water, dried and recrystallized from ethanol. Products were identified by ¹H and ¹³C NMR, MS and IR spectroscopy: data were in agreement with literature values.
14. Baghurst, D.; Mingos, D. *J. Chem. Soc., Chem. Commun.* **1992**, 674–677.