

## Synthetic Approach Toward the Partial Sequences of Betaglycan in the Linkage Region on Solid Support and in Solution Phase

Jun-ichi Tamura , Akihiro Yamaguchi , Junko Tanaka & Yuko Nishimura

To cite this article: Jun-ichi Tamura , Akihiro Yamaguchi , Junko Tanaka & Yuko Nishimura (2007) Synthetic Approach Toward the Partial Sequences of Betaglycan in the Linkage Region on Solid Support and in Solution Phase, Journal of Carbohydrate Chemistry, 26:2, 61-82, DOI: [10.1080/07328300701296810](https://doi.org/10.1080/07328300701296810)

To link to this article: <http://dx.doi.org/10.1080/07328300701296810>



Published online: 09 May 2007.



Submit your article to this journal [↗](#)



Article views: 59



View related articles [↗](#)



Citing articles: 4 View citing articles [↗](#)

# Synthetic Approach Toward the Partial Sequences of Betaglycan in the Linkage Region on Solid Support and in Solution Phase

Jun-ichi Tamura, Akihiro Yamaguchi, Junko Tanaka, and  
Yuko Nishimura

Department of Environmental Sciences, Faculty of Regional Sciences, Tottori University, Tottori, Japan

We have synthesized, for the first time, the partial sequence of the betaglycan composed of the tetraosyl hexapeptide, which was directly usable as a probe for enzymatic glycosyl transfer. Stepwise elongation afforded the corresponding tetraosyl trichloroacetimidate. The common glycosyl dipeptide:[ $\beta$ -D-GlcA-(1  $\rightarrow$  3)- $\beta$ -D-Gal-(1  $\rightarrow$  3)- $\beta$ -D-Gal-(1  $\rightarrow$  4)- $\beta$ -D-Xyl-(1  $\rightarrow$  O)-Ser-Gly] was synthesized by glycosylation of the corresponding tetraosyl trichloroacetimidate and Ser-Gly moiety. The glycosyl dipeptide was coupled with other core peptide parts in solution phase and on a solid support. These glycosyl hexapeptides were then transformed into the desired target compounds.

**Keywords** Proteoglycan, Glycosaminoglycan, Betaglycan, Heparin, Heparan sulfate

## INTRODUCTION

Glycosaminoglycan (GAG) is the saccharide part of proteoglycans (PGs) covalently linked to the core peptide through the hydroxyl group of the serine residue. GAG is composed of two parts: the linkage (tetrasaccharide) region and repeating disaccharide region. Biochemically, GAG elongates by a stepwise addition with the help of UDP-sugars and the corresponding glycosyl transferases.<sup>[1]</sup> GAG is classified into two categories based on the


Received 16 November 2006; accepted 19 February 2007.

Address correspondence to Jun-ichi Tamura, Department of Environmental Sciences, Faculty of Regional Sciences, Tottori University, Tottori 680-8551, Japan. E-mail: jtamura@rstu.jp


type of the hexosamine residues in the repeating disaccharide region (Fig. 1). The first transfer of the hexosamine residue,  $\alpha$ -GlcNAc or  $\beta$ -GalNAc, which is the fifth saccharide moiety from the reducing terminal, determines the type of GAG as either the heparin or chondroitin type. Most of the glycosyl transferases involved in the GAG biosynthesis have been found; however, the factors controlling the biosynthetic sorting mechanisms are still unclear.

To the best of our knowledge, two hypotheses on the sorting mechanisms, which are not consistent with each other, have been supported on the basis of the environment around the glycan and/or peptide part of PG. The common tetrasaccharide often possesses sulfates at specific positions. Sugahara and his coworkers advocated that no sulfate has been detected on the heparin type of GAG vs. that on the chondroitin type.<sup>[2]</sup> It seems that the sulfate in the linkage region of chondroitin might provide an important function for the  $\beta$ -GalNAc transferase. On the other hand, Esko's group reported that the hydrophobic and acidic amino acid clusters of the core peptide are often flanking the heparin type of GAG.<sup>[3]</sup> His group demonstrated the glycan elongation toward the heparin type of GAG using mutant peptides and glycosyl primers containing hydrophobic aglycons.<sup>[4]</sup> These results provided important evidence for the relationship between the selective biosynthesis of heparin/heparan sulfate and the specific character of the common tetrasaccharide and/or the core peptides; however, the transfer mechanism of the first hexosamine residue remains unclear.

Heparin/heparan sulfates are medically and pharmaceutically indispensable materials; therefore, the selective and effective biosynthesis of heparin/heparan sulfate is anticipated. These facts prompted us to elucidate the biosynthetic mechanisms of heparin/heparan sulfate from a chemical point of view. Figure 2 shows the partial structure of betaglycan, of which the core peptide contains acidic and hydrophobic amino acids. It is noteworthy that the Ser<sup>535</sup> of the betaglycan predominantly possesses the heparin type of GAG.<sup>[5]</sup> We selected the target compounds (**1** and **2**), which are partial sequences of betaglycan containing the above core peptide. We now report



**Figure 1:** Structure of glycosaminoglycans.




**Figure 2:** Target compounds. A partial sequence of betaglycan.

the detailed synthesis of these betaglycans, which can be the acceptors for the  $\alpha$ -GlcNAc transfer.

## RESULTS AND DISCUSSION

Based on a retrosynthetic analysis, we planned (1) the stepwise elongation by the coupling of each monosaccharide moiety from the reducing terminal, (2) the subsequent coupling of the obtained tetrasaccharide with the Ser-Gly moiety, (3) the coupling of the tetraosyl serylglycine with the tetrapeptide synthesized in the solution phase or on a solid support, and (4) the final deprotection to obtain the desired compounds.

As depicted in Scheme 1, in the presence of 4-methoxyphenol and TMSOTf, known tetraacetyl- $\beta$ -D-xylose (**3**)<sup>[6]</sup> was converted into the corresponding 4-methoxyphenyl  $\beta$ -D-xyloside (**4**) in a 90% yield. The acetyl groups of **4** were removed with Et<sub>3</sub>N to quantitatively give **5**. The hydroxyl groups of **5** at the 2 and 3 positions were isopropylidened to afford **6** in a 75% yield. The galactosyl imidate (**7**) was converted from the known 4-methoxyphenyl 2,4,6-tri-O-acetyl-3-O-allyl- $\beta$ -D-galactopyranoside<sup>[7]</sup> by the removal of the 4-methoxyphenyl group with CAN, then followed by trichloroacetimidation with CCl<sub>3</sub>CN and DBU in 79% and 71% yields, respectively. The 4-OH of **6** was glycosylated with **7** in the presence of TMSOTf in CH<sub>2</sub>Cl<sub>2</sub>. The following methanolysis afforded the disaccharide (**8**) in an 82% yield in two steps. The diol **8** was acylated with 4-methylbenzoyl chloride to produce **9** in a 77% yield. The position of the newly formed glycosidic linkage was evident from the acylation shifts of the H-2 and H-3 of the xylosyl residue. The allyl group of **9** was isomerized using an iridium catalyst, and the I<sub>2</sub> treatment gave the disaccharide acceptor **10** in a 95% yield in two steps. The disaccharide acceptor **10** was stereoselectively coupled with **7** in the presence of TMSOTf in CH<sub>2</sub>Cl<sub>2</sub> to afford the trisaccharide **11** in an 86% yield. The allyl group of **11** was removed in the same manner as above to give the trisaccharide acceptor **12** in an 85% yield in two steps.



**Scheme 1:** Reagents and conditions: i) 4-methoxyphenol, TMSOTf, MS4A,  $\text{CH}_2\text{Cl}_2$ ,  $-20^\circ\text{C}$ , 1 h; ii)  $\text{Et}_3\text{N}$ , MeOH,  $\text{H}_2\text{O}$ , rt, overnight; iii) 2-methoxypropene, camphorsulfonic acid, DMF,  $60^\circ\text{C}$ , 5 h; iv) TMSOTf, MSAW300,  $\text{CH}_2\text{Cl}_2$ ,  $-20^\circ\text{C}$ , 1 h; v) camphorsulfonic acid, MeOH,  $\text{CH}_2\text{Cl}_2$ , rt, 2 h; vi) 4-methylbenzoyl chloride, pyridine, rt, 1.5 h; vii) (1,5-cyclooctadiene)bis(methyldiphenylphosphine)iridium(I) hexafluorophosphate, THF,  $\text{H}_2$ , then  $\text{I}_2$ ,  $\text{H}_2\text{O}$ ,  $\text{NaHCO}_3$ , rt, 2.5 h; viii)  $\text{BF}_3 \cdot \text{OEt}_2$ , MSAW300,  $\text{CH}_2\text{Cl}_2$ ,  $-20^\circ\text{C} \sim \text{rt}$ , overnight; ix)  $\text{CuBr}_2$ , AgOTf, *n*- $\text{Bu}_4\text{NBr}$ , MS4A,  $\text{CH}_2\text{Cl}_2$ ,  $0^\circ\text{C} \sim \text{rt}$ , overnight; x) CAN,  $\text{CH}_3\text{CN}$ ,  $\text{H}_2\text{O}$ ,  $0^\circ\text{C}$ , 4 h; xi)  $\text{CCl}_3\text{CN}$ , DBU,  $\text{CH}_2\text{Cl}_2$ ,  $0^\circ\text{C}$ , 2 h; xii)  $(\text{Ph}_3\text{P})_4\text{Pd}(\text{O})$ , *N*-methylaniline, THF, rt, 4 h.


For the following coupling with the glucuronic acid moiety, we prepared two types of glycosyl donors: methyl thioglycoside **13**<sup>[8]</sup> and trichloroacetimidate **14**.<sup>[9]</sup> Although **13** was used to couple with **12** mediated by the NIS-TfOH system, no tetrasaccharide was obtained. However, AgOTf- $\text{CuBr}_2$ -*n*- $\text{Bu}_4\text{NBr}$  could afford the desired compound **15** in a 31% yield. The glycosylation of **12** with **14** using  $\text{BF}_3 \cdot \text{OEt}_2$  also afforded **15** in a 22% yield with 69% of the

acceptor being recovered. When TMSOTf was used as the promoter, the reaction afforded **15** in a 32% yield, while the residual acceptor was completely decomposed.

The obtained tetrasaccharide was converted into the corresponding hemiacetal **16** with CAN in a 93% yield, and brought to **17** using  $\text{CCl}_3\text{CN}$  in the presence of DBU in an 89% yield. The imide (**17**) was coupled with known seryl-glycine allyl esters (**18**<sup>[10]</sup> and **19**<sup>[10]</sup>) protected with Boc and Fmoc groups at their N-terminals, with the help of TMSOTf in  $\text{CH}_2\text{Cl}_2$  at  $-20^\circ\text{C}$  to give **20** and **21** in 72% and 74% yields, respectively. The allyl groups were removed with  $\text{Pd}(\text{PPh}_3)_4$  and *N*-methylmorpholine to yield the corresponding carboxylic acids **22** and **23** in 90% and 76% yields, respectively.

## Synthesis of the Peptide Part


As shown in Scheme 2, the tetrapeptide Trp-Pro-Asp-Gly was synthesized in the conventional manner (liquid phase). Commercially available Fmoc-proline was converted into the corresponding allyl ester (**24**) in an 86% yield. The Fmoc group of **24** was removed with morpholine, and coupled with Fmoc-(*N*<sup>t</sup>-Boc)-tryptophane using HOBt and HBTU in the presence of Hünig's base to give **25** in a 56% yield. The allyl ester was converted into the carboxylic acid **26** with  $\text{Pd}(\text{PPh}_3)_4$  and *N*-methylmorpholine in a 63% yield. On the other hand, the *Z*-(*O*-*t*-Bu)-aspartic acid and the glycine *tert*-butyl ester, both of which were commercially available, were coupled with WSCD · HCl and HOBt in the presence of  $\text{Et}_3\text{N}$  to yield **27** in a 96% yield.




**Scheme 2:** Reagents and conditions: i)  $\text{Cs}_2\text{CO}_3$ , EtOH, rt, 1 h, then  $\text{AlIBr}$ , DMF, rt, overnight; ii) Morpholine,  $\text{CH}_2\text{Cl}_2$ , rt, overnight; iii) HOBt, HBTU,  $\text{CH}_2\text{Cl}_2$ ,  $-20^\circ\text{C} \sim \text{rt}$ , overnight; iv)  $(\text{Ph}_3\text{P})_4\text{Pd}(\text{O})$ , *N*-methylaniline, THF, rt, 15 h; v)  $\text{Pd-C}$ ,  $\text{H}_2$ , MeOH, AcOH, rt, 3 h.

Hydrogenolysis of **27** was successfully performed to give the free amine **28** in a 93% yield. The following condensation of **26** and **28** were executed in the same manner as the synthesis of **27** to furnish the tetrapeptide **29** in an 85% yield. The Fmoc group of **29** was finally removed with morpholine to give the free amine **30**, which was coupled with **22** in the presence of HOBt and HBTU. As depicted in Scheme 3, the successful condensation afforded the tetraosyl hexapeptide **31** in a 63% yield. The following deprotection procedures—(1) treatment with a TFA cocktail and (2) saponification with M0.107 NaOMe in 50% MeOH—gave the desired compound **1** in an 83% yield after gel permeation chromatography.

Scheme 4 shows the synthesis of the corresponding glycopeptide on Sieber amide resin. The elongation was manually performed by the usual Fmoc procedure employing HOBt, HBTU, and *i*-Pr<sub>2</sub>EtN in NMP. The Fmoc group was removed using piperidine in NMP. The *N*-terminal of the tetrapeptide on the resin was coupled with **23** in the same manner as the peptide elongation. The obtained glycosylated peptide on the resin was subjected to the TFA cocktail to give the corresponding tetraosyl hexapeptide. Careful saponification with diluted base (1M NaOH-MeOH 1:100, and the following 5 mM NaOH)<sup>[11]</sup> treatment afforded the fully deprotected glycosyl peptide **2** in the amide



**Scheme 3:** Reagents and conditions: i) HOBt, HBTU, *i*-Pr<sub>2</sub>EtN, DMF, −20°C ~rt, overnight; ii) TFA, thioanisole, phenol, H<sub>2</sub>O, 1,2-ethanedithiol, trisopropylsilane, CH<sub>2</sub>Cl<sub>2</sub>, rt, 2 h; iii) NaOMe, MeOH, H<sub>2</sub>O, rt, 3 h.



**Scheme 4:** Reagents and conditions: i) piperidine (20%), NMP, ii) HOBT, HBTU,  $i$ -Pr<sub>2</sub>EtN, iii) TFA, thioanisole, phenol, H<sub>2</sub>O, 1,2-ethanedithiol, triisopropylsilane, CH<sub>2</sub>Cl<sub>2</sub>, rt, 3 h; iv) NaOMe, MeOH, H<sub>2</sub>O, rt.

form. Further purification was successfully performed by gel permeation chromatography and HPLC (C18). The final compounds **1** and **2** were in good agreement with the calculated value based on the mass spectrum.

It was revealed that compound **1** was a poor substrate for the chondroitin GalNAc T-1 and T-2<sup>[12]</sup> but a superior substrate for heparan sulfate elongation (EXT1/EXT2) as well as the disaccharide primer,  $\beta$ -D-GlcA(1  $\rightarrow$  3)  $\beta$ -D-Gal(1  $\rightarrow$  O)C<sub>2</sub>H<sub>4</sub>NHZ having a hydrophobic aglycon.<sup>[13]</sup>

In summary, we synthesized the partial sequences in the linkage region of betaglycan as primers for the biosynthesis of heparan sulfate. Both synthetic strategies in the liquid phase as well as on a solid support are available for the synthesis of the target tetraosyl hexapeptides:  $\beta$ GlcA- $\beta$ Gal- $\beta$ Gal- $\beta$ Xyl-SerGlyTrpProAspGly.

## EXPERIMENTAL

### General Methods

Optical rotations were measured at  $22 \pm 3^\circ\text{C}$  with a HORIBA polarimeter SEPA-200. <sup>1</sup>H NMR assignments were confirmed by two-dimensional HH-COSY experiments with a JEOL ECP 500 MHz spectrometer. Signal



assignments such as  $1^{\text{III}}$  stand for a proton at C-1 of sugar residue III. MALDI-TOF and ESI mass spectra were measured by Daltonics Autoflex II spectrometer (Bruker) and Q-TOF2 (Micromass) instrument, respectively. Silica gel chromatography, analytical TLC, and preparative TLC (PTLC) were done on a column of Silica Gel 60 (Merck), Silica Gel 60 N (spherical neutral), (Kanto Kagaku) or glass plates coated with Silica Gel F<sub>254</sub> (Merck), respectively. Gels for size-exclusion chromatography (Sephadex LH-20 and Biobeads S-X1) were purchased from Amersham Biosciences and BIO · RAD, respectively. Reversed phase short column (Bond Elut<sup>®</sup> C8) was purchased from Varian Inc. Molecular sieves (MS) were purchased from GL Science, Inc., and activated at 200°C under diminished pressure prior to use. All reactions in organic solvents were performed under dry Ar atmosphere. As a usual workup, the organic phase of the reaction mixture was washed with aq. NaHCO<sub>3</sub> and brine and dried over MgSO<sub>4</sub>.

#### 4-Methoxyphenyl 2,3,4-tri-O-acetyl- $\beta$ -D-xylopyranoside (4)

To a suspension of 1,2,3,4-tetra-O-acetyl- $\beta$ -D-xylopyranose<sup>[6]</sup> (**3**, 15.5 g, 48.7 mmol) and 4-methoxyphenol (6.70 g, 54.0 mmol) in the presence of activated MS 4A (2.5 g) in 1,2-dichloroethane (100 mL) was added TMSOTf (4.1 mL, 2.3 mmol) at  $-20^\circ\text{C}$  with stirring, and then the reaction temperature was gradually raised up to  $-14^\circ\text{C}$  within 1 h. Et<sub>3</sub>N (15 mL) and an excess amount of saturated NaHCO<sub>3</sub> were added. Insoluble materials were filtered on Celite. The organic phase was treated as described in the general methods. The crude materials were subjected to a column of silica gel (5:1–3:1 toluene:EtOAc) to give **4** (16.8 g) in 90% yield. A small amount of the product was recrystallized from *n*-hexane:EtOAc to give white powder:  $[\alpha]_{\text{D}} -41.5$  (*c* 0.66, CHCl<sub>3</sub>); m.p.  $150.4^\circ\text{C}$ ; <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  6.96–6.94 (m, 2H, Ar H), 6.84–6.81 (m, 2H, Ar H), 5.23 (dd, 1H,  $J_{2,3} = 8.29$  Hz,  $J_{3,4} = 7.81$  Hz, H-3), 5.16 (dd, 1H,  $J_{1,2} = 6.10$  Hz, H-2), 5.03 (d, 1H, H-1), 5.01 (m, 1H, H-4), 4.21 (dd, 1H,  $J_{4,5\text{eq}} = 4.88$  Hz,  $J_{\text{gem}} = 11.96$  Hz, H-5eq), 3.77 (s, 3H, OMe), 3.47 (dd, 1H,  $J_{4,5\text{ax}} = 8.29$  Hz, H-5ax), 2.10, 2.08, 2.08 (3s, 3H  $\times$  3, 3MeCO). Anal. Calcd for C<sub>18</sub>H<sub>22</sub>O<sub>9</sub>: C, 56.53; H, 5.81. Found: C, 56.51; H, 5.81.

#### 4-Methoxyphenyl $\beta$ -D-xylopyranoside (5)

Triacetate (**4**) (16.8 g, 44.0 mmol) was diluted with Et<sub>3</sub>N (75 mL), MeOH (150 mL), and H<sub>2</sub>O (75 mL) with stirring overnight. The volatiles were removed under diminished pressure and the residue was coevaporated with water and toluene. The residue was purified from a column of silica gel (1:1–1:5 *n*-hexane:EtOAc–100:1–40:1–20:1 EtOAc:MeOH) to give **5** quantitatively:  $[\alpha]_{\text{D}} -26.7$  (*c* 1.29, MeOH); <sup>1</sup>H NMR (CD<sub>3</sub>OD):  $\delta$  6.91–6.89 (m, 2H, Ar H), 6.74–6.72 (m, 2H, Ar H), 4.61 (d, 1H,  $J_{1,2} = 7.56$  Hz, H-1), 3.79 (dd, 1H,  $J_{4,5\text{a}} = 5.37$  Hz,  $J_{\text{gem}} = 11.47$  Hz, H-5a), 3.64 (s, 3H, OMe), 3.5–3.2 (m, 4H, H-2,3,4,5e). Anal. Calcd for C<sub>12</sub>H<sub>16</sub>O<sub>6</sub>: C, 56.24; H, 6.31. Found: C, 56.19; H, 6.31.

**4-Methoxyphenyl 2,3-O-isopropylidene  $\beta$ -D-xylopyranoside (6)**

To a solution of **5** (15.2 g, 59.4 mmol) in DMF (70 mL) were added a camphor sulfonic acid (2.1 g, 9.0 mmol) and 2-methoxypropene (12.0 mL, 125 mmol) with stirring at 60°C. After 2 h, 6 mL of 2-methoxypropene was added for 30 minutes. Forty minutes later, the reaction mixture was cooled to rt and Et<sub>3</sub>N (6.0 mL, 35 mmol) was added. The reaction mixture was extracted with EtOAc. The organic phase was treated as described in the general methods. The residue was eluted from a column of silica gel (5:1–4:1–1:1–1:5 toluene:EtOAc) to give **6** (13.2 g, 75%) as a syrup:  $[\alpha]_D -32.3$  (c 1.32, CHCl<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  7.04–7.01 (m, 2H, Ar H), 6.84–6.81 (m, 2H, Ar H), 5.18 (d, 1H,  $J_{1,2} = 5.86$  Hz, H-1), 4.15 (m, 1H, H-5eq), 4.11 (m, 1H, H-4), 3.77 (s, 3H, OMe), 3.62 (m, 2H, H-2,3), 3.38 (dd, 1H,  $J_{4,5ax} = 6.34$  Hz,  $J_{gem} = 11.47$  Hz, H-5ax), 2.49 (d, 1H,  $J_{4,OH} = 3.91$  Hz, OH-4), 1.51, 1.49 (2s, 3H  $\times$  2, 2CH<sub>3</sub>). Anal. Calcd for C<sub>15</sub>H<sub>20</sub>O<sub>6</sub>  $\cdot$  0.2H<sub>2</sub>O: C, 60.06; H, 6.87. Found: C, 59.88; H, 6.90.

**2,4,6-Tri-O-acetyl-3-O-allyl- $\alpha$ -D-galactopyranosyl trichloroacetimidate (7)**

To a solution of 4-methoxyphenyl 2,4,6-tri-O-acetyl-3-O-allyl- $\beta$ -D-galactopyranoside<sup>[7]</sup> (77.3 g, 171 mmol) in CH<sub>3</sub>CN (400 mL) and H<sub>2</sub>O (100 mL) was added cerium (IV) ammonium nitrate (CAN) (88.4 g, 161 mmol) at 0°C with stirring. After 3 h, additional CAN (175.57 g, 320.3 mmol) was supplied and the reaction mixture was stirred for another 2.5 h, then diluted with CHCl<sub>3</sub> and brine and extracted with CHCl<sub>3</sub>. The organic phase was washed with brine and the residue was eluted from a column of silica gel (2:1–3:2–1:1–1:3 *n*-hexane:EtOAc) to give the corresponding hemiacetal (47.0 g, 79%) as a syrup. The hemiacetal was diluted with CH<sub>2</sub>Cl<sub>2</sub> (350 mL). To the solution were added CCl<sub>3</sub>CN (22 mL) and DBU (8.0 mL, 54 mmol) at 0°C. The reaction mixture was stirred for 1.5 h and directly subjected to a column of silica gel (3:1~3:2~1:1 *n*-hexane:EtOAc) to give **7** (47.4 g, 71%) as a syrup, which was used for the next reaction without further purification: <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  8.63 (s, 1H, NH), 6.56 (s, 1H, H-1), 5.88–5.78 (m, 1H, CH=), 5.58 (d, 1H,  $J_{3,4} = 2.54$  Hz, H-4), 5.31–5.17 (m, 3H, H-2, =CH<sub>2</sub>), 4.35 (brt, 1H,  $J = 6.47$  Hz, H-5), 4.23–3.97 (m, 5H, H-3, 6, CH<sub>2</sub>O), 2.16, 2.05, 2.04 (3s, 3H  $\times$  3, 3MeCO).

**4-Methoxyphenyl O-(2,4,6-tri-O-acetyl-3-O-allyl- $\beta$ -D-galactopyranosyl)-(1  $\rightarrow$  4)- $\beta$ -D-xylopyranoside (8)**

To a solution of **6** (321.0 mg, 1.083 mmol) and **7** (977.5 mg, 1.992 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (15 mL) was added MS AW300 (300 mg). This mixture was stirred for 30 min at rt and cooled to –15°C. To this solution was added a TMSOTf (70  $\mu$ L, 0.38 mmol) with stirring and the reaction temperature was gradually raised to –10°C for 1 h. Then, Et<sub>3</sub>N (110  $\mu$ L, 0.79 mmol) and an excess

amount of aqueous  $\text{NaHCO}_3$  were added. Insoluble materials were filtered on Celite. The organic phase was treated as described in the general methods. To a solution of the crude products in MeOH and  $\text{CH}_2\text{Cl}_2$  (1:1, 46 mL) was added a camphor sulfonic acid (251.6 mg, 1.08 mmol) and the mixture stirred for 2 h and quenched with  $\text{Et}_3\text{N}$  (150  $\mu\text{L}$ , 1.08 mmol). The volatiles were removed under diminished pressure and the residue was subjected to a column of silica gel (4:1–2:1–1:1–2:3 toluene:EtOAc) to give **8** (518.1 mg) in 82% yield as a syrup:  $[\alpha]_{\text{D}} -3.9$  (*c* 1.50,  $\text{CHCl}_3$ );  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  7.01–6.97 (m, 2H, Ar H), 6.85–6.82 (m, 2H, Ar H), 5.82–5.73 (m, 1H, CH=), 5.42 (d, 1H,  $J_{3,4} = 3.42$  Hz, H-4<sup>II</sup>), 5.34–5.17 (m, 2H, =CH<sub>2</sub>), 5.10 (dd, 1H,  $J_{1,2} = 8.05$  Hz,  $J_{2,3} = 10.00$  Hz, H-2<sup>II</sup>), 4.90 (d, 1H,  $J_{1,2} = 6.59$  Hz, H-1<sup>I</sup>), 4.48 (d, 1H, H-1<sup>II</sup>), 4.24 (dd, 1H,  $J_{5,6a} = 4.88$  Hz,  $J_{\text{gem}} = 11.47$  Hz, H-6a<sup>II</sup>), 4.14–4.07 (m, 1H, one of OCH<sub>2</sub>), 4.09 (m, 1H, H-6b<sup>II</sup>), 3.95 (dd, 1H,  $J_{4,5\text{eq}} = 4.64$  Hz,  $J_{\text{gem}} = 11.95$  Hz, H-5eq<sup>I</sup>), 3.92–3.88 (m, 2H, H-5<sup>II</sup>, one of CH<sub>2</sub>O), 3.89 (m, 1H, OH-3<sup>I</sup>), 3.78 (s, 3H, OMe), 3.78–3.73 (m, 1H, H-3<sup>I</sup>), 3.67 (m, 2H, H-2<sup>I</sup>, 4<sup>I</sup>), 3.53 (dd, 1H, H-3<sup>II</sup>), 3.41 (dd, 1H,  $J_{4,5\text{ax}} = 8.54$  Hz, H-5ax<sup>I</sup>), 2.78 (d, 1H,  $J_{2,\text{OH}} = 3.66$  Hz, OH-2<sup>I</sup>), 2.14, 2.12, 2.12, (3s, 3H  $\times$  3, 3MeCO). Anal. Calcd for  $\text{C}_{27}\text{H}_{36}\text{O}_{14} \cdot 0.5 \text{H}_2\text{O}$ : C, 54.62; H, 6.30. Found: C, 54.68; H, 6.02.

**4-Methoxyphenyl O-(2,4,6-tri-O-acetyl-3-O-allyl- $\beta$ -D-galactopyranosyl)-(1  $\rightarrow$  4)-2,3-di-O-(4-methylbenzoyl)- $\beta$ -D-xylopyranoside (9)**

To a solution of **8** (1.71 g, 2.92 mmol) in pyridine (10 mL) was added a 4-methylbenzoyl chloride (1.50 mL, 11.6 mmol) dropwise with stirring. After 1.5 h MeOH (0.5 mL) was added to the reaction mixture at 0°C and diluted with  $\text{CHCl}_3$ . The organic phase was treated as described in general methods. The crude materials were subjected to a column of silica gel (10:1–7:1–6:1–4:1–1:1 toluene:EtOAc) to give **9** (1.85 g, 77%) as a syrup:  $[\alpha]_{\text{D}} +33.9$  (*c* 0.945,  $\text{CHCl}_3$ );  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  7.97–7.95 (m, 4H, Ar H), 7.24–7.22 (m, 4H, Ar H), 6.98–6.96 (m, 2H, Ar H), 6.82–6.80 (m, 2H, Ar H), 5.84–5.71 (m, 1H, CH=), 5.68 (brt, 1H,  $J = 6.30$  Hz, H-3<sup>I</sup>), 5.43 (dd, 1H,  $J_{1,2} = 5.04$  Hz,  $J_{2,3} = 6.18$  Hz, H-2<sup>I</sup>), 5.29 (brd, 2H,  $J = 4.58$  Hz, H-1<sup>I</sup>, 4<sup>II</sup>), 5.23–5.14 (m, 2H, =CH<sub>2</sub>), 5.05 (dd, 1H,  $J_{1,2} = 8.02$  Hz,  $J_{2,3} = 10.08$  Hz, H-2<sup>II</sup>), 4.57 (d, 1H, H-1<sup>II</sup>), 4.23 (dd, 1H,  $J_{4,5\text{eq}} = 3.90$  Hz,  $J_{\text{gem}} = 12.37$  Hz, H-5eq<sup>I</sup>), 4.10–4.06 (m, 1H, one of OCH<sub>2</sub>), 3.98 (m, 1H, H-4<sup>I</sup>), 3.87–3.84 (m, 1H, one of CH<sub>2</sub>O), 3.83 (dd, 1H,  $J_{5,6a} = 6.19$  Hz,  $J_{\text{gem}} = 10.99$  Hz, H-6a<sup>II</sup>), 3.76 (s, 3H, OMe), 3.71 (brt, 1H,  $J = 6.53$  Hz, H-5<sup>II</sup>), 3.65 (dd, 1H,  $J_{4,5\text{ax}} = 6.65$  Hz, H-5ax<sup>I</sup>), 3.59 (dd, 1H,  $J_{5,6b} = 6.65$  Hz, H-6b<sup>II</sup>), 3.46 (dd, 1H,  $J_{3,4} = 3.44$  Hz, H-3<sup>II</sup>), 2.40, 2.38 (2s, 3H  $\times$  2, 2PhMe), 2.03, 2.02, 1.96 (3s, 3H  $\times$  3, 3MeCO). Anal. Calcd for  $\text{C}_{43}\text{H}_{48}\text{O}_{16} \cdot \text{H}_2\text{O}$ : C, 61.56; H, 6.02. Found: C, 61.72; H, 5.87.

**4-Methoxyphenyl O-(2,4,6-tri-O-acetyl- $\beta$ -D-galactopyranosyl)-(1  $\rightarrow$  4)-2,3-di-O-(4-methylbenzoyl)- $\beta$ -D-xylopyranoside (10)**

A suspension of a catalytic amount of (1,5-cyclooctadiene)bis(methyldiphenylphosphine)iridium(I) hexafluorophosphate in THF (10 mL) was stirred

under H<sub>2</sub> atmosphere, which was then replaced with argon. This manipulation was repeated a few times. Then, a solution of **9** (2.01 g, 2.45 mmol) in THF (20 mL) was added to the above solution of the iridium complex. After stirring for 2 h, H<sub>2</sub>O (5 mL), NaHCO<sub>3</sub> (3.92 g, 46.7 mmol), and I<sub>2</sub> (1.72 g, 6.78 mmol) were added to the reaction mixture, and the stirring was continued for 2.5 h at 0°C. The reaction mixture was then diluted with CHCl<sub>3</sub>. The organic phase was washed with aq. NaHCO<sub>3</sub>, aq. Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>, and brine. The volatiles were removed under diminished pressure, and the crude materials obtained were eluted from a column of silica gel (4:1–3:1–2:1–1:1–1:2 toluene:EtOAc) to give **10** (1.82 g, 95%) as a syrup:  $[\alpha]_D^{25} +22.6$  (*c* 1.54, CHCl<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 7.94 (m, 4H, Ar H), 7.29–7.21 (m, 4H, Ar H), 6.98–6.96 (m, 2H, Ar H), 6.82–6.80 (m, 2H, Ar H), 5.69 (t, 1H,  $J_{2,3} = J_{3,4} = 6.34$  Hz, H-3<sup>I</sup>), 5.43 (dd, 1H,  $J_{1,2} = 4.88$  Hz, H-2<sup>I</sup>), 5.30 (d, 1H, H-1<sup>I</sup>), 5.20 (d, 1H,  $J_{3,4} = 3.42$  Hz, H-4<sup>II</sup>), 4.90 (dd, 1H,  $J_{1,2} = 8.05$  Hz,  $J_{2,3} = 10.00$  Hz, H-2<sup>II</sup>), 4.60 (d, 1H, H-1<sup>II</sup>), 4.26 (dd, 1H,  $J_{4,5eq} = 3.66$  Hz,  $J_{gem} = 12.20$  Hz, H-5eq<sup>I</sup>), 4.00 (m, 1H, H-4<sup>I</sup>), 3.85 (dd, 1H,  $J_{5,6a} = 6.34$  Hz,  $J_{gem} = 10.98$  Hz, H-6a<sup>II</sup>), 3.80–3.73 (m, 2H, H-3<sup>II</sup>, 5<sup>II</sup>), 3.75 (s, 3H, OMe), 3.68 (dd, 1H,  $J_{4,5ax} = 6.10$  Hz, H-5ax<sup>I</sup>), 3.59 (dd, 1H,  $J_{5,6b} = 6.83$  Hz, H-6b<sup>II</sup>), 2.51, 2.49 (2s, 3H × 2, 2PhMe), 2.06, 2.03, 1.94 (3s, 3H × 3, 3MeCO). Anal. Calcd for C<sub>40</sub>H<sub>44</sub>O<sub>16</sub> · H<sub>2</sub>O: C, 60.14; H, 5.82. Found: C, 60.00; H, 5.53.

**4-Methoxyphenyl O-(2,4,6-tri-O-acetyl-3-O-allyl-β-D-galactopyranosyl)-(1 → 3)-O-(2,4,6-tri-O-acetyl-β-D-galactopyranosyl)-(1 → 4)-2,3-di-O-(4-methylbenzoyl)-β-D-xylopyranoside (11)**

To a solution of **10** (7.40 g, 9.48 mmol) and **7** (7.12 g, 14.5 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (160 mL) was added MS AW300 (4.0 g). This mixture was stirred for 35 min at rt, and then cooled to –20°C. To this solution was added a TMSOTf (660 μL, 3.65 mmol) with stirring. After 1 h, the reaction was quenched with Et<sub>3</sub>N (1.0 mL, 7.0 mmol) and excess amount of aq. NaHCO<sub>3</sub>. The reaction was treated as described in the synthesis of **8**. The residue was subjected to a column of silica gel (5:1–4:1–3:1–2:1–1:2 *n*-hexane:EtOAc) to give **11** (9.06 g) in 86% yield as a syrup:  $[\alpha]_D^{25} +51.5$  (*c* 0.96, CHCl<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 7.98–7.94 (m, 5H, Ar H), 7.24–7.23 (m, 3H, Ar H), 6.97 (m, 2H, Ar H), 6.82 (m, 2H, Ar H), 5.80–5.70 (m, 1H, =CH), 5.68 (t, 1H,  $J_{2,3} = J_{3,4} = 6.10$  Hz, H-3<sup>I</sup>), 5.42 (dd, 1H,  $J_{1,2} = 4.88$  Hz, H-2<sup>I</sup>), 5.37 (d, 1H,  $J_{3,4} = 3.42$  Hz, H-4<sup>III</sup>), 5.30 (d, 1H, H-1<sup>I</sup>), 5.27 (d, 1H,  $J_{3,4} = 3.42$  Hz, H-4<sup>II</sup>), 5.19 (m, 2H, =CH<sub>2</sub>), 5.16 (dd, 1H,  $J_{1,2} = 7.81$  Hz,  $J_{2,3} = 9.51$  Hz, H-2<sup>II</sup>), 4.95 (dd, 1H,  $J_{1,2} = 8.05$  Hz,  $J_{2,3} = 10.00$  Hz, H-2<sup>III</sup>), 4.55 (d, 1H, H-1<sup>II</sup>), 4.45 (d, 1H, H-1<sup>III</sup>), 4.23 (dd, 1H,  $J_{4,5eq} = 3.66$  Hz,  $J_{gem} = 12.44$  Hz, H-5eq<sup>I</sup>), 4.10 (m, 2H, OCH<sub>2</sub>), 4.10 (dd,  $J_{5,6b} = 5.12$  Hz,  $J_{gem} = 12.69$  Hz, H-6b<sup>III</sup>), 3.96 (m, 1H, H-4<sup>I</sup>), 3.89–3.85 (m, 2H, H-6a<sup>III</sup>, 6b<sup>II</sup>), 3.84–3.75 (m, 3H, H-3<sup>II</sup>, 5<sup>II</sup>, 5<sup>III</sup>), 3.64 (dd, 1H,  $J_{4,5ax} = 6.10$  Hz, H-5ax<sup>I</sup>), 3.51 (dd, 1H,  $J_{5,6a} = 6.83$  Hz, H-6a<sup>II</sup>), 3.41 (dd, 1H, H-3<sup>III</sup>), 2.40, 2.39

(2s, 3H × 2, 2MePh), 2.14, 2.08, 2.06, 2.06, 2.04, 1.94 (6s, 3H × 6, 6MeCO). ESI-MS (positive) Calcd for C<sub>55</sub>H<sub>64</sub>O<sub>24</sub>Na [M + Na]<sup>+</sup>, 1131.4. Found: 1130.9.

**4-Methoxyphenyl O-(2,4,6-tri-O-acetyl-β-D-galactopyranosyl)-(1 → 3)-O-(2,4,6-tri-O-acetyl-β-D-galactopyranosyl)-(1 → 4)-2,3-di-O-(4-methylbenzoyl)-β-D-xylopyranoside (12)**

The iridium catalyst was suspended in THF and activated as described in the synthesis of **10**. Then, a solution of **11** (95.9 mg, 86.5 μmol) in THF (3 mL) was added to the iridium complex solution. After stirring for 45 min, H<sub>2</sub>O (1 mL), NaHCO<sub>3</sub> (145.3 mg, 1.730 mmol), and I<sub>2</sub> (43.9 mg, 0.174 mmol) were added to the reaction mixture at 0°C. The solution was stirred for 45 min at 0°C and treated as described in the synthesis of **10**. The obtained crude material was eluted from a column of silica gel (9:1–7:1–5:1–3:1–2:1–1:5 toluene:EtOAc) to give **12** (78.5 mg, 85%) as a syrup: [α]<sub>D</sub> +26.3 (c 1.01, CHCl<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 7.97–7.94 (m, 5H, Ar H), 7.24–7.22 (m, 3H, Ar H), 6.97 (m, 2H, Ar H), 6.81 (m, 2H, Ar H), 5.68 (t, 1H, *J*<sub>2,3</sub> = *J*<sub>3,4</sub> = 6.10 Hz, H-3<sup>I</sup>), 5.43 (dd, 1H, *J*<sub>1,2</sub> = 4.89 Hz, H-2<sup>I</sup>), 5.30–5.27 (m, 3H, H-1<sup>I</sup>, 4<sup>II</sup>, 4<sup>III</sup>), 5.16 (dd, 1H, *J*<sub>1,2</sub> = 8.05 Hz, *J*<sub>2,3</sub> = 10.00 Hz, H-2<sup>II</sup>), 4.83 (dd, 1H, *J*<sub>1,2</sub> = 8.05 Hz, *J*<sub>2,3</sub> = 10.00 Hz, H-2<sup>III</sup>), 4.58 (d, 1H, H-1<sup>I</sup>), 4.47 (d, 1H, H-1<sup>III</sup>), 4.23 (dd, 1H, *J*<sub>4,5eq</sub> = 3.66 Hz, *J*<sub>gem</sub> = 12.44 Hz, H-5eq<sup>I</sup>), 4.17–4.08 (m, 2H, H-6<sup>III</sup>), 3.97 (m, 1H, H-4<sup>I</sup>), 3.86 (dd, 1H, *J*<sub>5,6b</sub> = 5.37 Hz, *J*<sub>gem</sub> = 11.22 Hz, H-6b<sup>II</sup>), 3.79–3.71 (m, 3H, H-3<sup>II</sup>, 3<sup>III</sup>, 5<sup>II</sup>), 3.64 (dd, 1H, *J*<sub>4,5ax</sub> = 6.34 Hz, H-5ax<sup>I</sup>), 3.51 (dd, 1H, *J*<sub>5,6a</sub> = 7.07 Hz, H-6a<sup>II</sup>), 2.41, 2.39 (2s, 3H × 2, 2MePh), 2.18, 2.12, 2.07, 2.04, 2.04, 1.94 (6s, 3H × 6, 6MeCO). TOF-MS (positive) Calcd for C<sub>52</sub>H<sub>60</sub>O<sub>24</sub>Na [M + Na]<sup>+</sup>, 1091.3. Found: 1091.5.

**4-Methoxyphenyl O-[methyl 2,3,4-tri-O-(4-methylbenzoyl)-β-D-glucopyranosyl uronate]-(1 → 3)-O-(2,4,6-tri-O-acetyl-β-D-galactopyranosyl)-(1 → 3)-O-(2,4,6-tri-O-acetyl-β-D-galactopyranosyl)-(1 → 4)-2,3-di-O-(4-methylbenzoyl)-β-D-xylopyranoside (15)**

**Method A:** A suspension containing CuBr<sub>2</sub> (1.37 g, 6.14 mmol), AgOTf (1.58 g, 6.14 mmol), *n*-Bu<sub>4</sub>NBr (327 mg, 1.01 mmol), and MS 4A (5.0 g) in CH<sub>2</sub>Cl<sub>2</sub> (31 mL) was stirred in the dark at rt and cooled to 0°C after 30 min. Then, CH<sub>2</sub>Cl<sub>2</sub> (60 mL) and a solution of **13** (798.3 mg, 1.347 mmol) and **12** (713.0 mg, 0.667 mmol) were added dropwise to the suspension and stirred overnight, gradually raised to rt. The reaction was quenched with aq. NaHCO<sub>3</sub> and brine. Then, the reaction mixture was diluted with EtOAc and filtered on Celite. The organic phase was treated as usual. The obtained crude materials were eluted from a column of silica gel (3:1–3:2–1:1–1:10 *n*-hexane:EtOAc) to give **15** (338.8 mg, 31%) as a syrup together with 288.6 mg (41%) of recovered **12**.

**Method B:** To a solution of **14** (1.11 g, 1.57 mmol) and **12** (675.1 mg, 0.631 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (28 mL) was added MS AW300 (300 mg). This

mixture was stirred for 10 min at rt, and then cooled to  $-20^{\circ}\text{C}$ . To this solution was added a TMSOTf (85  $\mu\text{L}$ , 0.47 mmol) with stirring, and the same amount of TMSOTf was added after 2 h. One hour later, the reaction was quenched with  $\text{Et}_3\text{N}$  (260  $\mu\text{L}$ , 1.88 mmol) and an excess amount of aq.  $\text{NaHCO}_3$  was added. The reaction was treated as described in the synthesis of **8**. The residue was purified as above to give **15** (323.7 mg) in 32% yield..

**Method C:** To a solution of **14** (147.9 mg, 209.2  $\mu\text{mol}$ ) and **12** (96.5 mg, 90.3  $\mu\text{mol}$ ) in toluene (17.5 mL) and  $\text{CH}_2\text{Cl}_2$  (0.2 mL) was added MS AW300 (480 mg). This mixture was stirred for 30 min at rt, and then cooled to  $-20^{\circ}\text{C}$ . To this solution was added a  $\text{BF}_3 \cdot \text{OEt}_2$  (8  $\mu\text{L}$ , 0.09 mmol) with stirring and the same amount of  $\text{BF}_3 \cdot \text{OEt}_2$  was added after 1.5 h. The reaction mixture was stirred overnight while the temperature gradually was raised to rt. The reaction was worked up with an excess amount of aq.  $\text{NaHCO}_3$  and treated as above to give **15** (31.4 mg) in 22% yield together with 66.4 mg (69%) of recovered **12**:  $[\alpha]_{\text{D}} +26.9$  (c 1.01,  $\text{CHCl}_3$ );  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  7.97–7.93 (m, 4H, Ar H), 7.81–7.70 (m, 6H, Ar H), 7.23–7.07 (m, 10H, Ar H), 6.96 (m, 2H, Ar H), 6.81 (m, 2H, Ar H), 5.80 (t, 1H,  $J_{2,3} = J_{3,4} = 9.27$  Hz, H-3<sup>IV</sup>), 5.65 (t, 1H,  $J_{2,3} = J_{3,4} = 6.10$  Hz, H-3<sup>I</sup>), 5.64 (brt, 1H,  $J = 9.76$  Hz, H-4<sup>IV</sup>), 5.47 (d, 1H,  $J_{3,4} = 3.45$  Hz, H-4<sup>III</sup>), 5.41 (dd, 1H,  $J_{1,2} = 4.88$  Hz, H-2<sup>I</sup>), 5.37 (dd, 1H,  $J_{1,2} = 7.56$  Hz, H-2<sup>IV</sup>), 5.29 (d, 1H, H-1<sup>I</sup>), 5.22 (d, 1H,  $J_{3,4} = 3.42$  Hz, H-4<sup>II</sup>), 5.08 (dd, 1H,  $J_{1,2} = 8.05$  Hz,  $J_{2,3} = 10.00$  Hz, H-2<sup>II</sup>), 5.02 (dd, 1H,  $J_{1,2} = 8.05$  Hz,  $J_{2,3} = 10.24$  Hz, H-2<sup>III</sup>), 4.91 (d, 1H, H-1<sup>IV</sup>), 4.51 (d, 1H, H-1<sup>II</sup>), 4.37 (d, 1H, H-1<sup>III</sup>), 4.26 (d, 1H,  $J_{4,5} = 10.01$  Hz, H-5<sup>IV</sup>), 4.21 (dd, 1H,  $J_{4,5\text{eq}} = 3.91$  Hz,  $J_{\text{gem}} = 12.19$  Hz, H-5eq<sup>I</sup>), 4.16 (dd, 1H,  $J_{5,6\text{b}} = 5.85$  Hz,  $J_{\text{gem}} = 11.71$  Hz, H-6b<sup>III</sup>), 4.03 (dd, 1H,  $J_{5,6\text{a}} = 6.34$  Hz, H-6a<sup>III</sup>), 3.93 (m, 1H, H-4<sup>I</sup>), 3.85–3.81 (m, 2H, H-6b<sup>II</sup>, 3<sup>III</sup>), 3.79–3.73 (m, 3H, H-3<sup>II</sup>, 5<sup>II</sup>, 5<sup>III</sup>), 3.76 (s, 3H, MeOPh), 3.70 (s, 3H, COOMe), 3.61 (dd, 1H,  $J_{4,5\text{ax}} = 5.61$  Hz, H-5ax<sup>I</sup>), 3.51 (dd, 1H,  $J_{5,6\text{a}} = 7.07$  Hz,  $J_{\text{gem}} = 11.47$  Hz, H-6a<sup>II</sup>), 2.40, 2.37, 2.37, 2.36, 2.30 (5s, 3H  $\times$  5, 5MePh), 2.16, 2.09, 2.02, 1.93, 1.89, 1.65 (6s, 3H  $\times$  6, 6MeCO). Anal. Calcd for  $\text{C}_{83}\text{H}_{88}\text{O}_{33} \cdot 4\text{H}_2\text{O}$ : C, 59.13; H, 5.57. Found: C, 59.01; H, 6.13. ESI-MS (positive) Calcd for  $\text{C}_{83}\text{H}_{88}\text{O}_{33}\text{Na}$   $[\text{M} + \text{Na}]^+$ , 1635.5. Found: 1635.3.

***O*-[Methyl 2,3,4-tri-*O*-(4-methylbenzoyl)- $\beta$ -D-glucopyranosyl uronate]-(1  $\rightarrow$  3)-*O*-(2,4,6-tri-*O*-acetyl- $\beta$ -D-galactopyranosyl)-(1  $\rightarrow$  3)-*O*-(2,4,6-tri-*O*-acetyl- $\beta$ -D-galactopyranosyl)-(1  $\rightarrow$  4)-2,3-di-*O*-(4-methylbenzoyl)- $\alpha$ -D-xylopyranosyl trichloroacetimidate (**17**)**

To a solution of **15** (445.0 mg, 0.276 mmol) in  $\text{CH}_3\text{CN}$  (36 mL) and  $\text{H}_2\text{O}$  (9 mL) was added CAN (453.6 mg, 0.827 mmol) at  $0^{\circ}\text{C}$  with stirring. After 3 h, additional CAN (150.0 mg, 0.274 mmol) was supplied. The reaction mixture was stirred for another 1.5 h and treated as the synthesis of **7**. The residue was eluted from a column of silica gel (3:1–2:1–1:1–1:5–1:8 *n*-hexane:EtOAc) to give **16** (385.8 mg, 93%) as a syrup. The obtained hemiacetal (**16**) was diluted with

CH<sub>2</sub>Cl<sub>2</sub> (17 mL). To the solution were added CCl<sub>3</sub>CN (260 μL) and 1 drop of DBU at 0°C. The reaction mixture was stirred for 2 h and directly subjected to a column of silica gel (3:1–1:1–1:3–1:4 *n*-hexane:EtOAc) to give **17** (377.2 mg, 89%) as a syrup, which was used for the glycosylation without further purification.

***N*-Butoxycarbonyl-*O*-{[methyl 2,3,4-tri-*O*-(4-methylbenzoyl)-β-*D*-glucopyranosyl uronate]-(1 → 3)-*O*-(2,4,6-tri-*O*-acetyl-β-*D*-galactopyranosyl)-(1 → 3)-*O*-(2,4,6-tri-*O*-acetyl-β-*D*-galactopyranosyl)-(1 → 4)-2,3-di-*O*-(4-methylbenzoyl)-β-*D*-xylopyranosyl}-*L*-serylglycine allyl ester (20)**

To a solution of **17** (409.7 mg, 248.0 μmol) and **18** (225.2 mg, 744.9 μmol) in CH<sub>2</sub>Cl<sub>2</sub> (47 mL) was added MS AW300 (1.8 g). This mixture was stirred for 30 min at rt, and then cooled to –20°C. To this solution was added a TMSOTf (9 μL, 0.05 mmol) with stirring, and the same amount of TMSOTf was added after 1 h, and stirring was continued for another 1.5 h. Then, the reaction was treated as described in the synthesis of **8**. The residue was subjected to the columns of gel permeation (LH-20, 1:1 CHCl<sub>3</sub>:MeOH) and silica gel (5:1–3:1–2:1–1:1–1:2 toluene:EtOAc) to give **20** (318.7 mg) in 72% yield: [α]<sub>D</sub> +31.6 (*c* 1.26, CHCl<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 7.80–7.62 (m, 10H, Ar H), 7.12–7.00 (m, 10H, Ar H), 6.82 [m, 1H, NH(Gly)], 5.80 (m, 1H, =CH), 5.72 (dd, 1H, *J*<sub>2,3</sub> = 9.03 Hz, *J*<sub>3,4</sub> = 9.76 Hz, H-3<sup>IV</sup>), 5.57 (t, 1H, H-4<sup>IV</sup>), 5.49 (t, 1H, *J*<sub>2,3</sub> = *J*<sub>3,4</sub> = 7.81 Hz, H-3<sup>I</sup>), 5.40 (d, 1H, *J*<sub>3,4</sub> = 3.17 Hz, H-4<sup>III</sup>), 5.36 [m, 1H, NH(Ser)], 5.28 (dd, 1H, *J*<sub>1,2</sub> = 7.08 Hz, H-2<sup>IV</sup>), 5.20 (m, 2H, =CH<sub>2</sub>), 5.14 (d, 1H, *J*<sub>3,4</sub> = 1.71 Hz, H-4<sup>II</sup>), 5.12 (dd, 1H, *J*<sub>1,2</sub> = 6.10 Hz, H-2<sup>I</sup>), 4.95 (dd, 1H, *J*<sub>1,2</sub> = 8.05 Hz, *J*<sub>2,3</sub> = 7.81 Hz, H-2<sup>II</sup>), 4.92 (dd, 1H, *J*<sub>1,2</sub> = 7.81 Hz, *J*<sub>2,3</sub> = 10.49 Hz, H-2<sup>III</sup>), 4.84 (d, 1H, H-1<sup>IV</sup>), 4.65 (d, 1H, H-1<sup>I</sup>), 4.46 (m, 2H, OCH<sub>2</sub>), 4.37 (d, 1H, H-1<sup>II</sup>), 4.29 (d, 1H, H-1<sup>III</sup>), 4.19 (d, 1H, H-5<sup>IV</sup>), 4.15 (m, 1H, Serα), 4.09 (dd, 1H, *J*<sub>5,6b</sub> = 6.10 Hz, *J*<sub>gem</sub> = 10.97 Hz, H-6b<sup>III</sup>), 4.06 (dd, 1H, *J*<sub>4,5eq</sub> = 5.61 Hz, *J*<sub>gem</sub> = 13.17 Hz, H-5eq<sup>I</sup>), 3.97 (dd, 1H, *J*<sub>5,6a</sub> = 5.86 Hz, H-6a<sup>III</sup>), 3.89 (m, 1H, H-4<sup>I</sup>), 3.89 (dd, 1H, *J*<sub>a,NH</sub> = 5.37 Hz, *J*<sub>gem</sub> = 18.54 Hz, Glya), 3.79 (dd, 1H, *J*<sub>b,NH</sub> = 5.37 Hz, Glyb), 3.76 (m, 1H, H-3<sup>III</sup>), 3.73 (dd, 1H, *J*<sub>5,6b</sub> = 5.85 Hz, *J*<sub>gem</sub> = 11.22 Hz, H-6b<sup>II</sup>), 3.68 (brt, 1H, *J* = 6.10 Hz, H-5<sup>III</sup>), 3.63 (m, 1H, H-3<sup>II</sup>), 3.63 (s, 3H, COOMe), 3.57 (brt, 1H, *J* = 6.10 Hz, H-5<sup>II</sup>), 3.48 (m, 2H, Serβ), 3.45 (m, 2H, H-5a<sup>I</sup>, 6a<sup>II</sup>), 2.29 (s, 6H, 2MePh), 2.28 (s, 6H, 2MePh), 2.22 (s, 3H, MePh), 2.07, 2.01, 1.94, 1.85, 1.83, 1.59 (6s, 3H × 6, 6MeCO), 1.36 (s, 9H, *tert*-Bu). Anal. Calcd for C<sub>89</sub>H<sub>102</sub>N<sub>2</sub>O<sub>37</sub> · H<sub>2</sub>O: C, 59.06; H, 5.80; N, 1.55. Found: C, 59.08; H, 5.97; N, 1.43.

***N*-(9-Fluorenylmethoxycarbonyl)-*O*-{[methyl 2,3,4-tri-*O*-(4-methylbenzoyl)-β-*D*-glucopyranosyl uronate]-(1 → 3)-*O*-(2,4,6-tri-*O*-acetyl-β-*D*-galactopyranosyl)-(1 → 3)-*O*-(2,4,6-tri-*O*-acetyl-β-*D*-galactopyranosyl)-(1 → 4)-2,3-di-*O*-(4-methylbenzoyl)-β-*D*-xylopyranosyl}-*L*-serylglycine allyl ester (21)**

To a solution of **17** (434.7 mg, 263.2 μmol) and **19** (335.1 mg, 789.5 μmol) in CH<sub>2</sub>Cl<sub>2</sub> (50 mL) was added MS AW300 (2.3 g). This mixture was stirred

for 30 min at rt and then cooled to  $-20^{\circ}\text{C}$ . To this solution was added a TMSOTf (15  $\mu\text{L}$ , 83  $\mu\text{mol}$ ) with stirring. Additional TMSOTf (15 and 10  $\mu\text{L}$ ) were added after 45 and 100 min, respectively, and stirring was continued for another 1 h. The reaction was quenched and treated as described in the synthesis of **8**. The residue was subjected to the columns of gel permeation (LH-20, 1:1  $\text{CHCl}_3\text{:MeOH}$ ) and silica gel (3:1–1:2–1:4 *n*-hexane:EtOAc) to give **21** (427.3 mg) in 74% yield:  $[\alpha]_{\text{D}} +37.1$  (c 0.41,  $\text{CHCl}_3$ );  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  7.98–7.56 (m, 11H, Ar H), 7.70 (d, 2H,  $J = 8.02$  Hz, Ar H), 7.57 (brt, 1H,  $J = 7.45$  Hz, Ar H), 7.40 (brt, 1H,  $J = 7.10$  Hz, Ar H), 7.31–7.22 (m, 2H, Ar H), 7.20–7.14 (m, 9H, Ar H), 7.08 (d, 2H,  $J = 8.02$  Hz, Ar H), 6.91 [brs, 1H,  $\text{NH}(\text{Gly})$ ], 5.87–5.79 (m, 1H,  $=\text{CH}$ ), 5.79 (brt, 1H,  $J = 9.28$  Hz,  $\text{H-3}^{\text{IV}}$ ), 5.64 (t, 1H,  $J_{3,4} = J_{4,5} = 9.63$  Hz,  $\text{H-4}^{\text{IV}}$ ), 5.58 (t, 1H,  $J_{2,3} = J_{3,4} = 8.02$  Hz,  $\text{H-3}^{\text{I}}$ ), 5.51 [m, 1H,  $\text{NH}(\text{Ser})$ ], 5.47 (d, 1H,  $J_{3,4} = 3.43$  Hz,  $\text{H-4}^{\text{III}}$ ), 5.35 (m, 1H,  $\text{H-2}^{\text{IV}}$ ), 5.31–5.23 (m, 2H,  $=\text{CH}_2$ ), 5.21 (m, 2H,  $\text{H-2}^{\text{I}}$ ,  $4^{\text{II}}$ ), 5.00 (m, 2H,  $\text{H-2}^{\text{II}}$ ,  $2^{\text{III}}$ ), 4.91 (d, 1H,  $\text{H-1}^{\text{IV}}$ ), 4.73 (m, 1H,  $\text{H-1}^{\text{I}}$ ), 4.54–4.38 (m, 1H,  $\text{Ser}\alpha$ ), 4.51 (m, 2H,  $\text{OCH}_2$ ), 4.42 (m, 1H,  $\text{H-1}^{\text{II}}$ ), 4.37–4.31 (m, 1H,  $\text{Ser}\beta\alpha$ ), 4.36 (m, 1H,  $\text{H-1}^{\text{III}}$ ), 4.26 (d, 1H,  $\text{H-5}^{\text{IV}}$ ), 4.17 (m, 1H,  $\text{H-6b}^{\text{III}}$ ), 4.12 (dd, 1H,  $J_{4,5\text{eq}} = 7.10$  Hz,  $J_{\text{gem}} = 14.44$  Hz,  $\text{H-5eq}^{\text{I}}$ ), 4.06–3.68 [m, 3H,  $\text{OCH}_2$  (Fmoc),  $\text{H-9}(\text{Fmoc})$ ], 4.04 (dd, 1H,  $J_{5,6a} = 6.64$  Hz,  $J_{\text{gem}} = 11.91$  Hz,  $\text{H-6a}^{\text{III}}$ ), 3.99 (m, 1H,  $\text{H-4}^{\text{I}}$ ), 3.93 (m, 1H, Glyb), 3.90–3.54 (m, 1H,  $\text{Ser}\beta\text{b}$ ), 3.89 (m, 1H, Glya), 3.82 (m, 1H,  $\text{H-3}^{\text{III}}$ ), 3.77 (m, 1H,  $\text{H-6b}^{\text{II}}$ ), 3.75 (brt, 1H,  $J = 5.96$  Hz,  $\text{H-5}^{\text{III}}$ ), 3.70 (s, 3H,  $\text{COOMe}$ ), 3.68 (m, 1H,  $\text{H-3}^{\text{II}}$ ), 3.63 (brt, 1H,  $J = 6.30$  Hz,  $\text{H-5}^{\text{II}}$ ), 3.53 (m, 2H,  $\text{H-5a}^{\text{I}}$ ,  $6a^{\text{II}}$ ), 2.36 (s, 12H, 4MePh), 2.29 (s, 3H, MePh), 2.15, 2.08, 2.05, 2.01, 1.92, 1.91 (6s,  $3\text{H} \times 6$ , 6MeCO). ESI-MS (positive) Calcd for  $\text{C}_{99}\text{H}_{105}\text{N}_2\text{O}_{37}$   $[\text{M} + \text{H}]^+$ , 1913.6. Found: 1913.2, Calcd for  $\text{C}_{99}\text{H}_{104}\text{N}_2\text{O}_{37}\text{Na}$   $[\text{M} + \text{Na}]^+$ , 1935.6. Found: 1935.1.

***N*-Butoxycarbonyl-*O*-{[methyl 2,3,4-tri-*O*-(4-methylbenzoyl)- $\beta$ -*D*-glucopyranosyl uronate]-(1  $\rightarrow$  3)-*O*-(2,4,6-tri-*O*-acetyl- $\beta$ -*D*-galactopyranosyl)-(1  $\rightarrow$  3)-*O*-(2,4,6-tri-*O*-acetyl- $\beta$ -*D*-galactopyranosyl)-(1  $\rightarrow$  4)-2,3-di-*O*-(4-methylbenzoyl)- $\beta$ -*D*-xylopyranosyl)-*L*-serylglycine (22)}**

Tetrakis(triphenylphosphine)palladium(0) (22.0 mg, 19.0  $\mu\text{mol}$ ) and *N*-methylaniline (105  $\mu\text{L}$ , 96.9  $\mu\text{mol}$ ) were added to the solution of **20** (173.5 mg, 96.6  $\mu\text{mol}$ ) in THF (4 mL) with stirring. Additional palladium catalysts (22 and 16 mg) were supplied to the solution after 2 and 3 h, respectively. One hour later, volatiles were removed under diminished pressure and the residue was subjected to the columns of gel permeation (LH-20, 10:10:1  $\text{CHCl}_3\text{:MeOH:AcOH}$ ) and silica gel (1:1–1:3 *n*-hexane:EtOAc) to give **22** (151.7 mg) in 90% yield. This compound was used for the next reaction without further purification.



***N*-(9-Fluorenylmethoxycarbonyl)-O-([methyl 2,3,4-tri-O-(4-methylbenzoyl)- $\beta$ -D-glucopyranosyl uronate]-(1  $\rightarrow$  3)-O-(2,4,6-tri-O-acetyl- $\beta$ -D-galactopyranosyl)-(1  $\rightarrow$  3)-O-(2,4,6-tri-O-acetyl- $\beta$ -D-galactopyranosyl)-(1  $\rightarrow$  4)-2,3-di-O-(4-methylbenzoyl)- $\beta$ -D-xylopyranosyl)-L-serylglycine (23)**

Tetrakis(triphenylphosphine)palladium(0) (45 mg, 39  $\mu$ mol) and *N*-methylaniline (211  $\mu$ L, 1.95 mmol) were added to the solution of **21** (427.3 mg, 195.1  $\mu$ mol) in THF (8.2 mL) with stirring. Additional palladium catalyst (22 mg) was supplied to the solution after 1.5 h. One hour later, volatiles were removed under diminished pressure and the residue was subjected to the column of silica gel (1:1–1:2–1:3 *n*-hexane:EtOAc–100:1:0.3–50:1:0.3–30:1:0.3 EtOAc:MeOH:AcOH) to give **23** (320.0 mg) in 76% yield. This compound was used for the next reaction without further purification.

***N*-(9-Fluorenylmethoxycarbonyl)-proline allyl ester (24)**

To a solution of Fmoc-Pro-H (2.00 g, 5.93 mmol) in ethanol (100 mL) was added a 6 mL of aq. Cs<sub>2</sub>CO<sub>3</sub> (0.965 g, 2.96 mmol) with stirring at 20°C. After 50 min, the reaction mixture was evaporated with toluene and dried in vacuo for 3 h. DMF (50 mL) and 3-bromopropene (0.57 mL, 6.75 mmol) were added to the residue and stirred overnight at rt. The reaction mixture was diluted with EtOAc and treated as described in general methods. The crude materials were subjected to the column of silica gel (11:1–7:1–5:1–4:1 *n*-hexane:EtOAc) to give **24** (1.93 g, 5.11 mmol) in 86% yield: [ $\alpha$ ]<sub>D</sub> –54.4 (c 1.76, CHCl<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  7.78–7.75 (m, 2H, Ar H), 7.64–7.54 (m, 2H, Ar H), 7.42–7.26 (m, 4H, Ar H), 5.89 (m, 1H, =CH), 5.28 (m, 2H, =CH<sub>2</sub>), 4.65 (d, 2H, OCH<sub>2</sub>), 4.41 (m, Pro $\alpha$ ), 4.44 [m, 1H, one of CH<sub>2</sub> (Fmoc)], 4.27 [m, 2H, one of CH<sub>2</sub> (Fmoc), H-9], 3.67 (m, 1H, Pro $\delta$ b), 3.55 (m, 1H, Pro $\delta$ a), 2.28 (m, 1H, Pro $\beta$ b), 2.05 (m, 1H, Pro $\beta$ a), 1.90 (m, 2H, Pro $\gamma$ ). Anal. Calcd. for C<sub>23</sub>H<sub>23</sub>NO<sub>4</sub> · 0.3H<sub>2</sub>O: C, 72.21; H, 6.17; N, 3.66. Found: C, 72.64; H, 6.45; N, 3.64.

***N*-(9-Fluorenylmethoxycarbonyl)-N<sup>i</sup>-(butoxycarbonyl)-L-tryptophanyl-L-proline allyl ester (25)**

Morpholine (4.9 mL) was added to a solution of **24** (429.4 mg, 1.137 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) and stirred overnight. The crude mixture was evaporated to dryness with toluene and diluted with CH<sub>2</sub>Cl<sub>2</sub> (22 mL). To this solution was added HOBt (278 mg, 2.08 mmol) with stirring. This solution was cooled to –20°C and HBTU (430 mg, 1.13 mmol) was added. The cooling bath was removed, and the reaction mixture was stirred at rt for 30 min. Then, Fmoc-Trp(Boc)-H (544.6 mg, 1.034 mmol) was added. The reaction mixture was stirred overnight and diluted with CHCl<sub>3</sub>. Organic phase was washed with 1M HCl and aq. NaHCO<sub>3</sub>, brine and dried over MgSO<sub>4</sub>. The volatiles were removed under diminished pressure. The crude materials were subjected

to the column of silica gel (7:1–6:1–5:1–3:1–1:1 *n*-hexane:EtOAc) to give **25** (391.4 mg) in 56% yield:  $[\alpha]_{\text{D}} -26.3$  (*c* 1.12,  $\text{CHCl}_3$ );  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  8.14 (m, 1H, Ar H), 7.78–7.23 (m, 11H, Ar H), 5.93 (m, 1H, =CH), 5.65 [d, 1H,  $J_{\alpha,\text{NH}} = 8.54$  Hz, NH(Trp)], 5.30 (m, 2H, =CH<sub>2</sub>), 4.87 (ddd,  $J_{\alpha,\beta\text{a}} = 7.08$  Hz,  $J_{\alpha,\beta\text{b}} = 5.86$  Hz, Trp $\alpha$ ), 4.66 (d, 2H, OCH<sub>2</sub>), 4.55 (dd,  $J_{\alpha,\beta\text{a}} = 4.88$  Hz,  $J_{\alpha,\beta\text{b}} = 8.05$  Hz, Pro $\alpha$ ), 4.37–4.15 [m, 3H, CH<sub>2</sub> (Fmoc), 9H], 3.61 (m, 1H, Pro $\delta\text{b}$ ), 3.25 (m, 1H, Pro $\delta\text{a}$ ), 3.22 (dd, 1H,  $J_{\text{gem}} = 14.64$  Hz, Trp $\beta\text{b}$ ), 3.11 (dd, 1H, Trp $\beta\text{a}$ ), 2.20 (m, 1H, Pro $\beta\text{b}$ ), 1.97 (m, 1H, Pro $\beta\text{a}$ ), 1.91 (m, 2H, Pro $\gamma$ ), 1.63 (s, 9H, *t*-Bu). Anal. Calcd for  $\text{C}_{39}\text{H}_{41}\text{N}_3\text{O}_7 \cdot 2.5\text{H}_2\text{O}$ : C, 66.07; H, 6.55; N, 5.93. Found: C, 65.94; H, 6.17; N, 5.71. ESI-MS (positive) Calcd for  $\text{C}_{39}\text{H}_{42}\text{N}_3\text{O}_7$   $[\text{M} + \text{H}]^+$ , 664.3. Found: 664.5, Calcd for  $\text{C}_{39}\text{H}_{41}\text{N}_3\text{O}_7\text{Na}$   $[\text{M} + \text{Na}]^+$ , 686.3. Found: 686.4.

***N*-(9-Fluorenylmethoxycarbonyl)-*N*<sup>*i*</sup>-(butoxycarbonyl)-*L*-tryptophanyl-*L*-proline (**26**)**

Tetrakis(triphenylphosphine)palladium(0) (247.3 mg, 0.241 mmol) and *N*-methylaniline (2.3 mL, 21 mmol) were added to the solution of **25** (1.447 g, 2.135 mmol) in THF (18 mL) with stirring. After 15 h, volatiles were removed under diminished pressure and the residue was subjected to a column of silica gel (5:1 toluene:EtOAc–100:1:0.3–50:1:0.3 EtOAc:MeOH:AcOH) to give **26** (861.9 mg) in 63% yield. This compound was used for the next reaction without further purification:  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  8.11–8.09 (d, 1H, Ar H), 7.74–7.21 (m, 11H, Ar H), 5.81 [d, 1H,  $J_{\alpha,\text{NH}} = 8.78$  Hz, NH(Trp)], 4.85 (dd,  $J_{\alpha,\beta\text{a}} = 7.32$  Hz,  $J_{\alpha,\beta\text{b}} = 7.08$  Hz, Trp $\alpha$ ), 4.57 (dd,  $J_{\alpha,\beta\text{a}} = 3.66$  Hz,  $J_{\alpha,\beta\text{b}} = 8.29$  Hz, Pro $\alpha$ ), 4.36–4.14 [m, 3H, CH<sub>2</sub> (Fmoc), 9H], 3.59 (m, 1H, Pro $\delta\text{b}$ ), 3.16 (m, 3H, Pro $\delta\text{a}$ , Trp $\beta$ ), 2.22 (m, 1H, Pro $\beta\text{b}$ ), 2.03 (m, 1H, Pro $\beta\text{a}$ ), 1.85 (m, 2H, Pro $\gamma$ ), 1.62 (s, 9H, *t*-Bu).

***N*-(Benzyloxycarbonyl)-*O*-(*tert*-butyl)-*L*-aspartylglycine *tert*-butyl ester (**27**)**

Z-Asp(*O*-*t*-Bu)-H (200.0 mg, 0.586 mmol) and H-Gly-*t*-Bu · HCl (98.2 mg, 0.586 mmol) were dissolved in  $\text{CH}_2\text{Cl}_2$  (18 mL). To this was added  $\text{Et}_3\text{N}$  (81  $\mu\text{L}$ , 0.59 mmol) and HOBT (158.4 mg, 1.172 mmol) with stirring. This solution was cooled to  $-20^\circ\text{C}$ , and then, a solution of WSCD · HCl (146 mg, 0.762 mmol) in  $\text{CH}_2\text{Cl}_2$  (5 mL) was added. The reaction mixture was stirred overnight and gradually raised to rt, diluted with  $\text{CHCl}_3$ , and treated as described in the synthesis of **25**. The crude materials were subjected to a column of silica gel (5:1–1:2 *n*-hexane:EtOAc) to give **27** (246.1 mg) in 96% yield:  $[\alpha]_{\text{D}} +15.0$  (*c* 1.45,  $\text{CHCl}_3$ );  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ):  $\delta$  7.37–7.27 (m, 5H, Ar H), 6.96 [m, 1H, NH(Gly)], 5.98 [d, 1H,  $J_{\alpha,\text{NH}} = 8.25$  Hz, NH(Asp)], 5.14 (s, 2H, PhCH<sub>2</sub>), 4.57 (brs, 1H, Asp $\alpha$ ), 3.92 (m, 2H, Gly), 2.92 (brdd, 1H,  $J = 4.38$  and 16.95 Hz, Asp $\beta\text{a}$ ), 2.64 (dd, 1H,  $J_{\alpha,\beta\text{b}} = 6.42$ ,  $J_{\text{gem}} = 13.95$  Hz,

Aspβb), 1.46, 1.43 (2s, 9H × 2, *t*-Bu). Anal. Calcd. for C<sub>22</sub>H<sub>32</sub>N<sub>2</sub>O<sub>7</sub>: C, 60.54; H, 7.39; N, 6.42. Found: C, 60.31; H, 7.59; N, 6.27.

***N*-(9-Fluorenylmethoxycarbonyl)-*N*<sup>i</sup>-(butoxycarbonyl)-*L*-tryptophanyl-*L*-prolyl-*O*-(*tert*-butyl)-*L*-aspartylglycine *tert*-butyl ester (**29**)**

A solution of **27** (246.1 mg, 563.8 μmol) in MeOH (5 mL) and AcOH (0.1 mL) containing a catalytic amount of Pd-C was stirred vigorously for 3 h under H<sub>2</sub> atmosphere. The reaction mixture was filtered on Celite and the volatiles were removed under diminished pressure to afford **28** (191.2 mg, 93%), which was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (20 mL). To this solution were added **26** (404.1 mg, 633.6 μmol), Et<sub>3</sub>N (73 μL, 0.53 mmol), and HOBt (171 mg, 1.27 mmol) with stirring. This solution was cooled to −20°C, and then a solution of WSCD · HCl (157 mg, 819 μmol) in CH<sub>2</sub>Cl<sub>2</sub> (6 mL) was added. The reaction mixture was stirred overnight gradually raised to rt, diluted with CHCl<sub>3</sub>, and treated as described in the synthesis of **25**. The crude materials were subjected to a column of silica gel (5:1–3:1–1:1–1:3–1:5 *n*-hexane:EtOAc) to give **29** (414.6 mg) in 85% yield: [α]<sub>D</sub> −20.7 (*c* 1.09, CHCl<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>): δ 7.79–7.21 [m, 13H, Ar H, NH (Asp)], 7.22 [m, 1H, NH (Gly)], 5.76 (d, 1H, *J*<sub>α,β</sub> = 5.50 and 8.02 Hz, Proα), 4.30 [dd, 1H, one of CH<sub>2</sub> (Fmoc)], 4.23 [dd, 1H, one of CH<sub>2</sub> (Fmoc)], 4.14 [m, 1H, H-9 (Fmoc)], 4.11 (m, 1H, Glya), 3.82 (m, 1H, Glyb), 3.79 (m, 1H, Proδb), 3.54 (m, 1H, Proδa), 3.28 (d, 2H, Trpβ), 3.00 (dd, 1H, *J*<sub>α,βa</sub> = 4.35 Hz, *J*<sub>gem</sub> = 16.95 Hz, Aspβa), 2.59 (dd, 1H, *J*<sub>α,βb</sub> = 5.96 Hz, Aspβb), 2.18 (m, 3H, Proβb, Trpβa, Proγb), 2.03 (m, 1H, Proγa), 1.66, 1.61, 1.46, (3s, 9H × 3, *t*-Bu). Anal. Calcd for C<sub>50</sub>H<sub>61</sub>N<sub>5</sub>O<sub>11</sub> · 3H<sub>2</sub>O: C, 62.41; H, 7.03; N, 7.28. Found: C, 62.22; H, 6.77; N, 7.07.

***N*-Butoxycarbonyl-*O*-{[methyl 2,3,4-tri-*O*-(4-methylbenzoyl)-β-*D*-glucopyranosyl uronate]-(1 → 3)-*O*-(2,4,6-tri-*O*-acetyl-β-*D*-galactopyranosyl)-(1 → 3)-*O*-(2,4,6-tri-*O*-acetyl-β-*D*-galactopyranosyl)-(1 → 4)-2,3-di-*O*-(4-methylbenzoyl)-β-*D*-xylopyranosyl}-*L*-serylglycyl-*N*<sup>i</sup>-(butoxycarbonyl)-*L*-tryptophanyl-*L*-prolyl-*O*-(*tert*-butyl)-*L*-aspartylglycine *tert*-Butyl Ester (**31**)**

Morpholine (0.9 mL) was added to a solution of **29** (110.5 mg, 119.8 μmol) in CH<sub>2</sub>Cl<sub>2</sub> (3.5 mL) and stirred overnight. The crude mixture was evaporated to dryness with toluene, and the residue was subjected to a column of silica gel (1:1 *n*-hexane:EtOAc–10:1:0.3 EtOAc:MeOH:Et<sub>3</sub>N) to afford **30** (83.6 mg) quantitatively. A part of **30** (41.3 mg, 59.0 μmol) was dissolved in DMF (130 μL), and HOBt (7.9 mg, 58 μmol) was added to the solution with stirring. This solution was cooled to −20°C and HBTU (12.0 mg, 31.6 μmol) was added. The cooling bath was removed, and the reaction mixture was stirred at rt for 30 min. Then, **22** (47.8 mg, 27.2 μmol) and *i*-Pr<sub>2</sub>EtN (10.3 μL, 59.1 μmol) in DMF (132 mL) were added. The reaction mixture was stirred overnight and diluted with EtOAc. Organic phase was treated as described

in the synthesis of **25**. The volatiles were removed under diminished pressure. The crude materials were subjected to a column of gel permeation (1:1 CHCl<sub>3</sub>:MeOH) to give **31** (41.5 mg) in 63% yield: [ $\alpha$ ]<sub>D</sub> +78 (c 0.1, CHCl<sub>3</sub>); <sup>1</sup>H NMR (CDCl<sub>3</sub>):  $\delta$  8.13–7.05 (m, 25H, Ar H), 7.45 [d, 1H,  $J_{\alpha,\text{NH}}$  = 8.25 Hz, NH (Asp)], 7.18 [m, 1H, NH (Gly)], 6.94 [br, 1H, NH (Gly)], 6.78 [br, 1H, NH (Trp)], 5.80 (t, 1H,  $J_{2,3} = J_{3,4} = 9.39$  Hz, H-3<sup>IV</sup>), 5.64 (brt, 1H,  $J = 9.26$  Hz, H-4<sup>IV</sup>), 5.53 (dd, 1H,  $J_{2,3} = 7.79$  Hz,  $J_{3,4} = 8.25$  Hz, H-3<sup>I</sup>), 5.47 (d, 1H,  $J_{3,4} = 2.75$  Hz, H-4<sup>III</sup>), 5.37 [m, 1H, NH (Ser)], 5.36 (dd, 1H,  $J_{1,2} = 7.10$  Hz, H-2<sup>IV</sup>), 5.20 (d, 1H,  $J_{3,4} = 3.21$  Hz, H-4<sup>II</sup>), 5.17 (d, 1H,  $J_{1,2} = 6.18$  Hz, H-2<sup>I</sup>), 5.06 (m, 1H, Trp $\alpha$ ), 4.99 (m, 2H, H-2<sup>II</sup>, 2<sup>III</sup>), 4.92 (d, 1H, H-1<sup>IV</sup>), 4.79 (m, 1H, Asp $\alpha$ ), 4.66 (d, 1H, H-1<sup>I</sup>), 4.49 (t, 1H,  $J = 6.42$  Hz, Pro $\alpha$ ), 4.40 (d, 1H,  $J_{1,2} = 8.02$  Hz, H-1<sup>III</sup> or <sup>II</sup>), 4.37 (d, 1H,  $J_{1,2} = 8.02$  Hz, H-1<sup>II</sup> or <sup>III</sup>), 4.28 (m, 1H, Ser $\alpha$ ), 4.27 (d, 1H,  $J_{4,5} = 9.85$  Hz, H-5<sup>IV</sup>), 4.16 (dd, 1H,  $J_{5,6b} = 5.73$  Hz,  $J_{\text{gem}} = 11.62$  Hz, H-6b<sup>III</sup>), 4.06 (m, 1H, H-5eq<sup>I</sup>), 4.03 (dd, 1H,  $J_{5,6a} = 5.27$  Hz, H-6a<sup>III</sup>), 4.00 (m, 1H, Glya), 3.92 (m, 1H, H-4<sup>I</sup>), 3.83 (dd,  $J_{2,3} = 10.31$  Hz, H-3<sup>III</sup>), 3.81 (m, 1H, Glyb), 3.8–3.5 (m, 2H, H-6<sup>II</sup>), 3.76 (brt,  $J = 6.56$  Hz, H-5<sup>III</sup>), 3.75 (m, 2H, Gly), 3.70 (s, 3H, COOMe), 3.69 (m, 1H, H-3<sup>II</sup>), 3.68 (m, 1H, Pro $\delta$ b), 3.61 (brt, 1H,  $J = 7.10$  Hz, H-5<sup>II</sup>), 3.55 (m, 2H, Ser $\beta$ ), 3.47 (dd, 1H,  $J_{4,5\text{ax}} = 8.01$  Hz,  $J_{\text{gem}} = 12.14$  Hz, H-5ax<sup>I</sup>), 3.37 (m, 1H, Pro $\delta$ a), 3.24 and 3.15 (m, 2H, Trp $\beta$ ), 2.90 (dd, 1H,  $J_{\alpha,\beta a} = 4.81$  Hz,  $J_{\text{gem}} = 16.96$  Hz, Asp $\beta$ a), 2.62 (dd, 1H,  $J_{\alpha,\beta b} = 5.95$  Hz, Asp $\beta$ b), 2.36, 2.35 (2s, 3H  $\times$  2, 2MePh), 2.34 (s, 6H, 2MePh), 2.29 (s, 3H, MePh), 2.15 (s, 6H, 2MeCO), 2.12 (m, 2H, Pro $\beta$ ), 2.09, 2.02, 1.93, 1.91 (4s, each 3H, 4MeCO), 1.92 (m, 2H, Pro $\gamma$ ), 1.66, 1.44, 1.40, 1.39 (4s, each 9H, 4*t*-Bu). ESI-MS (positive) Calcd. for C<sub>121</sub>H<sub>148</sub>N<sub>7</sub>O<sub>45</sub> [M + H]<sup>+</sup>, 2418.9. Found: 2418.8.

***O*-{ $\beta$ -D-Glucopyranuronosyl-(1  $\rightarrow$  3)-O- $\beta$ -D-galactopyranosyl-(1  $\rightarrow$  3)-O- $\beta$ -D-galactopyranosyl-(1  $\rightarrow$  4)- $\beta$ -D-xylopyranosyl}-L-serylglycyl-L-tryptophanyl-L-prolyl-L-aspartylglycine (1)**

To a solution of **31** (30.4 mg, 12.5  $\mu$ mol) in CH<sub>2</sub>Cl<sub>2</sub> (0.75 mL) was added a TFA cocktail (40.7:2.5:2.5:2.5:1.3:50 trifluoroacetic acid:thioanisole:phenol: H<sub>2</sub>O:1,2-ethanedithiol:triisopropylsilane:CH<sub>2</sub>Cl<sub>2</sub>) (0.75 mL) and stirred for 2 h. Volatiles were removed with toluene under diminished pressure and the residue was eluted from Bond Elut<sup>®</sup> C8 (H<sub>2</sub>O–1:1–1:2 H<sub>2</sub>O:MeOH–MeOH–CH<sub>2</sub>Cl<sub>2</sub>). The crude product was diluted with 50% MeOH (3 mL) and 0.107M NaOMe (1.4 mL) was added to the solution during 3 h keeping less than pH 9. Then, the solution was neutralized with 50% AcOH and the volatiles were removed under diminished pressure. The crude materials were subjected to a column of gel permeation (LH-20, 1% AcOH) to give **1** (14.8 mg) in 83% yield. <sup>1</sup>H NMR (D<sub>2</sub>O):  $\delta$  (selected) 7.71–7.20 (m, 5H, Ar H), 4.96 (t,  $J = 6.9$  Hz, Trp $\alpha$ ), 4.75 (dd,  $J = 6.0$  and  $7.0$  Hz, Asp $\alpha$ ), 4.70 (d,  $J_{1,2} = 7.9$  Hz, H-1<sup>IV</sup>), 4.66 (dd,  $J = 5.8$  and  $7.8$  Hz, Asp $\alpha'$ ), 4.59 (d,  $J_{1,2} = 7.9$  Hz, H-1<sup>III</sup>), 4.57 (m, Trp $\alpha'$ ), 4.56 (d,  $J_{1,2} = 7.5$  Hz, H-1<sup>II</sup>), 4.46 (d,  $J_{1,2} = 7.8$  Hz, H-1<sup>I</sup>), 4.42

(d,  $J_{1,2} = 7.3$  Hz, H-1' <sup>1</sup>), 4.41 (m, Pro $\alpha$ , Ser $\alpha$ ), 4.33 (dd,  $J = 5.0$  and 11.0 Hz, Ser $\beta$ a), 4.30 (t,  $J = 4.1$  Hz, Ser $\alpha'$ ), 4.24 (m, Ser $\beta$ a'), 4.14 (s, H-4<sup>IIorIII</sup>), 4.10 (s, H-4<sup>IIIorII</sup>), 4.10 (m, Ser $\beta$ b), 4.04 (m, Ser $\beta$ b'), 3.78 (m, H-3<sup>IIorIII</sup>), 3.73 (m, H-2<sup>II</sup>, 2<sup>III</sup>, Pro $\delta$ a), 3.63 (m, H-3<sup>IIIorII</sup>), 3.42 (m, H-2<sup>IV</sup>), 3.37 (m, H-2<sup>I</sup>, Pro $\delta$ b), 3.33 (m, H-2' <sup>1</sup>), 3.31 (m, Trp $\beta$ a), 3.30 (m, Pro $\delta$ b'), 3.14 (m, Trp $\beta$ b), 2.94 (dd,  $J_{gem} = 19.1$  Hz, Asp $\beta$ a), 2.90 (dd,  $J_{gem} = 19.1$  Hz, Asp $\beta$ a'), 2.79 (dd, Asp $\beta$ b), 2.77 (dd, Asp $\beta$ b'), 2.22 (m, Pro $\beta$ a), 1.92 (m, Pro $\beta$ b, Pro $\gamma$ ), 1.64, 1.44 (m, Pro $\beta'$ , Pro $\gamma$ a'), 0.97 (m, Pro $\gamma$ b'); FAB-MS (positive) Calcd. for C<sub>50</sub>H<sub>69</sub>N<sub>7</sub>O<sub>30</sub>Na<sub>3</sub> [M + H]<sup>+</sup>, 1316.4. Found: 1316.3. Calcd. for C<sub>50</sub>H<sub>68</sub>N<sub>7</sub>O<sub>30</sub>Na<sub>4</sub> [M + Na]<sup>+</sup>, 1338.4. Found: 1338.4.

## General Procedure for Solid-Phase Synthesis

Glycopeptides were synthesized manually for 100 mg of Sieber amide resin (52  $\mu$ mol) as follows. The Fmoc group was removed with 20% of piperidine/NMP (2 mL) (2  $\times$  3 min and 1  $\times$  20 min), and was monitored by Kaiser ninhydrin test. After washing with NMP (2.2 mL) (6  $\times$  1 min) and CH<sub>2</sub>Cl<sub>2</sub> (2.2 mL) (3  $\times$  1 min), it was dried in vacuo. The *N*-terminal free resin or peptide-on-resin was swollen in NMP (2 mL) and shaken overnight with corresponding Fmoc amino acid (130  $\mu$ mol), HOBt (130  $\mu$ mol), HBTU (130  $\mu$ mol), and *i*-Pr<sub>2</sub>EtN (260  $\mu$ mol). Coupling was monitored by Kaiser test. The resin was washed with NMP (2.2 mL) (3  $\times$  1 min) and CH<sub>2</sub>Cl<sub>2</sub> (2.2 mL) (3  $\times$  1 min) and dried in vacuo.

### *O*-{ $\beta$ -D-Glucopyranuronosyl-(1 $\rightarrow$ 3)-*O*- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 3)-*O*- $\beta$ -D-galactopyranosyl-(1 $\rightarrow$ 4)- $\beta$ -D-xylopyranosyl}-L-serylglycyl-L-tryptophanyl-L-prolyl-L-aspartylglycineamide (2)

Fmoc-Gly-H (46.4 mg, 156  $\mu$ mol), Fmoc-Asp(O-*t*-Bu)-H (64.1 mg, 156  $\mu$ mol), Fmoc-Pro-H (52.6 mg, 156  $\mu$ mol), and Fmoc-Trp(Boc)-H (82.1 mg, 156  $\mu$ mol) were coupled in turn on Sieber amide resin (300 mg, 156  $\mu$ mol) with HOBt (21.3 mg, 156  $\mu$ mol), HBTU (53.3 mg, 156  $\mu$ mol), and *i*-Pr<sub>2</sub>EtN (55  $\mu$ L, 312  $\mu$ mol) in NMP (2.7 mL). A part of the *N*-terminal of H-Trp(Boc)-Pro-Asp(O-*t*-Bu)-Gly-resin (67.4 mg, 27  $\mu$ mol) was coupled with **23** (116 mg, 54  $\mu$ mol) as described in the general procedure. The resultant resin was exposed to TFA cocktail (3.5 mL) with shaking for 7.5 h and evaporated. The residue was eluted from Bond Elut<sup>®</sup> C8 (1:0–0:1 H<sub>2</sub>O–MeOH) and subjected to a column of gel permeation (LH-20, 1% AcOH) to give crude glycopeptide (26.5 mg) still having acyl groups. Saponification was performed with 1 N NaOH–MeOH (1:100, 6.2 mL) overnight. The reaction mixture was diluted with H<sub>2</sub>O (6 mL), neutralized with 0.1 N AcOH after 5 h, and evaporated to dryness. The residue was subjected to further saponification with 5 mM NaOH (5 mL) for 4 days. The reaction mixture was neutralized in the same manner. The crude products were purified by HPLC (C18, A~B = 10%CH<sub>3</sub>CN + 0.1%CF<sub>3</sub>COOH~

90%CH<sub>3</sub>CN + 0.1%CF<sub>3</sub>COOH) to give **2**. TOF-MS (positive) Calcd. for C<sub>50</sub>H<sub>71</sub>N<sub>8</sub>O<sub>29</sub>Na [M + Na]<sup>+</sup>, 1271.4. Found: 1271.4, Calcd. for C<sub>50</sub>H<sub>70</sub>N<sub>8</sub>O<sub>29</sub>Na<sub>2</sub> [M – H + 2Na]<sup>+</sup>, 1293.4. Found: 1293.4.

## ACKNOWLEDGEMENT

This study was supported by a Grant-in-Aid for Scientific Research (C), 12660098, from the Ministry of Education, Science, Culture and Sport of Japan. The authors thank Ms. Miyuki Tanmatsu for the elemental analysis measurements (Research Center for Bioscience and Technology, Division of Instrumental Analysis, Tottori University).

## REFERENCES

- [1] (a) Sugahara, K.; Kitagawa, H. Recent advances in the study of the biosynthesis and functions of sulfated glycosaminoglycans. *Curr. Opin. Struct. Biol.* **2000**, *10*, 518–521; (b) Sugahara, K.; Kitagawa, H. Heparin and heparan sulfate biosynthesis. *IUBMB Life* **2002**, *54*, 163–175; (c) Silbert, J.E.; Sugumaran, G. Biosynthesis of chondroitin/dermatan sulfate. *IUBMB Life* **2002**, *54*, 177–186; (d) Kitagawa, H.; Izumikawa, T.; Uyama, Y.; Sugahara, K. Molecular cloning of a chondroitin polymerizing factor that cooperates with chondroitin synthase for chondroitin polymerization. *J. Biol. Chem.* **2003**, *278*, 23666–23671; (e) Funderbruch, L.L. Keratan sulfate biosynthesis. *IUBMB Life* **2002**, *54*, 187–194; (f) DeAngelis, P.L. Microbial glycosaminoglycan glycosyltransferases. *Glycobiology* **2002**, *12*, 9R–16R.
- [2] (a) Sugahara, K.; Yamada, S.; de Yoshida, K.; Waard, P.; Vliegthart, J.F.G. A novel sulfated structure in the carbohydrate-protein linkage region isolated from porcine intestinal heparin. *J. Biol. Chem.* **1992**, *267*, 1528–1533; (b) Sugahara, K.; Tsuda, H.; Yoshida, K.; Yamada, S.; de Beer, T.; Vliegthart, J.F.G. Structure determination of the octa- and decasaccharide sequences isolated from the carbohydrate-protein linkage region of porcine intestinal heparin. *J. Biol. Chem.* **1995**, *270*, 22914–22923.
- [3] Esko, J.D.; Zhang, L. Influence of core protein sequence on glycosaminoglycan assembly. *Curr. Opin. Struct. Biol.* **1996**, *6*, 663–670.
- [4] Zhang, L.; Esko, J.D. Amino acid determinants that drive heparan sulfate assembly in a proteoglycan. *J. Biol. Chem.* **1994**, *269*, 19295–19299.
- [5] López-Casillas, F.; Payne, H.M.; Andres, J.L.; Massagué, J. Betaglycan can act as a dual modulator of TGF-beta access to signaling receptors: mapping of ligand binding and GAG attachment sites. *J. Cell. Biol.* **1994**, *124*, 557–568.
- [6] Hudson, C.C.; Johnson, J.M. The isomeric tetracetates of xylose, and observations regarding the acetates of melibiose, trehalose and sucrose. *J. Am. Chem. Soc.* **1915**, *37*, 2748–2753.
- [7] Tamura, J.; Nishihara, J. Synthesis of phosphorylated and sulfated glycosyl serines in the linkage region of the glycosaminoglycans. *J. Org. Chem.* **2001**, *66*, 3074–3083.
- [8] Nakano, T.; Ito, Y.; Ogawa, T. Total synthesis of a sulfated glucuronyl glycosphingolipid, IV<sup>3</sup>GlcA(3-SO<sub>3</sub>)nLcOse<sub>4</sub>Cer, a carbohydrate epitope of neural cell adhesion molecules. *Tetrahedron Lett.* **1990**, *31*, 1597–1600.

- [9] Goto, F.; Ogawa, T. Synthesis of sulfated glycohexaose of linkage region of chondroitin 4-sulfate:  $\beta$ -D-GlcA-(1  $\rightarrow$  3){( $\text{SO}_3\text{Na} \rightarrow 4$ )}- $\beta$ -D-GalNAc-(1  $\rightarrow$  4)- $\beta$ -D-GlcA-(1  $\rightarrow$  3)- $\beta$ -D-Gal-(1  $\rightarrow$  3)- $\beta$ -D-Gal-(1  $\rightarrow$  4)-D-Xyl. *Tetrahedron Lett.* **1992**, 33, 6841–6844.
- [10] Rio, S.; Beau, J.-M.; Jacquinet, J.-C. Synthesis of glycopeptides from the carbohydrate-protein linkage region of proteoglycans. *Carbohydr. Res.* **1991**, 219, 71–90.
- [11] Liebe, B.; Kunz, H. Solid-phase synthesis of a sialyl-Tn-glycoundecapeptide of the MUC1 repeating unit. *Helv. Chim. Acta.* **1997**, 80, 1473–1482.
- [12] Uyama, T.; Kitagawa, H.; Tanaka, J.; Tamura, J.; Ogawa, T.; Sugahara, K. Molecular cloning and expression of a second chondroitin N-acetylgalactosaminyl transferase involving in the initiation and elongation of chondroitin/dermatan sulfate. *J. Biol. Chem.* **2003**, 278, 3072–3078.
- [13] Kim, B.-T.; Kitagawa, H.; Tanaka, J.; Tamura, J.; Sugahara, K. In vitro heparan sulfate polymerization. *J. Biol. Chem.* **2003**, 278, 41618–41623.