

TETRAHEDRON

Synthesis and Synthetic Applications of 3-Amino-∆⁵-piperidein-2-ones: Synthesis of Methionine-derived Pseudopeptides

M. Angels Estiarte,^a Marcus V. N. de Souza,^{a,b} Xavier del Rio,^a Robert H. Dodd,^b Mario Rubiralta,^a Anna Diez^{a#}

a. Laboratori de Química Orgànica. Facultat de Farmàcia. Universitat de Barcelona. 08028-Barcelona, Spain.
 b. Institut de Chimie des Substances Naturelles. CNRS. 91198 Gif-sur-Yvette Cédex. France.

Received 18 February 1999; revised 1 June 1999; accepted 17 June 1999

Abstract- A method is described for the efficient preparation of 3-amino- Δ^5 -piperidein-2ones 3. A synthetic application of enamides 3 has been achieved by methylthiolation on the C5 position to obtain the target methionine-derived pseudopeptides 2. © 1999 Elsevier Science Ltd. All rights reserved. **Keywords:** peptide mimetics, enamides, lactams, piperideinones

Introduction

In the context of our studies on the synthesis of conformationally restricted pseudopeptides¹ presenting a 3-amino-2-piperidone backbone,² and more specifically on the preparation of methionine derivatives as potential inhibitors of the hepatic transport of glutathione,³ we prepared compound 1 (Scheme 1) and its C3 epimer, derived from methionine and (*R*)-phenylglycinol.⁴ We now present the synthesis of protected pseudopeptides 2, with the double purpose of submitting them to pharmacological testing⁵ and of studying structure-activity relationships through introduction into longer peptide chains of known activity.⁶



Scheme 1

For the preparation of 1, we chose to introduce the SMe group on the C5 of a Δ^5 -piperidein-2-one, and to perform the C3 amination in the last steps of the synthesis. However, we could not apply the same strategy to the synthesis of the amino acid derivatives 2 because the C3 amination would be problematic. Thus, the use of a strong base such as *sec*-BuLi would racemize the amino acid labelled aa₂ in scheme 1, and the need for a free hydroxyl group on the side chain of the lactam nitrogen atom to obtain the C3 enolate conveniently,⁷ would limit the range of amino acids aa₂.

We therefore envisaged the use of a 3-amino- Δ^5 -piperidein-2-one type **3** as our new building block. In principle, compounds **3** could be easily prepared from commercially available glutamic acid, with the advantage of presenting the amino group in a prefixed configuration, and with the possibility of functionalising not only the C5 but also the C4 and C6 positions.

Results and Discussion

We chose to use di-*tert*-butylserine as the aa₂ because of its side chain functionality, and, in light of our previous work,^{2c,4} we first planned to prepare the corresponding enamide 3 (Prot = Alloc; R = ¹Bu; R' = CH₂O¹Bu) by reduction of imide 6. Imide 6 was prepared by reaction of anhydride 5⁸ with di-*tert*-butylserine with final addition of AcCl (Scheme 2). The subsequent reduction of imide 6 with NaBH₄ in THF yielded alcohol 7 as the only product, derived from the reduction of the intermediate open-chain aldehyde.⁹ Alcohol 7 was identified by the presence of two broad doublets at δ 5.80 and 6.98 in its ¹H NMR spectrum, corresponding to the amide and carbamate NH protons respectively, and by the multiplet centered at δ 3.70 due to the methylene group bearing the alcohol. The ¹³C NMR spectrum also showed the presence of the C5 methylene group (δ 62.1), and the structure was confirmed by mass spectrometry.



Scheme 2

Treatment of 6 with DIBAH gave an equimolar mixture of alcohol 7 and 6-hydroxylactams 8, which were separated by column chromatography. Hydroxylactams 8 were obtained as a mixture of the C6 epimers, as shown by the presence of split signals in the ¹³C NMR spectrum. The most relevant signals for assigning the structure were: i) the methine carbon at δ 83.2, which corresponded to C6 and indicated that the reduction had occurred; ii) the quaternary carbon of the *tert*-butoxycarbonyl at δ 81.6 and 82.4; and iii) the quaternary carbon of the *tert*-butoxy group at δ 72.9. When LiEt₃BH was used as the reducing agent we obtained the epimeric mixture of hydroxylactams 8 as the only isolable products. Unfortunately, all attempts to eliminate the C6 hydroxyl group of compound 8 to obtain the desired enamide type 3 were unsuccessful, and we recovered only the starting material or decomposition products when the reaction conditions were forced.¹⁰

In view of these results we decided to use aldehyde 9^{11} as the Δ^5 -piperidein-2-one precursor, by condensing it directly with di-*tert*-butylserine (Scheme 3). The reaction in refluxing C₆H₆ in a Dean-Stark trap in the presence of Et₃N (1 equivalent) gave the expected enamide, 11, but in very low yield (6%), and as a 4:1 mixture of epimers on C3. The most characteristic spectral data of compounds 11 were the signals at δ 5.21 and 6.50 in the ¹H NMR spectrum, corresponding to the double bond protons, and the olefinic methine carbons at δ 104.2 (C5) and 129.1 (C6) in the ¹³C NMR.



As an alternative, we coupled the racemic phthaloylthioester 12^{12} with di-*tert*-butylserine using the standard DCC-HOBt conditions, to obtain amides **13a,b** in 54% yield (Scheme 4). The subsequent reduction of the thioester with Et₃SiH and 10% Pd-C¹³ yielded the corresponding aldehyde that was directly cyclized using *p*-TsOH to obtain enamides **19a,b** in 45% yield.^{14,15} The most relevant data for the structural assignment of the enamides were the double bond NMR signals (see Experimental) and the absence of the thioester phenyl ring.

In order to establish the validity of the method, we applied the three-step sequence to prepare the {Met-Val} and {Met-Leu} pseudopeptides. The best results were obtained using a mixture of DMF and CH₂Cl₂ as the solvent for the coupling reaction to obtain amides **14** and **15**. The following reduction and cyclisation gave the corresponding enamides **20** and **21** as equimolar mixtures of the C3 epimers. The two diastereomers of both **20** and **21** were separated by column chromatography, and all enamides showed the characteristic double bond signals in the NMR spectra ($\delta_H \sim 5.3$ and 6.2; $\delta_C \sim 105$ and 126).¹⁵



Scheme 4

Once the synthesis of enamides of type 3 had been established, we proceeded to the preparation of the {Met-aa} derivatives 24-26 (Scheme 5). The methylthio group was introduced by treatment of enamides 19a,b, 20a, and 21a with MeSH in the presence of AIBN.⁴ The resulting type 2 piperidones, 24-26, showed the characteristic methyl group ($\delta_H \sim 2.2$; $\delta_C \sim 14$), and the loss of the double bond. From the four possible diastereomers of {Met-Ser} lactams 24¹⁵ (Scheme 6), only two were observed in the NMR spectra, in the proportion 3:1. This indicated that each pair, the 3,5-*trans*- (a and c) and the 3,5-*cis*-substituted (b and d), adopts the same ring conformation.



In the ¹H NMR spectrum of compounds 24 (*trans:cis* = 3:1), the 3-H proton appeared as a double doublet (J = 10 and 8 Hz), indicating that it is pseudoaxially disposed in both *cis* and *trans* diastereomers. A sample of only the major *trans* pair of diastereomers, **a** and **c**, was obtained by column chromatography. The signal corresponding to the pseudoaxial 4-H proton of these isomers was a double doublet of doublets (J = 10 and J = 10).

13, 10, and 6 Hz), which, together with the multiplicity observed for the corresponding 5-H proton (a double triplet of J = 7 and 6 Hz) indicated that the SMe group is pseudoaxial. Hence, we identified the major pair of piperidones **24a** and **24c** as the 3,5-*trans* diastereomers, and the minor pair as the 3,5-*cis*.¹⁶



Scheme 6

The methylthiolation of pure 20a in a sealed tube yielded a 3:1 mixture of piperidones *trans-*25a and *cis-*25b in 80% yield,¹⁵ from which the major isomer was obtained pure by column chromatography. As in the case of compound 20a, the relative disposition of the substituents was inferred from the signal multiplicity of the 3-H, the axial 4-H, and 5-H protons in the ¹H NMR spectrum, which clearly indicated that the phthalimido group was equatorial and that the methylthio substituent was axial in the major *trans* isomer. Similarly, the methylthiolation of pure 21a¹⁵ gave a 3:1 proportion of *trans-*26a and *cis-*26b. Since the 3:1 proportion is maintained regardless of the aa₂, we think that the steric hindrance that the C3 phthalimido group exerts on one side of the piperideinone ring is responsible for the stereoselection observed.

Compounds **24-26** are currently being tested as potential inhibitors of the hepatic transport of glutathione. Enamides of type **3** should allow us to functionalise the lactam ring on positions C4, C5, and C6 in the near future, and therefore provide access to a wide range of 3-aminopiperidin-2-one pseudodipeptides.

EXPERIMENTAL

General. Melting points were determined in a capillary tube on a Büchi apparatus. Optical rotations were measured with a Perkin-Elmer 241 polarimeter, at 23°C. ¹H and ¹³C NMR spectra were recorded on a Varian Gemini-200 instrument (200 MHz) and 2D NMR COSY experiments were performed on a Varian XL-500 instrument (500 MHz). Unless otherwise noted, NMR spectra were registered in CDCl₃, and chemical shifts are expressed in parts per million (δ) relative to internal Me₄Si. IR spectra were recorded on a Nicolet FT-IR spectrophotometer. Mass spectra were determined on a Hewlett-Packard 5988A mass spectrometer, either by chemical ionization (CIMS) or electronic impact

(EIMS). Flash column chromatography was carried out on SiO_2 (silica gel 60, 35-70 µm, SDS). TLC was performed on SiO_2 (silica gel 60 F254, Macherey-Nagel) and developed with the solvent described in each case for flash chromatography. The spots were located by UV light and KMnO₄. Purification of reagents and solvents was effected according to standard methods. Prior to concentration under reduced pressure, all extracts were dried over anhydrous Na₂SO₄ powder. Microanalyses were performed on a Carlo Erba 1106 analyzer by the Departament de Química Orgànica i Biològica CID, Barcelona.

(aS,3S)-3-Allyloxycarbonylamino-N-[2-tert-butoxy-1-(tert-butoxycarbonyl)ethyl]piperidin-2,6-

dione (6). To a solution of commercial L-glutamic acid (6 g, 40.3 mmol) in dioxane-H₂O (2:1, 120 ml) at 0°C, 2N NaOH (40 ml) was added (pH = 10). A solution of allyl chloroformate (5.2 ml, 48.9 mmol) in dioxane (20 ml) was slowly added, and the pH was readjusted with 2N NaOH. The reaction was stirred at 0°C for 1 h and at room temperature for 4 days. The reaction mixture was cooled to 0°C, acidified by addition of 1N aqueous NaHSO₄, and extracted with AcOEt. The organic extracts, dried and evaporated, yielded (*S)-N*-allyloxycarbonylglutamic acid as a yellow oil (7.82 g, 84%), which was used without further purification. IR (CHCl₃) 3400-2400 (CO₂H), 1730 (br s, CO), 1540 (C=C) cm⁻¹; ¹H NMR 2.20-2.30 (m, 2H, β -H), 2.50-2.60 (m, 2H, γ -H), 4.47 (q, *J* = 6 Hz, 1H, α -H), 4.58 (d, *J* = 5 Hz, 2H, OCH₂), 5.23 (dd, *J* = 10 and 1 Hz, 1H, CH=CH_A), 5.32 (br d, *J* = 17 Hz, 1H, CH=CH_B), 5.49 (br d, *J* = 6 Hz, 1H, NH), 5.85-5.98 (m, 1H, CH=CH₂); ¹³C NMR 26.9 (C- β), 29.6 (C- γ), 52.8 (C- α), 66.1 (OCH₂), 118.1 (CH=CH₂), 132.3 (CH=CH₂), 155.4 (NHCOAlloc), 176.0 and 177.9 (CO acids). CIMS *m/z* 232 (M⁺+1), 260 (M⁺+29).

To a solution of the allyloxycarbonylglutamic acid (10 g, 43.3 mmol) in dry THF (216 ml), cooled at -20°C and under N₂ atmosphere, Et₃N (6.6 ml, 47.6 mmol) was added. After 15 min at -20°C, a solution of MsCl (3.7 ml, 47.6 mmol) in dry THF (130 ml) and Et₃N (13.2 ml, 95.2 mmol) were sequentially added. After stirring for 4 h, the white precipitate of Et₃N·HCl was filtered off, and the solvent was evaporated to give (*S*)-*N*-**allyloxycarbonylglutamic anhydride (5)**⁸ as a yellow oil (quantitative). The crude product contained some Et₃N·HCl, but the anhydride was used without further purification due to its instability. ¹H NMR 2.26-2.36 (m, 2H, β-H), 2.91-2.94 (m, 2H, γ-H), 4.55-4.72 (m, 3H, OCH₂ and α-H), 5.20 (d, *J* = 10 Hz, 1H, CH=CH_A), 5.25, (d, *J* = 16 Hz, 1H, CH=CH_B), 5.80-5.95 (m, 1H, CH=CH₂); ¹³C NMR 22.9 (C-β), 29.7 (C-γ), 50.8 (C-α), 66.0 (OCH₂), 117.9 (CH=*C*H₂), 132.3 (*C*H=CH₂), 156.0 (NHCOAlloc), 165.5 and 166.8 (CO anhydride).

A solution of anhydride 5 (8.8 g, 41.52 mmol) and di-*tert*-butyl-L-serine (3.9 g, 18.9 mmol) in dry CHCl₃ (400 ml) was refluxed for 24 h. The CHCl₃ was evaporated, AcCl (110 ml, 1.27 mol) was added, and the mixture was refluxed for 5 h. The solvent was evaporated, and the residue was dissolved in CH₂Cl₂ (75 ml) and washed with H₂O. The organic extracts, dried and evaporated, yielded a yellow oil that was chromatographed (hexane:AcOEt, 1:1 to 1:9) to obtain *N*-acetylated di-*tert*-butylserine (700 mg, 15%) and pure imide **6** (3.5 g, 45%). [α]_D = -46.2 (c = 1.01, CHCl₃). mp 64-65°C (AcOEt). IR (CHCl₃) 3345 (NH), 1794, 1732 and 1689 (CO), 1535 (C=C); ¹H NMR 1.15 (s, 9H, OC(CH₃)₃), 1.47 (s, 9H, CO₂C(CH₃)₃), 2.20-2.30 (m, 2H, 4-H), 2.50 (ddd, *J* = 17, 8, and 4 Hz, 1H, 5-H_A), 2.78 (dt, *J* = 17 and 10 Hz, 1H, 5-H_B), 3.52 (dd, *J* = 9 and 3 Hz, 1H, β -H_B), 4.58-4.65 (m, 2H, 3-H and α -H), 4.73 (d, *J* = 8 Hz, 2H, OCH₂), 5.25 (d, *J* = 10 Hz, 1H, CH=CH_A), 5.40 (d, *J* = 10 Hz, 1H, CH=CH_B), 5.87-6.01 (m, 1H, CH=CH₂), 6.65 (d, *J* = 8 Hz, 1H, NH); ¹³C NMR 22.3 (C-4), 27.5 (OC(CH₃)₃), 27.9 (CO₂C(CH₃)₃), 31.3

10179

(C-5), 53.0 (C- α), 59.7 (C-3), 61.9 (C- β), 67.2 (OCH₂), 73.2 (OC(CH₃)₃), 82.0 (CO₂C(CH₃)₃), 118.7 (CH=CH₂), 131.2 (CH=CH₂), 151.2 (NHCOAlloc), 169.0 (C-2), 170.0 (C-6), 173.3 (CO₂C(CH₃)₃). CIMS *m*/z 413 (M⁺+1), 441 (M⁺+29), 453 (M⁺+41). Anal. Calcd for C₂₀H₃₂N₂O₇: C, 58.24; H, 7.82; N, 6.79. Found: C, 58.29; H, 7.86; N, 6.65.

Reduction of imide 6. Method A (7): To a solution of imide 6 (150 mg, 0.36 mmol) in dry THF (12 ml), at -20°C and under N2 atmosphere, NaBH4 (68 mg, 1.81 mmol) was added portionwise, and the mixture was stirred at the same temperature for 5 h. If the showed that there was still some starting material, NaBH₄ (68 mg, 1.81 mmol) was added and the reaction was stirred for 2 h more. The reaction was guenched by addition of saturated aqueous NH₄Cl. The solvent was evaporated and the residue, dissolved in CH₂Cl₂ (10 ml), was washed with saturated aqueous NH₄Cl (2 x 10 ml). The organic extracts, dried and evaporated, gave a yellow oil which was chromatographed (hexane:Et₂O, 1:1 to 3:7) to isolate hydroxyamide 7 (75 mg, 50%). [α]_D = +173.5 (c = 0.99, CHCl₃). IR (CHCl₃) 3400-3200 (OH), 1735 and 1657 (CO), 1530 (C=C) cm⁻¹; ¹H NMR (CDCl₃) 1.15 (s, 9H, OC(CH₃)₃), 1.46 (s, 9H, CO₂C(CH₃)₃), 1.63-1.73 (m, 2H, 4-H), 1.86 (q, J = 7 Hz, 1H, 3-H_A), 1.95 (q, J = 7 Hz, 1H, 3-H_B), 3.52 (dd, J = 9 and 3 Hz, 1H, β -H_A), 3.60 (br s, 1H, OH), 3.65-3.74 (m, 2H, 5-H), 3.79 (dd, J = 9 and 3 Hz, 1H, β -H_B), 4.38 (dd, J = 14 and 6 Hz, 1H, 2-H), 4.54-4.64 (m, 3H, α -H and OCH₂), 5.20 (dd, J = 10 and 1 Hz, 1H, CH=CH_A), 5.30 (dd, J = 17 and 1 Hz, 1H, CH=CH_B), 5.80 (br d, J = 108 Hz, 1H, NH), 5.83-5.98 (m, 1H, CH=CH₂), 6.98 (br d, J = 8 Hz, 1H, NH); ¹³C NMR (CDCl₃) 27.2 $(OC(CH_3)_3)$, 27.8 (C-3), 27.9 $(CO_2C(CH_3)_3)$, 30.2 (C-4), 53.1 (C- α), 54.1 (C-2), 61.9 (C- β), 62.1 (C-5), 65.7 (OCH₂), 73.1 (OC(CH₃)₃), 81.9 (CO₂C(CH₃)₃), 117.6 (CH=CH₂), 132.6 (CH=CH₂), 156.0 (NHCOAlloc), 169.2 (C-1), 171.3 (CO₂C(CH₃)₃). CIMS m/z 417 (M⁺+1), 445 (M⁺+29), 457 (M⁺+41). Anal. Calcd. for C₂₀H₃₆N₂O₇: C, 57.67; H, 8.71; N, 6.73. Found: C, 57.46; H, 8.86; N, 6.46.

<u>Method B (7 and 8)</u>: To a solution of imide 6 (1 g, 2.42 mmol) in dry THF (12 ml), at -78°C and under N₂ atmosphere, DIBAH (7.28 ml, 7.28 mmol) was slowly added and the mixture was stirred for 1 h at this temperature. The reaction was quenched with saturated aqueous NH₄Cl. The solvent was evaporated, and the residue, dissolved in AcOEt (50 ml), was washed with saturated aqueous NH₄Cl (2 x 25 ml) and with brine. The organic extracts, dried and evaporated, gave a yellow oil which was chromatographed (hexane:Et₂O, 1:1 to 3:7 gradient) to yield a mixture of pure hydroxyamide 7 (lower Rf, 31%) and hydroxylactams 8a,b (higher Rf, 39%, a:b = 2.3:1)

Method C (8): To a solution of imide 6 (1 g, 2.42 mmol) in dry THF (12 ml), cooled to -78°C and under N₂ atmosphere, LiEt₃BH (6 ml, 6 mmol) was slowly added and the mixture was stirred for 2 h at the same temperature. The reaction was quenched by addition of saturated aqueous NaHCO₃ and 30% H₂O₂ (2 ml) to destroy the excess hydride. The solvent was evaporated, and the residue was dissolved in Et₂O (50 ml) and washed with H₂O. The organic extracts, dried and evaporated, yielded hydroxylactams 8 as a yellow oil (40-50%): IR (CHCl₃) 3400-3200 (OH), 1750, 1716 and 1656 (CO), 1530 (C=C) cm⁻¹; ¹H NMR (from the diastereomeric mixture **a**,**b**) 1.14 (s, 9H, OC(CH₃)₃), 1.45 (s, 9H, CO₂C(CH₃)₃), 1.90-2.05 (m, 1H, 5-H_A), 2.10-2.20 (m, 1H, 5-H_B), 2.15-2.30 (m, 1H, 4-H), 3.50-3.55 (m, 1H, β-H_A), 3.75-3.80 (m, 1H, β-H_B), 4.35 (br s, 1H, 3-H), 4.55-4.60 (m, 1H, α-H), 4.65 (br s, 2H, OCH₂), 5.15-5.30 (m, 2H, CH=CH₂), 5.60 and 5.70 (2 br s, 1 H each, 6-H), 5.80-6.00 (C*H*=CH₂), 6.95 and 7.20 (2 br s, 1H each, NH); ¹³C NMR 27.3 (OC(CH₃)₃), 27.9 (CO₂C(CH₃)₃), 28.6 (C-4), 31.7 and 33.8 (C-5), 53.0 and 53.3 (C-α), 60.9 and 61.1 (C-3), 61.9 and 62.1 (C-β), 66.4 (OCH₂), 72.9 (OC(CH₃)₃), 81.6 and 82.4 (CO₂C(CH₃)₃), 83.2 (C-6), 117.7 and 117.9

 $(CH=CH_2)$, 132.3 ($CH=CH_2$), 154.5 (NHCOAlloc), 168.9 (C-2), 172.0 ($CO_2C(CH_3)_3$). CIMS *m/z* 415 (M⁺+1), 443 (M⁺+29), 455 (M⁺+41). Anal. Calcd for $C_{20}H_{34}N_2O_7$: C, 57.97; H, 8.21; N, 6.76. Found: C, 58.02; H, 8.43; N, 6.57.

(αS,3RS)-N-[2-tert-Butoxy-1-(tert-butoxycarbonyl)ethyl]-3-benzyloxycarbonyl-amino-∆5-

piperidein-2-ones (11a,b). To a solution of di-*tert*-butyl serine-hydrochloride (183 mg, 0.72 mmol) and Et₃N (0.1 ml, 0.72 mmol) in dry C₆H₆ (2 ml), at 10°C and under N₂ atmosphere, a solution of aldehyde 9¹² (200 mg, 0.72 mmol) in dry C₆H₆ (4 ml) was added. The mixture was refluxed for 48 h in a Dean-Stark trap. Once cold, the mixture was filtered and the solvent was evaporated. The resulting oil was a complex mixture from which enamides 11 (6%) were the only isolable product by column chromatography (hexane:Et₂O, 8:2 to 7:3), as a 1:4 C3 epimeric mixture. IR (CHCl₃) 3400 (NH), 1731 and 1678 (CO) cm⁻¹; ¹H NMR (CDCl₃) 1.16 and 1.28[°] (2s, OC(CH₃)₃), 1.45 and 1.58^{*} (2s, CO₂C(CH₃)₃), 2.25 (tt, *J* = 15 and 3 Hz, 1H, 4-H_A), 2.32^{*} (td, *J* = 15 and 7 Hz, 1H, 4-H_A), 2.84 (dt, *J* = 15 and 7 Hz, 4-H_B), 3.64 (dd, *J* = 10 and 3 Hz, 1H, 6-H_A), 3.70^{*} (br t, *J* = 6 Hz, 1H, β-H_A), 3.91 (dd, *J* = 10 and 6 Hz, 2H, β-H_B), 4.23^{*} (m, 1H, 3-H), 4.34 (dt, *J* = 14 and 7 Hz, 1H, 3-H), 5.12 (br s, α-H and CH₂C₆H₅), 5.21 (br t, *J* = 6 Hz, 5-H), 5.79 (br d, *J* = 4 Hz, NH), 6.50 (dd, *J* = 8 and 3 Hz, 6-H), 7.35 (br s, C₆H₅); ¹³C NMR (CDCl₃) 27.3 and 27.7^{*} (OC(CH₃)₃), 27.8 and 27.9^{*} (CO₂C(CH₃)₃), 29.3 (C-4), 50.9 (C-α), 56.9 (C-3), 61.4 (C-β), 66.8 (CH₂C₆H₅), 73.5 (OC(CH₃)₃), 82.2 (CO₂C(CH₃)₃), 104.2 (C-5), 128.0 (C₆H₅-o), 128.5 (C₆H₅-m), 129.1 (C₆H₅-ρ and C-6), 136.4 (C₆H₅-*i*), 168.3 (C-2), 179.0 (CO). EIMS *m/z* (%) 446 (M⁺, 0.1), 373 (7), 317 (4), 96 (40), 57 (100).

4-Phenylthiocarbonyl-2-(N-phthaloyl)-N-[1-(tert-butoxycarbonyl)-2-tert-butoxyethyl]butanamides

(13a,b). To a solution of thioester 1213 (500 mg, 1.35 mmol) in dry CH2Cl2 (7 ml) at 0°C and in N2 atmosphere, HOBt (183 mg, 1.35 mmol) and DCC (335 mg, 1.62 mmol) were added at a 5 min interval. After 15 min, di-tert-butyl serine hydrochloride (343 mg, 1.35 mmol) and Et₃N (0.37 ml, 2.71 mmol) were added. The intense yellow suspension was stirred for 2 h at room temperature. The mixture was filtered, and washed with 1N HCI. The organic phase, dried and evaporated, gave a yellow oil which was chromatographed (hexane:AcOEt, 8:2 to 1:1) to yield amide 13 as a white foam (54%). IR (CHCl₃) 3300 (NH), 1790, 1720, 1686 and 1530 (CO) cm⁻¹; ¹H NMR (CDCl₃) 1.03 and 1.11 (s, 9H each, OC(CH₃)₃), 1.42 and 1.43 (2s, 9H each, CO₂C(CH₃)₃), 2.60-2.72 (m, 4H, 3-H), 2.73-2.82 (m, 4H, 4-H), 3.49 and 3.53 (2dd, J = 8 and 3 Hz, 1H each, β -H_A), 3.73 and 3.75 (2dd, J = 6 and 3 Hz, 1H each, β -H_B), 4.55 and 4.60 (2dt, J = 8 and 3 Hz, 1H each, α -H), 4.91 (br t, J = 8 Hz, 2H, 2-H), 6.73 and 6.85 (2br d, J = 8 Hz, 1H each, NH), 7.30-7.45 (m, C₆H₅), 7.72-7.80 (m, 4H, Pht-α), 7.85-7.92 (m, 4H, Pht-β); ¹³C NMR (CDCl₃) 24.3 (C-3), 27.2 (OC(CH₃)₃), 27.8 $(CO_2C(CH_3)_3)$, 40.0 (C-4), 52.9 (C-2), 53.5 (C- α), 61.8 (C- β), 72.8 and 72.9 (OC(CH_3)_3), 81.9 (CO₂C(CH₃)₃), 123.5 (Pht-α), 129.0 (C₆H₅-o), 129.3 (C₆H₅-m and -p), 132.2 (Pht-quaternary), 134.3 (Phtβ), 134.4 (C₆H₅-i), 167.3 and 167.4 (C-1), 167.6 and 167.7 (CO₂C(CH₃)₃), 195.9 and 196.0 (C-5). EIMS m/z (%) 568 (M+,1), 495 (2), 459 (3), 403 (2), 329 (46), 301 (21), 186 (48), 57 (100). Anal. Calcd for C₃₀H₃₆N₂O₇S: C, 63.36; H, 6.38; N, 4.93; S, 5.64. Found: C, 63.29; H, 6.34; N, 4.91; S, 5.54.

The star indicates the signals corresponding to the minor isomer.

4-Phenylthiocarbonyl-2-(N-phthaloyl)-N-[1-(methoxycarbonyl)-2-methylpropyl]butanamides (14a,b). To a solution of thioester 12 (1 g, 2.7 mmol) in dry DMF-CH2Cl2 (5:100) at room temperature, a solution of HOBt (0.43 g, 3.25 mmol) in dry DMF (5 ml) was added at room temperature. After 5 min DCC (0.82 g, 3.95 mmol) was added and the resulting yellow solution was stirred for 10 min before the addition of L-valine methyl ester (0.42 g, 3.2 mmol). The white suspension was stirred for 12 h at room temperature. The precipitated ureas were filtered off, the solvent was evaporated, and the residue was redissolved in Et₂O. The organic solution was washed with H₂O, citric acid (pH = 4), and 1N HCl, then dried and evaporated to give an oil which was chromatographed (hexane:AcOEt, 7:3) to obtain the amides 14 (54%), as a 1:1 epimeric mixture. IR (CHCl₃) 3376 (NH), 1777, 1718, 1696 and 1532 (CO) cm⁻¹; ¹H NMR (CDCl₃) 0.84 and 0.89 (2d, J = 7 Hz, y-H), 2.15-2.19 (m, β -H), 2.61-2.68 (m, 3-H), 2.76 (t, J = 4 Hz, 4-H), 3.68 and 3.69 (2s, 3H each, CO₂CH₃), 4.50 and 4.53 (2dd, J = 5 and 3 Hz, 1H each, α-H), 4.85 (t, J = 7 Hz, 2-H), 6.60 (br d, J = 8 Hz, NH), 7.30-7.45 (m, C₆H₅), 7.72-7.80 (m, Pht-α), 7.85-7.92 (m, Pht-β); ¹³C NMR (CDCl₃) 17.6 and 18.9 (C-γ), 24.5 (C-3), 31.1 (C-β), 40.0 (C-4), 52.2 (CO₂CH₃), 53.5 (C-2), 57.5 (C-α), 123.5 (Pht-α), 129.0 (C₆H₅-*o*), 129.3 (C₆H₅-m and -p), 132.2 (Pht-quaternary), 134.3 (Pht-β), 134.4 (C₆H₅-i), 168.0 (C-1), 172.0 (CO₂CH₃), 196.4 (C-5). EIMS m/z (%) 483 (M++1, 0.1), 373 (24), 313 (100), 186 (71), 138 (61). Anal. Cald for C25H26N2O6S: C, 62.23; H, 5.43; N, 5.81; S, 6.65. Found: C, 62.08; H, 5.61; N, 5.88; S, 6.35.

4-Phenylthiocarbonyl-2-(N-phthaloyl)-N-[1-(methoxycarbonyl)-3-methylbutyl]butanamides (15a,b).

Operating as above, from thioester 12 (1 g, 2.7 mmol), HOBt (0.36 g, 2.7 mmol), DCC (0.66 g, 3.24 mmol), and L-leucine methyl ester (0.32 g, 3.2 mmol) in DMF-CH₂Cl₂ (105 ml), compounds 15 (68%) were obtained, as a 1:1 epimeric mixture, after column chromatography (hexane:AcOEt, 7:3): IR (CHCl₃) 3369 (NH), 1776, 1719 and 1533 (CO) cm⁻¹; ¹H NMR (CDCl₃) 0.87-0.92 (m, δ -H), 1.48-1.67 (m, β -H and γ -H), 2.64-2.80 (m, 3-H and 4-H), 3.64 and 3.67 (2s, 3H each, CO₂CH₃), 4.57-4.67 (m, α -H), 4.82-4.96 (m, 2-H), 6.61 and 6.65 (2 d, *J* = 8 Hz, NH), 7.30-7.45 (m, C₆H₅), 7.72-7.80 (m, Pht- α), 7.85-7.92 (m, Pht- β); ¹³C NMR (CDCl₃) 21.7 and 22.6 (C- δ), 24.5 (C-3), 24.7 (C- γ), 39.9 (C- β), 41.2 (C-4), 50.8 (C-2), 52.3 (CO₂CH₃), 53.2 (C- α), 123.5 (Pht- α), 129.0 (C₆H₅- σ), 129.3 (C₆H₅-m and -p), 132.2 (Pht-*quaternary*), 134.3 (Pht- β), 134.4 (C₆H₅-i), 167.9 (C-1), 173.1 (*C*O₂CH₃), 196.4 (C-5). EIMS *m*/*z* 497 (M⁺+1, 0.1), 387 (12), 355 (20), 327 (65), 259 (28), 186 (100). Anal. Calcd for C₂₆H₂₈N₂O₆S: C, 62.89; H, 5.68; N, 5.64. Found: C, 62.62; H, 5.68; N, 5.69.

$(\alpha S, 3RS) - N-[1-(tert-Butoxycarbonyl)-2-tert-butoxyethyl]-3-phthaloyl-<math>\Delta^5$ -piperidein-2-ones

(19a,b). To a solution of amide 13 (1.3 g, 2.28 mmol) in dry CH₂Cl₂-CH₃CN (1:1, 23 ml), in the presence of 4Å molecular sieves and 10% Pd-C (130 mg) Et₃SiH (1.27 ml, 8.01 mmol) was slowly added. After 30 min, the reaction mixture was filtered and the solvent evaporated. The resulting aldehyde was dissolved in toluene (230 ml), *p*-TsOH (87 mg, 0.45 mmol) was added, and the mixture was refluxed in a Dean-Stark trap for 1 h. The reaction mixture was filtered, the solvent evaporated, and the resulting oil was chromatographed (hexane:AcOEt, 8:2) to obtain enamides **19a,b** as a white foam (1:1 epimeric mixture, 45%). IR (CHCl₃) 1780, 1750 and 1680 (CO), 1650 (C=C); ¹H NMR (CDCl₃) 1.19 (s, OC(CH₃)₃), 1.46 and 1.47* (2s, 9H each, CO₂C(CH₃)₃), 2.30-2.45 (m, 4-H_A), 3.29 and 3.37 (2tt, *J* = 16 and 2 Hz, 1H each, 4-H_B), 3.68 (dt, *J* = 10 and 3 Hz, β -H_A), 3.92 (dd, *J* = 10 and 6 Hz, β -H_B), 5.02 (dd, *J* = 6 and 3 Hz, α -H), 5.06-5.15 (m, 3-H), 5.15-5.23 (m, 5-H), 6.52 and 6.56 (2dd, *J* = 8 and 3 Hz, 6-H), 7.70-7.75 (m, Pht- α), 7.85-7.90 (m, Pht- β); ¹³C NMR

(CDCl₃) 24.1 and 24.2 (C-4), 27.2 and 27.3 (OC(CH₃)₃), 27.7 and 27.9 (CO₂C(CH₃)₃), 49.4 and 49.5 (C-3), 56.8 and 57.7 (C- α), 61.4 and 61.6 (C- β), 73.4 (OC(CH₃)₃), 82.0 (CO₂C(CH₃)₃), 102.7 and 102.8 (C-5), 123.5 (Pht- α), 129.5 and 129.9 (C-6), 132.0 (Pht-*quaternary*), 134.0 (Pht- β), 165.5 (Pht-CO), 167.7 (C-2). EIMS *m*/*z* (%) 442 (M⁺, 2), 386 (4), 369 (2), 331 (8), 313 (13), 183 (17), 96 (43), 57 (100). Anal. calcd for C₂₄H₃₀N₂O₆: C, 65.14; H, 6.83; N, 6.33. Found: C, 65.03; H, 6.97; N, 6.18.

 $(\alpha S, 3RS)-N-[1-(Methoxycarbonyl)-2-methylpropyl]-3-phthaloyl-\Delta^5-piperidein-2-ones (20a,b).$ To a solution of amide 14 (700 mg, 1.45 mmol) in dry acetone (30 ml) and in the presence of 4Å molecular sieves and 10% Pd-C (60 mg), Et₃SiH (0.8 ml, 6.6 mmol) was slowly added. The resulting mixture was stirred at room temperature for 20 min, the solvent was evaporated, and the residue was dissolved in toluene (100 ml). p-TsOH (55 mg, 0.29 mmol) was added and the reaction was refluxed for 3 h. The reaction mixture was filtered, and the solvent evaporated to yield an oil, which was carefully chromatographed (hexane:AcOEt, 8:2) to afford the pure isomers of the enamides, as colorless oils. Enamide (α *S*,3*S*^{*})-20a (lower Rf, 22%): [α]_D = -158 (c = 1.00, CHCl₃). IR (CHCl₃) 1779, 1720 and 1689 (CO), 1389 (C=C) cm⁻¹; ¹H NMR (CDCl₃) 0.90 and 1.01 (2d, J = 6 Hz, 3H each, γ -H), 2.25-2.33 (m, 2H, β -H), 2.36 (ddd, J = 16, 8 and 7 Hz, 1H, 4-H_A), 3.25 (dddd, J= 16, 15, 3 and 2 Hz, 1H, 4-H_B), 3.76 (s, 3H, CO₂CH₃), 4.96 (d, J = 12 Hz, 1H, α-H), 5.09 (dd, J = 16 and 6 Hz, 1H, 3-H), 5.25 (ddd, J = 9, 7 and 2 Hz, 1H, 5-H), 6.34 (dd, J = 9 and 3 Hz, 1H, 6-H), 7.70-7.75 (m, 2H, Pht-α), 7.85-7.90 (m, 2H, Pht-β); ¹³C NMR (CDCl₃) 18.5 and 19.7 (C-γ), 24.5 (C-4), 28.7 (C-β), 49.2 (C-3), 52.2 (CO₂CH₃), 61.4 (C-α), 104.4 (C-5), 123.5 (Pht-α), 126.5 (C-6), 132.0 (Pht-quaternary), 134.0 (Pht-β), 165.5 (Pht-CO), 167.6 (C-2), 170.7 (CO2CH3). EIMS m/z: 357 (M++1), 385 (M++29), 398 (M++41). Anal. Cald for C19H20N2O5: C, 64.04; H, 5.62; N, 7.86. Found: C, 63.98; H, 5.87; N, 7.78. Enamide (aS,3R*)-20b (higher Rf, 22%): $[\alpha]_D = + 34$ (c = 1.00, CHCl₃). ¹H NMR (CDCl₃) 1.00 and 1.02 (2d, J = 6 Hz, 3H each, γ -H), 2.15-2.27 (m, 2H, β -H), 2.41 (ddd, J = 16, 8 and 7 Hz, 1H, 4-H_A), 3.29 (dddd, J = 16, 15, 3 and 2 Hz, 1H, 4-H_B), 3.73 (s, 3H, CO₂CH₃), 4.96 (d, J = 10 Hz, 1H, α -H), 5.12 (dd, J = 15 and 7 Hz, 1H, 3-H), 5.32 (ddd, J = 9, 7 and 2 Hz, 1H, 5-H), 6.43 (dd, J = 9 and 3 Hz, 1H, 6-H), 7.70-7.75 (m, 2H, Pht- α), 7.85-7.90 (m, 2H, Pht- β); ¹³C NMR (CDCl₃) 18.4 and 19.4 (C- γ), 24.1 (C-4), 29.4 (C- β), 49.1 (C-3), 52.1 (CO₂CH₃), 60.5 (Cα), 105.2 (C-5), 123.5 (Pht-α), 126.6 (C-6), 132.0 (Pht-quaternary), 134.0 (Pht-β), 165.5 (Pht-CO), 167.6 (C-2), 171.0 (CO₂CH₃).

$(\alpha S, 3RS)-N-[1-(Methoxycarbonyl)-3-methylbutyl]-3-phthaloyl-<math>\Delta^5$ -piperidein-2-ones (21a,b).

Operating as above, from compound **15** (800 mg, 1.61 mmol), 4Å molecular sieves, 10% Pd-C (60 mg), Et₃SiH (0.8 ml, 6.6 mmol), in acetone (30 ml), and *p*-TsOH (60 mg, 0.3 mmol) in toluene (100 ml), an equimolar mixture of enamides **21a,b** was obtained, as a colorless oil. After repeated column chromatography (hexane:AcOEt, 8:2), diastereomers **21a** (32%) and **21b** (9%) were isolated pure, and a mixture fraction (39%, **a:b** = 1:3) was also obtained. Enamide (α *S*,3*S*²,-**21a** (higher Rf): [α]_D = -96 (c = 0.5, CHCl₃). ¹H NMR (CDCl₃) 0.98 (d, *J* = 6 Hz, 6H, δ -H), 1.68-1.81 (m, 3H, β -H and γ -H), 2.41 (dt, *J* = 16 and 7 Hz, 1H, 4-H_A), 3.27 (tt, *J* = 16 and 3 Hz, 1H, 4-H_B), 3.72 (s, 3H, CO₂CH₃), 5.12 (dd, *J* = 15 and 7 Hz, 1H, 3-H), 5.32 (br t, *J* = 13 Hz, 1H, α -H), 5.29-5.37 (m, 1H, 5-H), 6.19 (dd, *J* = 8 and 3 Hz,1H, 6-H), 7.70-7.75 (m, 2H, Pht- α), 7.85-7.90 (m, 2H, Pht- β); ¹³C NMR (CDCl₃) 21.0 and 23.1 (C- δ), 24.1 (C-4), 24.6 (C- γ), 38.6 (C- β), 49.1 (C-3), 52.4 (CO₂CH₃), 53.1 (C- α), 105.3 (C-5), 123.4 (Pht- α), 126.5 (C-6), 131.9 (Pht-*quaternary*), 134.0 (Pht- β), 165.7 (Pht-CO), 167.5

10183

(C-2), 171.7 (CO_2CH_3). EIMS *m/z* (%) 370 (M⁺, 2), 339 (1), 311 (4), 223 (25), 167 (100). Anal. Calcd for $C_{20}H_{22}N_2O_5$: C, 64.85; H, 5.94; N, 7.56. Found: C, 64.50; H, 5.97; N, 7.36. Enamide (α*S*,3*R*^{*})-21b (lower Rf): IR (CHCl₃) 1779, 1743 and 1686 (CO), 1350 cm⁻¹; ¹H NMR (CDCl₃) 0.88 (d, *J* = 6 Hz, 6H, δ-H), 1.44-1.82 (m, 3H, β-H and γ-H), 2.36-2.50 (m, 1H, 4-H_A), 3.16-3.34 (m, 1H, 4-H_B), 3.74 (s, 3H, CO₂CH₃), 5.03-5.17 (m, 1H, 3-H), 5.24-5.36 (m, 2H, α-H and 5-H), 6.13-6.21 (m, 1H, 6-H), 7.71-7.77 (m, 2H, Pht-α), 7.85-7.88 (m, 2H, Pht-β); ¹³C NMR (CDCl₃) 21.3 and 22.3 (C-δ), 24.4 (C-4), 24.5 (C-γ), 37.9 (C-β), 48.9 (C-3), 52.2 (CO₂CH₃), 54.2 (C-α), 104.6 (C-5), 123.5 (Pht-α), 126.3 (C-6), 132.0 (Pht-*quaternary*), 134.0 (Pht-β), 165.5 (Pht-CO), 167.5 (C-2), 171.2 (*CO*₂CH₃).

 $(\alpha S, 3RS, 5RS)$ -N-[1-(*tert*-Butoxycarbonyl)-2-*tert*-butoxyethyl]-3-phthaloyl-5-methylthiopiperidin-2one (24a-d). To a solution of enamides 19a,b (160 mg, 0.36 mmol), in dry THF (1.6 ml) cooled to -78°C, AIBN (catalytic amount) and CH3SH (4 ml, 280 mmol), were added. The mixture was refluxed for 5 h. Once cooled (0°C), the reaction mixture was filtered and the solvent evaporated. The resulting foam was chromatographed (hexane:AcOEt, 8:2) to yield compounds 24a-d (70%; trans:cis = 3:1) as a white solid. A sample of trans-24 diastereomers was obtained by repeating the chromatography (a and c, higher Rf, 47%): IR (CHCl₃) 1790, 1727 and 1662 (CO) cm⁻¹; ¹H NMR (CDCl₃) 1.18 (s, OC(CH₃)₃), 1.45 (s, CO₂C(CH₃)₃), 2.15-2.20 (m, 4- H_{A}), 2.17 (s, SCH₃), 2.73 (ddd, J = 13, 10 and 6 Hz, 4-H_B), 3.36 (dt, J = 7 and 6 Hz, 5-H), 3.63-3.75 (m, 6- $H_A i \beta - H_A$), 3.86-3.95 (m, $\beta - H_B$), 4.00-4.15 (m, 6- H_B), 4.99 (dd, J = 6 and 3 Hz, $\alpha - H$), 5.17 (dd, J = 10 and 8 Hz, 3-H), 7.62-7.65 (m, Pht-α), 7.75-7.80 (m, Pht-β); ¹³C NMR (CDCl₃) 14.3 (SCH₃), 27.4 (OC(CH₃)₃), 27.9 (CO₂C(CH₃)₃), 30.5 (C-4), 39.3 (C-5), 47.0 (C-3), 50.7 (C-6), 58.3 (C- α), 61.4 (C- β), 73.3 (OC(CH₃)₃), 81.8 (CO₂C(CH₃)₃), 123.3 (Pht-α), 132.0 (Pht-quaternary), 133.9 (Pht-β), 165.4 (Pht-CO), 167.7 (C-2), 168.0 (CO₂C(CH₃)₃). Lactams cis-24 (b and d, from a 3:1 trans:cis mixture): ¹H NMR (CDCl₃) 1.18 (s, OC(CH₃)₃), 1.46 (s, CO₂C(CH₃)₃), 2.17 (s, SCH₃), 2.15-2.20 (m, 4-H_A), 2.75-2.80 (m, 4-H_B), 3.35-3.41 (m, 5-H), 3.63-3.75 (m, 6-H_A i β-H_A), 3.86-3.95 (m, 2H, β-H_B), 4.00-4,15 (m, 6-H_B), 4.92 (dd, J = 6 and 3 Hz, α -H), 5.12 (dd, J = 10 and 8 Hz, 3-H), 7.62-7.65 (m, Pht- α), 7.75-7.80 (m, Pht- β); ¹³C NMR (CDCl₃) 14.3 (SCH₃), 27.4 (OC(CH₃)₃), 27.9 (CO₂C(CH₃)₃), 30.5 (C-4), 39.3 (C-5), 47.1 (C-3), 50.6 (C-6), 58.8 (C-α), 61.4 (C-β), 73.3 (OC(CH₃)₃), 81.7 (CO₂C(CH₃)₃), 123.3 (Pht-α), 132.0 (Pht-quaternary), 133.9 (Pht-β), 166.5 (Pht-CO), 167.7 (C-2), 168.0 (CO₂C(CH₃)₃). EIMS m/z (%) 490 (M⁺, 1), 361 (10), 313 (12), 291 (17), 246 (22), 57 (100). Anal. Calcd for C₂₅H₃₃N₂O₆S: C, 61.35; H, 6.74; N, 5.72; S, 6.54. Found: C, 61.31; H, 7.18; N, 5.59; S, 6.21.

(α S,3S*,5RS)-N-[1-(Methoxycarbonyl)-2-methylpropyl]-3-phthaloyl-5-methylthiopiperidin-2-ones

(25). To a solution of enamide 20a (0.2 g, 0.56 mmol) in dry toluene (5 ml), AIBN (catalytic amount) and CH₃SH (excess) were added at 0°C. The reaction mixture was heated in a sealed tube at 80°C. The reaction was cooled to 0°C, and a tic control was done every 5 h. If there was any starting enamide left, some more AIBN (catalytic amount) and more CH₃SH was added, and the reaction was continued (80°C). The excess CH₃SH and the toluene were evaporated to obtain a yellow oil that was flash chromatographed (hexane:AcOEt, 1:1) to yield a 3:1 mixture of methylthio derivatives *trans*-25a and *cis*-25b (80%) Lactam *trans*-25a (higher Rf, 60%): [α]_D = -107.2 (c = 1.00, CHCl₃). ¹H NMR (CDCl₃) 0.95 and 0.96 (2d, *J* = 7 Hz, 3H each, γ -H), 2.15-2.20 (m, 2H, 4-H_A and β -H), 2.18 (s, 3H, SCH₃), 2.69 (ddd, *J* = 13, 10 and 4 Hz, 1H, 4-

H_B), 3.29-3.33 (m, 1H, 5-H), 3.35 (dd, J = 12 and 3 Hz, 1H, 6-H_A), 3.74 (s, 3H, CO₂CH₃), 3.86 (dd, J = 12 and 3 Hz, 1H, 6-H_B), 4.80 (d, J = 10 Hz, 1H, α-H), 5.08 (dd, J = 10 and 8 Hz, 1H, 3-H), 7.70-7.77 (m, 2H, Pht-α), 7.82-7.88 (m, 2H, Pht-β); ¹³C NMR (CDCl₃) 14.6 (SCH₃), 18.9 and 19.6 (C-γ), 27.6 (C-β), 30.5 (C-4), 39.5 (C-5), 47.0 (C-3), 47.9 (C-6), 51.9 (CO₂CH₃), 61.7 (C-α), 123.4 (Pht-α), 132.0 (Pht-*quaternary*), 134.0 (Pht-β), 165.4 (Pht-CO), 167.6 (C-2), 171.0 (CO_2 CH₃). EIMS *m/z* (%) 404 (M⁺, 5), 357 (96), 345 (60), 297 (100), 246 (77). Lactam *cis*-25b (lower Rf, 20%) IR (CHCl₃) 1776, 1717 and 1681 (CO) cm⁻¹; ¹H NMR (CDCl₃) (from a diastereomeric mixture **a:b**, 3:1) 0.95 and 0.96 (2d, J = 7 Hz, 3H each, γ-H), 2.15-2.25 (m, 2H, 4-H_A and β-H), 2.16 (s, 3H, SCH₃), 2.48 (q, J = 11 Hz, 1H, 4-H_B), 3.07 (tt, J = 11 and 3 Hz, 1H, 5-H), 3.33-3.45 (m, 1H, 6-H_A), 3.58 (dd, J = 12 and 3 Hz, 1H, 6-H_B), 3.78 (s, 3H, CO₂CH₃), 4.70 (d, J = 10 Hz, 1H, α-H), 5.08 (dd, J = 10 and 8 Hz, 1H, 3-H), 7.70-7.79 (m, 2H, Pht-α), 7.80-7.90 (m, 2H, Pht-β); ¹³C NMR (CDCl₃) 14.0 (SCH₃), 22.9 and 23.6 (C-γ), 28.8 (C-β), 30.3 (C-4), 38.6 (C-5), 46.9 (C-3), 47.9 (C-6), 51.9 (CO₂CH₃), 61.6 (C-α), 123.4 (Pht-α), 132.0 (Pht-*quaternary*), 134.0 (Pht-β), 165.4 (Pht-CO), 167.6 (C-2), 171.0 (CO_2CH_3).

(aS,3S*,5RS)-N-[1-(Methoxycarbonyl)-3-methylbutyl]-3-phthaloyl-5-methylthiopiperidin-2-ones

(26). Operating as above, from enamide 21a (0.2 g, 0.54 mmol), CH₃SH (excess), and AIBN (catalytic amount), in dry toluene (5 ml), compounds 26 were obtained (95%), as a 3:1 C-5 epimeric mixture of lactams trans-26a and cis-26b. Pure samples were obtained by repeating the chromatography (hexane: AcOEt, 6:4). Lactam trans-26a (lower Rf, major isomer, 61%): [α]_D = -68.0 (c = 0.3, CHCl₃). IR (CHCl₃) 1774, 1720, 1717 and 1656 (CO) cm⁻¹; ¹H NMR (CDCl₃) 0.84 and 0.89 (2d, J = 7 Hz, 3H each, δ -H), 1.50-1.70 (m, 3H, β -H and γ -H), 2.13 (s, 3H, SCH₃), 2.12-2.24 (m, 1H, 4-H_A), 2.73 (ddd, J = 13, 10 and 5 Hz, 1H, 4-H_B), 3.26 (ddd, J = 12, 6 and 1 Hz, 1H, 6-H_A), 3.35-3.45 (m, 1H, 5-H), 3.69 (s, 3H, CO₂CH₃), 3.80 (dd, J = 12 and 4 Hz, 1H, 6-H_B), 5.08 (dd, J = 10 and 8 Hz, 1H, 3-H), 5.26 (t, J = 8 Hz, 1H, α -H), 7.65-7.70 (m, 2H, Pht- α), 7.75-7.80 (m, 2H, Pht-β); ¹³C NMR (CDCl₃) 14.6 (SCH₃), 21.4 and 23.1 (C-δ), 24.9 (C-γ), 30.7 (C-4), 37.4 (C-β), 39.6 (C-5), 46.9 (C-3), 47.6 (C-6), 52.4 (CO₂CH₃), 54.2 (C-α), 123.4 (Pht-α), 132.0 (Pht-quaternary), 134.0 (Phtβ), 165.4 (Pht-CO), 167.7 (C-2), 171.7 (CO₂CH₃). EIMS m/z (%) 418 (M⁺, 4), 371 (M⁺-SMe, 93), 359 (43), 311 (94), 246 (100), 167 (36). Lactam *cis*-26b (higher Rf, minor isomer, 20%): [α]_D = -58.9 (c = 0.3, CHCl₃). ¹H NMR (CDCl₃) 0.91 and 0.97 (2d, J = 7 Hz, 3H each, δ-H), 1.50-1.60 (m, 1H, γ-H), 1.63-1.86 (m, 2H, β-H), 2.19 (s, 3H, SCH₃), 2.28-2.42 (m, 1H, 4-H_A), 3.01-3.12 (m, 1H, 4-H_B), 3.46-3.52 (m, 2H, 6-H_A and 5-H), 3.66-3.76 (m, 1H, 6-H_B), 3.77 (s, 3H, CO₂CH₃), 4.88 (dd, J = 12 and 6 Hz, 1H, 3-H), 5.14 (dd, J = 10 and 4 Hz, 1H, α-H), 7.62-7.65 (m, 2H, Pht-α), 7.75-7.80 (m, 2H, Pht-β); ¹³C NMR (CDCl₃) 14.2 (SCH₃), 21.3 and 23.3 (C-δ), 24.7 (C-γ), 32.6 (C-4), 36.6 (C-β), 38.6 (C-5), 48.6 (C-3), 49.2 (C-6), 52.3 (CO₂CH₃), 55.0 (Cα), 123.4 (Pht-α), 132.0 (Pht-quaternary), 134.0 (Pht-β), 165.4 (Pht-CO), 167.5 (C-2), 170.0 (CO₂CH₃). EIMS m/z (%) 418 (M+, 4), 371 (M+-SMe, 22), 359 (93), 311 (53), 246 (42), 167 (100).

ACKNOWLEDGEMENTS

Support for this research has been provided by the CIRIT (Generalitat de Catalunya) through grants QFN95-4703, 1995SGR-00429, and 1997SGR-00075, and by the DGICYT (Ministerio de Educación y Cultura, Spain) through grants PB97-0976 and 2FD97-0293. We also thank the Ministerio de Educación y Cultura for a fellowship given to M.A. Estiarte.

REFERENCES AND NOTES

- To whom the correspondence should be addressed. Phone: 34-93-4024540. FAX: 34-93-4021896. E-mail: adiez@farmacia.far.ub.es
- a. For a general review, see: Giannis, A.; Kolter, T. Angew. Chem. Int. Ed. Engl., 1993, 32, 1244-1267. b. Luthman, K.; Hacksell, U. "Peptides and Peptidomimetics" in "A Textbook of Drug Design and Development", 2nd Ed. Krogsgaard-Larsen, P. and Madsen U. Eds., Overseas Publishers Association. Amsterdam, 1996.
- a. Freidinger, R.M.; Perlow, D.S.; Veber, D.F. J. Org. Chem., 1982, 47, 104-109. b. For a phenylalanine analogue, see: Wyss, C.; Batra, R.; Lehmann, C.; Sauer, S.; Giese, B. Angew. Chem. Int. Ed. Engl., 1996, 35, 2529-2531. c. For tryptophan analogues, see: Rodríguez, R.; Diez, A.; Rubiralta, M.; Giralt, E. Heterocycles, 1996, 43, 513-517.
- 3. For a review, see: Fernández-Checa, J.C.; Yi, J-R.; García-Ruiz, C.; Ookhtens, M.; Kaplowitz, N. Semin. Liver Dis., 1996, 16, 147-158.
- 4. Rodríguez, R.; Estiarte, M.A.; Diez, A.; Rubiralta, M. Tetrahedron, 1996, 52, 7727-7736.
- The biological assays of the {Met-aa} compounds as potential inhibitors of the hepatic transport of glutathione are performed by Dr. J.-C. Fernández-Checa and his group at the Hospital Clínic Provincial de Barcelona.
- a. Since valine tends to generate β-sheets, and leucine α-helixes,^{6b} the pseudopeptides will also be evaluated as conformation inducers. b. Pons, M.; Royo, M. "Diseño de estructuras peptidicas" in "Péptidos en biología y biomedicina", Andreu, D.; Rivas, L. Eds., CSIC. Madrid, 1997. Pp. 198-200, and references cited therein.
- a. Romo, D.; Meyers, A.I. *Tetrahedron*, **1991**, *47*, 9503-9569. b. Micouin, L.; Varea, T.; Riche, C.; Chiaroni,
 A.; Quirion, J.-C.; Husson, H.-P. *Tetrahedron Lett.*, **1994**, *35*, 2529-2532.
- a. The preparation of anhydride 5 has been reported: Vassel, B. U.S. Patent 2806859, Sept. 17, 1957; *Chem. Abstr.*, 1957, 3855a. However, it was not isolated nor characterized. We prepared anhydride 5 b y allyloxycarbonylation^{8b} of glutamic acid, followed by ring closure with Et₃N and MsCl.^{8c} Anhydride 5 was observed to be unstable, and was used without purification. b. For a general method for allyloxycarbonylation of amino acids, see: Fox, S.W.; Wax, H. *J. Am. Chem. Soc.*, 1950, *72*, 5087-5090. c. Chandrasekaran, S.; Nangia, A. *J. Chem. Res.*, 1984, 100.
- 9. Speckamp, N.; Hiemstra, H. Tetrahedron, 1985, 41, 4367-4416.
- The following assays were performed: a. TFA, see reference 4. b. *p*-TsOH, see: Utermoehlen, C.M.; Singh, M.; Lehr, R.E. *J. Org. Chem.*, **1987**, *52*, 5574-5582. c. MsCl/Et₃N and DBU, see: Williams, R.M.; Maruyama, L.K. *J. Org. Chem.*, **1987**, *52*, 4044-4047. d. POCl₃/pyridine, see: Menha, G.; Murthy, A.N.; Reddy, D.S.; Reddy, A.V. *J. Am. Chem. Soc.*, **1986**, *108*, 3443-3452. d. NH₄Br, see: Nyberg, K. *Synthesis*, **1976**, 545-546.
- 11. Fukuyama, T.; Liu, G.; Linton, S.D.; Lin, S.C.; Nishino, H. Tetrahedron, 1993, 34, 2577-2580.
- 12. Nagai, U.; Sato, K. Tetrahedron Lett., 1985, 26, 647-650.
- 13. Fukuyama, T.; Lin, S.C.; Li, L. J. Am. Chem. Soc., 1990, 112, 7050-7051.

14. On one occasion enamides 19 (16%) were obtained together with oxazolidones 22 (29 %) and 23 (12 %), resulting from the loss of the *tert*-butyl protecting groups in the acid medium and cyclisation with the intermediate acyliminium salt. This explains the moderate yield of the reaction. In the absence of *p*-TsOH the reaction did not improve.



- 15. The absolute stereochemistry of the 3-position has not been determined. However, for the sake of clarity, we have called isomers **a** and **b** 3S* and isomers **c** and **d** 3R*.
- 16. This assignment was consistent with the major isomer observed in the phenylglycinol series,⁴ whose stereochemistry has now been confirmed by X-ray crystallography.