Dibutyltin Dibromide-catalyzed Trimethylsilylcyanation of Aldehydes and Imines

Akira Yanagisawa,* Takuya Matsumoto, Takayoshi Arai, and Kazuhiro Yoshida Department of Chemistry, Graduate School of Science, Chiba University, Chiba 263-8522

(Received January 28, 2009; CL-090094; E-mail: ayanagi@faculty.chiba-u.jp)

Dibutyltin dibromide efficiently catalyzes addition of trimethylsilyl cyanide to aldehydes and aldimines. Various trimethylsilylated cyanohydrins and α -aminonitriles are obtained in moderate to high yield by this method.

Organotin(IV) reagents such as trialkyltin enolates and allylic trialkyltins are known to have moderate nucleophilicity and have been often utilized for carbon-carbon bond forming reactions; however, their use in organic synthesis is somewhat limited because of their toxicity.¹ To overcome this difficulty, we attempted to develop the above-mentioned reactions promoted by a catalytic amount of organotin(IV) compounds and as a result, we found an aldol reaction of alkenyl trichloroacetates catalyzed by dibutyltin dimethoxide, which is efficiently recycled with the assistance of methanol.² By taking advantage of this catalytic system, the Mannich-type reaction of aldimines³ and the Michael addition to *p*-benzoquinone⁴ have been also achieved. As for these reactions, noteworthy is the fact that organotin(IV) enolates are generated catalytically and immediately undergo addition to the corresponding electrophiles. In addition, we have found a dibutyltin oxide-catalyzed allyl-transfer reaction from tertiary homoallylic alcohols to aldehydes, in which a catalytic amount of an allylic tin compound is formed in situ.⁵ Our interest in the catalytic ability of these organotin(IV) compounds has led us to undertake a further investigation of related tin species. Here, we wish to describe a cyanation of aldehydes and imines with trimethylsilyl cyanide catalyzed by dibutyltin dibromide (Scheme 1).

Cyanohydrins are versatile synthetic intermediates for diverse natural products, which are further transformable into α hydroxycarboxylic acids or α -hydroxyketones.⁶ In order to construct the cyanohydrin moiety efficiently, various metal catalysts have been applied to silylcyanation of aldehydes.⁷ As for organotin(IV) compounds, tributyltin cyanide has been found to catalyze addition of triethylsilyl cyanide to aldehydes⁸ and relevant acylcyanation.9 In 1996, Whitesell and Apodaca have reported that dibutyltin dichloride is an effective catalyst for trimethylsilylcyanation of aldehydes and ketones in the absence of solvent.¹⁰ They measured a 1:1 mixture of Bu₂SnCl₂ and Me₃SiCN in CDCl₃ by ¹HNMR at room temperature and detected Me₃SiCl (ca. 5%) which suggests a tin cyanide as a catalytically active species. This result prompted us to examine Bu_2SnBr_2 (1), which is anticipated to show higher catalytic activity than that of Bu₂SnCl₂.¹¹ So, we carried out a reaction of benzaldehyde (2a) with Me₃SiCN (3) in the presence of 10 mol % Bu₂SnBr₂ (1) in CH₂Cl₂ under more diluted conditions than Whitesell's condi-

Scheme 1. Bu₂SnBr₂-catalyzed cyanation of aldehydes and imines.

Table 1. Screening of tin catalysts in trimethylsilylcyanation of benzaldehyde (2a) with cyanotrimethylsilane $(3)^a$

O II		tin catalyst (0.1 equiv)	OSiMe ₃
Ph H 2a	3 (2 equiv)	CH ₂ Cl ₂ , r.t., 24 h	Ph CN 4a
Entry	Tin catalyst		Yield/% ^b
1	Bu_2SnBr_2 (1)		52
2	Bu_2SnCl_2		14
3	$Bu_2Sn(OMe)_2$		3
4	Bu ₂ SnO		3
5	Bu ₃ SnBr		<1

^aThe reaction was carried out employing cyanotrimethylsilane (**3**, 2 equiv), benzaldehyde (**2a**, 1 equiv), and tin catalyst (0.1 equiv) in CH_2Cl_2 at room temperature for 24 h. ^bIsolated yield.

Table 2. Solvent effect on trimethylsilylcyanation of benzaldehyde (2a) with cyanotrimethylsilane (3) catalyzed by Bu_2SnBr_2 (1)^a

		Bu ₂ SnBr ₂ (1, 0.1 equiv)	OSiMe ₃
Ph H 2a	Ph H 3 (2 equiv) 2a	solvent, r.t., 24 h	Ph CN 4a
Entry		Solvent	Yield/% ^b
1		CH ₂ Cl ₂	52
2		THF	78
3°		THF	<1
4		MeCN	
5		DMF	<1
6	MeOH		<1
7		Toluene	<1

^aThe reaction was carried out employing cyanotrimethylsilane (**3**, 2 equiv), benzaldehyde (**2a**, 1 equiv), and dibutyltin dibromide (**1**, 0.1 equiv) in a specified solvent at room temperature for 24 h. ^bIsolated yield. ^cBu₂SnCl₂ was used as a catalyst instead of Bu₂SnBr₂ (**1**).

tions and compared its catalytic activity with other organotin(IV) reagents (Table 1). As a result, Bu_2SnBr_2 (1) afforded the desired trimethylsilylated cyanohydrin **4a** in 52% yield in the reaction at room temperature for 24 h (Entry 1). In contrast, Bu_2SnCl_2 indicated lower reactivity under similar reaction conditions (Entry 2). $Bu_2Sn(OMe)_2$ and Bu_2SnO , which were effective catalysts for the aforementioned aldol reaction² and allyl-transfer reaction,⁵ had almost no catalytic activity in the present cyanation (Entries 3 and 4).

In order to obtain a more satisfactory yield, we then investigated solvent effects (Table 2). Among the solvents tested, THF was found to furnish a higher yield than that given in CH_2Cl_2 (Entries 1 and 2), while Bu_2SnCl_2 did not promote the cyanation at all in the former solvent (Entry 3). Acetonitrile was also a promising solvent (Entry 4), however, use of DMF, MeOH, and toluene resulted in totally unacceptable yields (Entries 5–7).

Table 3. Bu_2SnBr_2 (1)-catalyzed trimethylsilylcyanation of various aldehydes **2** with trimethylsilyl cyanide (**3**)^a

0 + R H 3		Bu ₂ SnBr ₂ (1, 0.1 equiv)	OSiMe ₃
	3 (2 equiv)	THF, r.t., 24 h	R CN 4
Entry	Aldehyde		Yield/% ^b
1	PhCHO (2a)		78
2	p-MeOC ₆ H ₄ CHO (2b)		65
3	<i>o</i> -MeOC ₆ H ₄ CHO (2c)		70
4	2-C ₁₀ H ₇ CHO (2d)		74
5	(E)-PhCH=CHCHO $(2e)$		68
6	$Ph(CH_2)_2CHO(2f)$		74
7	<i>c</i> -C ₆ H ₁₁ CHO (2 g)		68

^aThe reaction was carried out using cyanotrimethylsilane (**3**, 2 equiv), aldehyde (**2**, 1 equiv), and dibutyltin dibromide (**1**, 0.1 equiv) in THF at room temperature for 24 h. ^bIsolated yield.

Table 4. Bu_2SnBr_2 (1)-catalyzed cyanation of various aldimines 5 with trimethylsilyl cyanide (3)^a

N ^F		Bu ₂ SnBr ₂ (1 , 0.1 equiv)	HN ^{R²}
R ¹ 人 5	3 (2 equiv)	THF, r.t., 24 h	R ¹ CN 6
Entry	\mathbb{R}^1	\mathbb{R}^2	Yield/% ^b
1	Ph	Ph (5a)	99
2	Ph	p-BrC ₆ H ₄ (5b)	>99
3	Ph	p-MeOC ₆ H ₄ (5c)	>99
4	p-CF ₃ C ₆ H ₄	Ph (5d)	96
5	p-MeOC ₆ H ₄	Ph (5e)	>99
6	o-MeOC ₆ H ₄	Ph (5f)	95
7	<i>p</i> -MeOC ₆ H ₄	p-BrC ₆ H ₄ (5g)	95
8	1-Naphthyl	Ph (5h)	98
9	2-Furyl	Ph (5i)	97

^aThe reaction was carried out using cyanotrimethylsilane (**3**, 2 equiv), aldimine (**5**, 1 equiv), and dibutyltin dibromide (**1**, 0.1 equiv) in THF at room temperature for 24 h. ^bIsolated yield.

Under thus optimized reaction conditions, we finally performed cyanation of various aldehydes 2 with cyanotrimethylsilane (3) catalyzed by Bu₂SnBr₂ (1) in THF. The results are shown in Table 3. Not only aromatic aldehydes 2a–2d but α,β -unsaturated aldehyde 2e and aliphatic aldehydes 2f and 2g were also efficiently silylcyanated and the corresponding α trimethylsiloxynitriles 4 were obtained in good yield. In the case of cinnamaldehyde (2e), only a 1,2-adduct was observed (Entry 5).

We further attempted addition of cyanotrimethylsilane (3) to aldimines 5 using Bu₂SnBr₂ (1) as a catalyst. The Strecker-type reaction is a beneficial route to α -amino acids or 1,2-diamines via α -aminonitriles.^{7c,12} First, we tested the optimized reaction conditions for the above-mentioned trimethylsilylcyanation of aldehydes. When *N*-benzylideneaniline (5a) was treated with two equivalents of Me₃SiCN (3) in the presence of 0.1 equiv of Bu₂SnBr₂ (1) in THF at room temperature for 24 h, the targeted α -aminonitrile 6a was obtained in 99% yield (Table 4, Entry 1). Thus, we carried out the cyanation of various aldimines 5 under the reaction conditions and Table 4 indicates representative examples. As a result, in addition to electron-deficient imines 5b and 5d, electron-rich imines 5c, 5e, and 5f also gave



Figure 1. A probable catalytic cycle.

the corresponding cyanated product **6** almost quantitatively (Entries 2–6).

A possible catalytic mechanism for the present cyanation of aldehydes is shown in Figure 1.¹⁰ First, Bu_2SnBr_2 (1) reacts with cyanotrimethylsilane (3) to give Me₃SiBr (7) and dibutyltin bromide cyanide (8). Subsequently, the tin cyanide 8 is allowed to add to aldehyde 2, yielding tin alkoxide of cyanohydrin 9. Finally, silylation of the tin alkoxide 9 with 3 completes the catalytic cycle by regenerating the tin bromide cyanide 8 and furnishing the desired product 4. An alternative catalytic mechanism in which dibutyltin dicyanide is recycled in place of $Bu_2Sn(CN)Br$ (8) is also possible.

In summary, we have achieved a Bu₂SnBr₂-catalyzed addition reaction of Me₃SiCN to aldehydes and imines. The tin catalyst is efficiently regenerated via Bu₂Sn(CN)Br under the mild reaction conditions and various trimethylsilylated cyanohydrins and α -aminonitriles can be obtained in satisfactory yield. The procedure is operationally simple and can be performed using readily available chemicals.

This work was supported by a Grant-in-Aid for Scientific Research on Priority Areas "Advanced Molecular Transformations of Carbon Resources" from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

References and Notes

- For reviews, see: a) A. G. Davies, Organotin Chemistry, VCH, Weinheim, 1997, p. 185. b) A. Orita, J. Otera, in Main Group Metals in Organic Synthesis, ed. by H. Yamamoto, K. Oshima, Wiley-VCH, Weinheim, 2004, Vol. 2, p. 621.
- 2 A. Yanagisawa, T. Sekiguchi, Tetrahedron Lett. 2003, 44, 7163.
- 3 A. Yanagisawa, H. Saito, M. Harada, T. Arai, *Adv. Synth. Catal.* 2005, 347, 1517.
- 4 A. Yanagisawa, Y. Izumi, T. Arai, Chem. Lett. 2008, 37, 1092.
- 5 A. Yanagisawa, T. Aoki, T. Arai, Synlett 2006, 2071.
- 6 For reviews, see: a) M. North, Synlett 1993, 807. b) F. Effenberger, Angew. Chem., Int. Ed. Engl. 1994, 33, 1555. c) R. J. H. Gregory, Chem. Rev. 1999, 99, 3649. d) M. North, Tetrahedron: Asymmetry 2003, 14, 147.
- For reviews, see: a) M. Shibasaki, M. Kanai, K. Funabashi, *Chem. Commun.* 2002, 1989. b) J.-M. Brunel, I. P. Holmes, *Angew. Chem., Int. Ed.* 2004, 43, 2752. c) M. Kanai, N. Kato, E. Ichikawa, M. Shibasaki, *Synlett* 2005, 1491. d) T. R. J. Achard, L. A. Clutterbuck, M. North, *Synlett* 2005, 1828.
- 8 M. Scholl, G. C. Fu, J. Org. Chem. 1994, 59, 7178.
- 9 M. Scholl, C.-K. Lim, G. C. Fu, J. Org. Chem. 1995, 60, 6229.
- 10 J. K. Whitesell, R. Apodaca, Tetrahedron Lett. 1996, 37, 2525.
- Otera, Orita, and co-workers have reported that bis(pentafluorophenyl)tin dibromide catalyzes trimethylsilylcyanation of an aldehyde, see: J. Chen, K. Sakamoto, A. Orita, J. Otera, *Synlett* **1996**, 877.
- For reviews, see: a) S. Kobayashi, H. Ishitani, *Chem. Rev.* 1999, 99, 1069.
 b) H. Gröger, *Chem. Rev.* 2003, 103, 2795.