CHEMISTRY LETTERS, pp. 1059-1062, 1986.

© 1986 The Chemical Society of Japan

Catalytic Action of Metal Salts in the Amination of 5,8-Quinolinediones

Katsuhira YOSHIDA,^{*} Mayumi YAMAMOTO, and Miwa ISHIGURO Department of Chemistry, Faculty of Science, Kochi University, Akebono-cho, Kochi 780

The catalytic effects of some metal ions on the amination of 5,8-quinolinedione have been investigated. The reactions were specifically promoted by nickel(II) ion, and 6-aminated products were exclusively obtained. The metal chelate complex was proposed as an activated intermediate.

The reactivities of organic ligands coordinated to metal ions are often quite different from those of the free ligands. Recently, we have reported the novel direct alkylamination of d-aminoanthraquinones promoted by metal chelate formation.¹⁾ We now report a catalytic action of metal salt in the amination of 5,8-quinoline-dione(1) and the amine displacement of 6-piperidino-5,8-quinolinedione(2).



The reaction of 2 with butylamine was found to give 6-butylamino-5,8-quinolinedione(4); the amine exchange reaction proceeded smoothly at 30 °C in the presence of some metal ions, but did not proceed at all without metal ion. The results are summarized in Table 1. Metal chlorides were arranged in the order of the catalytic activity as follows: Ni(II)>Cu(II) > Co(II) > Al(III). The activity of the Ni(II) ion was particularly superior to those of other metal ions. For the presence of

Run	Metal salt	Mole ratio [Metal salt]/[<u>2</u>]	Time h	Recovery/% ^{b)}	Yield/% ^{b)}
1	none	0	50	86	0
2	CuCl ₂	1.0	1	0	73
3	CuCl ₂	0.1	25	trace	43
4	CoCl ₂	0.1	45	4	43
5	NiCl ₂	0.1	2.5	0	83
6	AlCl ₃	0.1	25	trace	7

Table 1. Effect of metal chlorides on the reaction of 2 with butylamine^{a)}

a) Reactant(<u>2</u>)(0.5 g) was stirred in pyridine(15 ml) with butylamine(5 ml) and metal salt at 30 °C.

b) Isolated yield after column chromatography.



Fig. 1. Influence of added Ni(II) ion on the absorption spectra of 2 (----: 2.5x10⁻⁴ M) in chloroform(10 vol% DMF).

Table 2. Influence of added metal ions on the absorption spectra of $\frac{2}{2}$ (2.5 x 10⁻⁴ M) in chloroform(10 vol% DMF)

	•			
Metal salt	Mole ratio [Metal salt]/[<u>2]</u>	<u>λmax</u> nm	<u>Δλmax</u> nm	ΔΑ
none		470 ^{a)}		
NiCl ₂	1.0	503	33	0.80
CuCl ₂	1.0	500	30	0.77
CoCl ₂	1.0	500	30	0.63

a) $\frac{2}{2}$ (2.5x10⁻⁴ M); λ max= 470 nm, A= 1.206.

0.1 mole ratio of nickel(II) chloride, a 83% yield of $\underline{4}$ was obtained after 2.5 h (Run 5). However, the displacement of piperidino group was not observed when 7-piperidino-5,8-quinolinedione($\underline{3}$) and 2-piperidino-1,4-naphthoquinone($\underline{10}$) were used instead of $\underline{2}$. These results suggest that the formation of a metal chelate($\underline{11}$), in which the 1nitrogen atom and 8-carbonyl group of $\underline{2}$, probably increases the electrophilicity of the 6-carbon atom and facilitates the nucleophilic substitution of 6-piperidino group by butylamine. This idea was supported by another experimental results; the absorption spectra of $\underline{2}$ are affected by the addition

of metal ions, causing red shift($\Delta \lambda$ max) of absorption maximum with increase of absorption intensity(Δ A)(Fig. 1). As shown in Table 2, the values of $\Delta \lambda$ max and Δ A for the metal ions are compatible with the order of catalytic activity in the amine exchange reaction. The absorption spectra of <u>10</u>, which did not undergo the amine exchange reaction, were not affected at all by the addition of metal ions.

From the above results, it is expected that the reaction of 1 with amines can be well catalyzed by NiCl₂. The effects of NiCl₂ on the amination of 1 were examined and summarized in Table 3. Isomeric mixtures of the products, 6- and

Run	Amine	Mole ratio [NiCl ₂]/[<u>1</u>]	Time h	Yield ^{d)}
7 ^{a)}	piperidine	none ^{c)}	2.0	<u>2</u> 48 , <u>3</u> 10
8 ^{a)}	11	1.0	1.5	<u>2</u> 75 , <u>3</u> 6
9 ^{b)}	p-N,N-dimethyl- aminoaniline	none ^{c)}	6.0	<u>6</u> 43 , <u>7</u> 37
10 ^{b)}	"	0.5	2.0	<u>6</u> 71 , <u>7</u> 0
11 ^{b)}	p-nitroaniline	none ^{C)}	15.0	no reaction
12 ^{b)}	11	0.1	7.0	<u>8</u> 36 , <u>9</u> 0

Table 3. Effect of added NiCl₂ on the amination of <u>1</u>

a) Reactant(<u>1</u>)(3.14 mmol) was stirred in chloroform(70 ml) at 0 °C with piperidine(9.42 mmol) and NiCl₂ under air atmosphere.

b) Reactant(1)(3.14 mmol) was stirred in 95% ethanol(150 ml) at 30 °C with the substituted aniline(9.42 mmol) and NiCl₂ under air atmosphere.

c) Reactions were conducted without NiCl₂.

d) Isolated yield after column chromatography.

7-amino derivatives of <u>1</u>, were obtained without metal ion(Runs 7 and 9),²⁾ whereas the 6-amino derivatives were preferentially obtained in the presence of NiCl₂(Runs 8 and 10). Atmospheric oxygen plays an important role as an oxidizing agent for the completion of the amination of <u>1</u>; the adducts formed by the Michael type addition of amines to <u>1</u> can be dehydrogenated by both the unchanged <u>1</u> and atmospheric oxygen to give the aminated products. The electrophilicity of 6-position of <u>1</u> would be so strongly increased by the formation of nickel chelate that the 6-amination of <u>1</u> can proceed even in the reaction with p-nitroaniline(Runs 11 and 12).

The present work was partially supported by a Grant-in-Aid for Scientific Research (No. 60750806) from the Ministry of Education, Science and Culture.

References

- K. Yoshida, M. Matsuoka, Y. Yamashita, and T. Kitao, Bull. Chem. Soc. Jpn., <u>53</u>, 2552 (1980); K. Yoshida, M. Matsuoka, Y. Yamashita, S. Nagamori, and T. Kitao, ibid., <u>53</u>, 3725 (1980); K. Yoshida, M. Matsuoka, Y. Yamashita, T. Okugawa, and T. Kitao, Dyes and Pigments, <u>2</u>, 125 (1981); M. Matsuoka, T. Takei, I. Nakamura, K. Yoshida, and T. Kitao, Bull. Chem. Soc. Jpn., <u>54</u>, 2225 (1981); K. Yoshida, K. Nojima, and Y. Yamashita, Mem. Fac. Sci. Kochi Univ., <u>5</u>, 1 (1984).
- 2) The reaction of <u>1</u> with aromatic amines to produce the 6- and 7-amino derivatives is known.³⁾ Y. T. Pratt has found that cerium ion catalyzes the reaction and the 6-arylamino isomers are preferentially obtained.⁴⁾
- R. Long and K. Schofield, J. Chem. Soc., <u>1953</u>, 3919; V. Petrow and B. Sturgeon, ibid., <u>1954</u>, 570.
- 4) Y. T. Pratt, J. Org. Chem., 27, 3905 (1962).

(Received March 28, 1986)