

## AN EXTREMELY MILD AND GENERAL METHOD FOR THE STEREOCONTROLLED CONSTRUCTION OF 1,2-*CIS*-GLYCOSIDIC LINKAGES VIA *S*-GLYCOPYRANOSYL PHOSPHORODIAMIDIMIDOTHIOATES

Shun-ichi Hashimoto, Takeshi Honda, and Shiro Ikegami\*

Faculty of Pharmaceutical Sciences, Teikyo University


Sagamiko, Kanagawa 199-01, Japan

**Summary:** A highly stereocontrolled construction of 1,2-*cis*-glycosidic linkages under extremely mild reaction conditions has been developed by using *S*-glycopyranosyl *N,N,N',N'*-tetramethyl-*N''*-phenylphosphorodiamidimidothioates with a non-participating *O*-2-benzyl group as glycosyl donors in the presence of 2,6-lutidinium *p*-toluenesulfonate and tetrabutylammonium iodide.

The growing significance of glycosides and oligosaccharides as constituents of biologically important compounds such as saponins, cardenolides, antibiotics, glycolipids, and glycoproteins has sparked considerable interest in expeditious methods for the stereocontrolled construction of the glycosidic linkages. Despite the recent advances<sup>1,2</sup> in this field, however, there still exists a continuing demand for appreciable developments in terms of mildness, efficacy, generality, and stereocontrol.

Considering that the leaving group of glycosyl donors is one of the most fundamental parameters responsible for the selectivity and yield of glycosidation reactions, our efforts centered on the development of new glycosyl donors. We were intrigued by the feasibility of using the phosphorus-containing leaving groups,<sup>3</sup> on the prospect that a number of variations of substituents on the phosphorus atom could make "tailor-made glycosyl donors" readily available. We recently developed efficient methods for the stereocontrolled construction of 1,2-*trans*-glycosidic linkages by the device of shelf-stable glycosyl donors incorporating diphenyl phosphate<sup>4</sup> or *P,P*-diphenyl-*N*-(*p*-toluenesulfonyl)phosphinimide<sup>5</sup> as leaving groups. Herein, we wish to report on an extremely mild and general procedure for the stereocontrolled construction of 1,2-*cis*-glycosidic linkages exploiting *S*-glycopyranosyl *N,N,N',N'*-tetramethyl-*N''*-phenylphosphorodiamidimidothioates with a non-participating *O*-2-benzyl group as glycosyl donors.


Construction of 1,2-*cis*-glycosidic linkages in a stereocontrolled manner has been the focus of much recent attention.<sup>6</sup> We envisaged that this challenging problem could be solved by judicious choice of substituents on the phosphorus atom in the leaving groups and promoters in favor of S<sub>N</sub>2-like mechanism, thus focusing on the readily available 1,2-*trans*-related 1-thio-glycopyranose bearing a non-participating benzyl group on *O*-2 as the source of the glycosyl donors. After considerable experimentation, it was found that coupling of *S*-(2,3,4,6-tetra-*O*-benzyl-β-D-glucopyranosyl)-*N,N,N',N'*-tetramethyl-*N''*-phenylphosphorodiamidimidothioate **17** (1.1 equiv.) with suitably protected glycosides **6**, **7**, **9**, and **10** (1.0 equiv.) in toluene in the presence of 2,6-lutidinium *p*-toluenesulfonate (LPTS)<sup>9</sup> (1.15 equiv.) and pulverized molecular sieves 4A at 23 °C proceeded under kinetic control to lead to the preponderant formation of 1,2-*cis*-linked disaccharides (entries 1, 4, 7, 9, and 11 in Table 1). The combination of LPTS as a promoter<sup>10</sup> and toluene



as a solvent proved to be the superior choice for allowing considerable levels of 1,2-*cis*-selectivity as well as extremely mild reaction conditions, with which the acid-labile groups such as epoxy, acetal, or *O*-tert-butyltrimethylsilyl groups are completely comparable.

Although the 1,2-*cis*-glycosidation reactions developed here involved several features,<sup>11</sup> the stereoselectivity was not necessarily satisfactory in view of the requisite levels for current synthetic carbohydrate chemistry. In an effort to enhance the degrees of the stereoselectivity, we reasoned that if this method could be coupled with Lemieux's glycosidation method<sup>6a</sup> via halide ion catalyzed process, the new version would constitute an extremely mild procedure for the highly stereocontrolled construction of 1,2-*cis*-glycosidic linkages. However, the glycosidation of 1 with 6 or 7 in the presence of tetrabutylammonium bromide, the most widely used additive,<sup>6c,j,k</sup> took much longer reaction times than the original method, though 1,2-*cis*-selectivity of up to 96% was attained (entries 2 and 5). In stark contrast, we were pleased to find that this goal could be readily achieved by the use of tetrabutylammonium iodide as an additive. As seen from Table 1, the new version required almost the same reaction times as the original procedure, but afforded even higher levels of 1,2-*cis*-selectivity (entries 3, 6, 8, 10, and 12).<sup>14</sup> It should be noted that this is the first successful use of tetrabutylammonium iodide as an additive. Apart from  $\alpha$ -D-glucosylation, the method was advantageously extended to  $\alpha$ -D-galactosylation and  $\alpha$ -L-fucosylation (entries 18–22), in which toluene- $\text{CH}_2\text{Cl}_2$  (1:1) proved to be the solvent of choice. Furthermore, the mildness and generality of this method was demonstrated by a range of variations possible in glycosyl donors and acceptors.

In conclusion, the present method based on *S*-glycopyranosyl phosphorodiamidimidothioates constitutes an exceptionally mild procedure for the highly stereocontrolled construction of 1,2-*cis*-glycosidic linkages, and thus represent a promising addition to the existing methods. Further extension of this method to the construction of carbohydrate-containing natural products and analogues of medicinal importance are currently in progress in these laboratories.

Table 1. 1,2-*cis*-Glycosidation reaction of *S*-glycopyranosyl phosphorodiamidimidothioates 1-5

| entry           | donor | acceptor        | additive            | time, h | yield, <sup>a</sup> % | $\alpha : \beta$ <sup>b</sup> | $\delta$ $^{13}\text{C}$ c,d | $[\alpha]_{\text{D}}^{23}/^{\circ}(\text{c, CHCl}_3)^{\text{c}}$ | ref. |
|-----------------|-------|-----------------|---------------------|---------|-----------------------|-------------------------------|------------------------------|------------------------------------------------------------------|------|
| 1               | 1     | 6               | —                   | 16      | 85                    | 80:20                         | 97.3                         | +58.2 (1.12)                                                     | 6b   |
| 2               | 1     | 6               | Bu <sub>4</sub> NBr | 48      | 82                    | 94:6                          |                              |                                                                  |      |
| 3               | 1     | 6               | Bu <sub>4</sub> NI  | 16      | 83                    | 91:9                          |                              |                                                                  |      |
| 4               | 1     | 7               | —                   | 55      | 71                    | 86:14                         | 96.6                         | +46.2 (1.20)                                                     | 6b   |
| 5               | 1     | 7               | Bu <sub>4</sub> NBr | 96      | 72                    | 96:4                          |                              |                                                                  |      |
| 6               | 1     | 7               | Bu <sub>4</sub> NI  | 55      | 72                    | 93:7                          |                              |                                                                  |      |
| 7               | 1     | 9               | —                   | 55      | 72                    | 87:13                         | 95.5                         | +50.0 (0.97)                                                     |      |
| 8               | 1     | 9               | Bu <sub>4</sub> NI  | 55      | 76                    | 91:9                          |                              |                                                                  |      |
| 9               | 1     | 10              | —                   | 55      | 75                    | 86:14                         | 96.3                         | +24.7 (0.83)                                                     |      |
| 10              | 1     | 10              | Bu <sub>4</sub> NI  | 55      | 77                    | 93:7                          |                              |                                                                  |      |
| 11              | 1     | 11              | —                   | 55      | 72                    | 87:13                         | 96.9                         | +29.9 (1.31)                                                     | 6e   |
| 12              | 1     | 11              | Bu <sub>4</sub> NI  | 55      | 73                    | 92:8                          |                              |                                                                  |      |
| 13              | 1     | 12              | Bu <sub>4</sub> NI  | 55      | 73                    | 94:6                          | 98.0                         | +44.9 (1.15)                                                     | 6b   |
| 14              | 1     | 14              | Bu <sub>4</sub> NI  | 55      | 80                    | 93:7                          | 98.0                         | +32.9 (1.12)                                                     | 6i   |
| 15              | 1     | 15 <sup>e</sup> | Bu <sub>4</sub> NI  | 55      | 79                    | >99:1                         | 94.8                         | +284.7 (0.55)                                                    |      |
| 16              | 2     | 11              | Bu <sub>4</sub> NI  | 55      | 74                    | >99:1                         | 96.3                         | +18.4 (1.12)                                                     |      |
| 17              | 3     | 11              | Bu <sub>4</sub> NI  | 55      | 72                    | 94:6                          | 96.3                         | +28.0 (1.02)                                                     |      |
| 18 <sup>f</sup> | 4     | 8               | Bu <sub>4</sub> NI  | 55      | 70                    | 93:7                          | 97.5                         | +18.2 (1.50)                                                     | 6k   |
| 19 <sup>f</sup> | 4     | 11              | Bu <sub>4</sub> NI  | 55      | 71                    | 92:8                          | 98.3                         | +15.3 (0.38)                                                     |      |
| 20 <sup>f</sup> | 4     | 13              | Bu <sub>4</sub> NI  | 10      | 73                    | 91:9                          | 97.6                         | +3.2 (1.32)                                                      | 6g   |
| 21 <sup>f</sup> | 5     | 8               | Bu <sub>4</sub> NI  | 55      | 72                    | 93:7                          | 97.7                         | -38.1 (1.12)                                                     | 6k   |
| 22 <sup>f</sup> | 5     | 12              | Bu <sub>4</sub> NI  | 55      | 70                    | 94:6                          | 96.8                         | -95.3 (0.73)                                                     | 6a   |


<sup>a</sup> Isolated total yield. <sup>b</sup> Determined by HPLC and  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ ) analysis. <sup>c</sup> Values for the 1,2-*cis*-linked glycosides or disaccharides purified by flash chromatography on silica gel. <sup>d</sup> Chemical shifts for the anomeric centers newly formed. <sup>e</sup> C-7 hydroxy group was glucosylated. <sup>f</sup> Performed in toluene- $\text{CH}_2\text{Cl}_2$  (1:1).

Acknowledgement: We are grateful to Misses Y. Ikenoya and H. Ishido for spectroscopic measurements. We also thank Kyowa Hakko Kogyo Co. Ltd. for a generous gift of adriamycin. This research was supported in part by grants from the Japan Research Foundation for Optically Active Compounds, the Japan Science Society, and the Ministry of Education, Science and Culture of Japan.

## References and Notes

- For recent reviews, see: (a) H. Paulsen, *Angew. Chem. Int. Ed. Engl.*, **21**, 155 (1982); (b) R. R. Schmidt, *ibid.*, **25**, 212 (1986); (c) H. Kunz, *ibid.*, **26**, 294 (1987).
- For other more recent works, see: (a) F. Dasgupta and P. J. Garegg, *Carbohydr. Res.*, **177**, C13 (1988); (b) Y. Ito and T. Ogawa, *Tetrahedron Lett.*, **29**, 1061 (1988); (c) M. Nishizawa, Y. Kan, and H. Yamada, *ibid.*, **29**, 4597 (1988); (d) D. R. Mootoo, P. Konradsson, U. Udodong, and B. Fraser-Reid, *J. Am. Chem. Soc.*, **110**, 5583 (1988); (e) K. Suzuki, H. Maeta, and T. Matsumoto, *Tetrahedron Lett.*, **30**, 4853 (1989); (f) A. G. M. Barrett, B. C. B. Bezuidenhout, A. F. Gasiccki, A. R. Howell, and M. A. Russell, *J. Am. Chem. Soc.*, **111**, 1392 (1989); (g) D. Kahne, S. Walker, Y. Cheng, and D. V. Engen, *ibid.*, **111**, 6881 (1989); (h) P. Konradsson, D. R. Mootoo, R. E. McDevitt, and B. Fraser-Reid, *J. Chem. Soc., Chem. Commun.*, 270 (1990).
- (a) T. Inazu, H. Hosokawa, and Y. Satoh, *Chem. Lett.*, 297 (1985); (b) M. Michalska and J. Michalski, *Heterocycles*, **28**, 1249 (1989).
- S. Hashimoto, T. Honda, and S. Ikegami, *J. Chem. Soc., Chem. Commun.*, 685 (1989).
- S. Hashimoto, T. Honda, and S. Ikegami, *Heterocycles*, **30**, 775 (1990).
- (a) R. U. Lemieux, K. B. Hendriks, R. V. Stick, and K. James, *J. Am. Chem. Soc.*, **97**, 4056 (1975); (b) J.-R. Pouigny, J.-C. Jacquinot, M. Nassr, D. Duchet, M.-L. Milat, and P. Sinaÿ, *ibid.*, **99**, 6762 (1977); (c) J. Leroux and A. S. Perlin, *Carbohydr. Res.*, **67**, 163 (1978); (d) T. Mukaiyama, T. Nakatsuka, and S. Shoda, *Chem. Lett.*, 487 (1979); (e) R. R. Schmidt and J. Michel, *Angew. Chem. Int. Ed. Engl.*, **19**, 731 (1980); (f) T. Mukaiyama, Y. Murai, and S. Shoda, *Chem. Lett.*, 431 (1981); (g) P. J. Garegg, C. Ortega, and B. Samuelsson, *Acta Chem. Scand., Ser. B*, **35**, 631 (1981); (h) S. Koto, N. Morishima, M. Owa, and S. Zen, *Carbohydr. Res.*, **130**, 73 (1984); (i) S. Hashimoto, M. Hayashi, and R. Noyori, *Tetrahedron Lett.*, **25**, 1379 (1984); (j) F. Andersson, P. Flüggedi, P. J. Garegg, and M. Nashed, *ibid.*, **27**, 3919 (1986); (k) S. Sato, M. Mori, Y. Ito, and T. Ogawa, *Carbohydr. Res.*, **155**, C6 (1986); (l) C. P. Fei and T. H. Chan, *Tetrahedron Lett.*, **28**, 849 (1987); (m) W. Szeja, *Synthesis*, 223 (1988); (n) G. V. Reddy, V. R. Kulkarni, and H. B. Mercyala, *Tetrahedron Lett.*, **30**, 4283 (1989); (o) N. K. Kochetkov, E. M. Klimov, and N. N. Malysheva, *ibid.*, **30**, 5459 (1989); (p) G. H. Veeneman and J. H. van Boom, *ibid.*, **31**, 275 (1990).
- The 1,2-*trans*-related S-glycopyranosyl phosphorodiamidimidothioates 1-5 were prepared from the corresponding 2,3,4,6-tetra-O- or 2,3,4-tri-O-protected glycopyranoses in 56-63% overall yields by the following sequence: (1) MsCl, 2,6-lutidine, CH<sub>2</sub>Cl<sub>2</sub>, 23 °C, 10 h; (2) KSC(=S)OEt, benzene-EtOH (2:1), 23 °C, 8 h; (3) NaOMe, MeOH, 40 °C, 1 h; (4) *n*-BuLi (1.1 equiv.), THF, -78 °C, 0.5 h; PhN=P(NMe<sub>2</sub>)<sub>2</sub>Cl<sup>8</sup> (1.1 equiv.), HMPA, 23 °C, 3 h. These donors can be stored without decomposition in the freezer (at -20 °C) for several months.
- V. A. Gilyarov, R. V. Kudryavtsev, and M. I. Kabachnik, *J. Gen. Chem. USSR (Engl. Transl.)*, **38**, 351 (1968).
- T. J. Nitz and L. A. Paquette, *Tetrahedron Lett.*, **25**, 3047 (1984).
- The corresponding 2,4,6-collidinium salt (CPTS) was almost equally effective, but the use of the pyridinium salt (PPTS) met with less success.
- Based on the results summarized in the table (*vide infra*), the stereochemical outcome of the glycosidations can be explained as follows: The glycosidations of 1 and its  $\alpha$ -anomer 1' are presumed to proceed via  $\beta$ - and  $\alpha$ -sulfonium salts formed by protonation of the nitrogen atom with LPTS, which are in equilibrium via glycopyranosyl tosylate<sup>12</sup> in favor of the  $\beta$ - salt due to the reverse anomeric effect,<sup>13</sup> followed by the back side attack with acceptor alcohols on these intermediates. In less polar toluene, the nucleophilic displacement by cyclohexylmethanol is competitive with the anomeric equilibration of the salts, and thus much higher 1,2-*cis*-selectivity is obtained from 1 than from 1'. On the other hand, the attack of cyclohexanol is slow compared to the equilibration, 1 exhibiting slightly higher selectivity than 1'. In more polar dichloromethane, since the equilibrium is reached prior to the attack by alcohols, the almost unchanged ratios are attained regardless of the anomeric configuration of the donors and alcohols.

| entry | donor | ROH                                                         | solvent                         | time, h | yield, % | $\alpha : \beta$ |
|-------|-------|-------------------------------------------------------------|---------------------------------|---------|----------|------------------|
| 1     | 1     | <i>c</i> -C <sub>6</sub> H <sub>11</sub> CH <sub>2</sub> OH | toluene                         | 10      | 83       | 82 : 18          |
| 2     | 1'    | <i>c</i> -C <sub>6</sub> H <sub>11</sub> CH <sub>2</sub> OH | toluene                         | 10      | 84       | 60 : 40          |
| 3     | 1     | <i>c</i> -C <sub>6</sub> H <sub>11</sub> OH                 | toluene                         | 15      | 85       | 88 : 12          |
| 4     | 1'    | <i>c</i> -C <sub>6</sub> H <sub>11</sub> OH                 | toluene                         | 15      | 83       | 78 : 22          |
| 5     | 1     | <i>c</i> -C <sub>6</sub> H <sub>11</sub> CH <sub>2</sub> OH | CH <sub>2</sub> Cl <sub>2</sub> | 10      | 78       | 77 : 23          |
| 6     | 1'    | <i>c</i> -C <sub>6</sub> H <sub>11</sub> CH <sub>2</sub> OH | CH <sub>2</sub> Cl <sub>2</sub> | 10      | 79       | 76 : 24          |
| 7     | 1     | <i>c</i> -C <sub>6</sub> H <sub>11</sub> OH                 | CH <sub>2</sub> Cl <sub>2</sub> | 15      | 77       | 79 : 21          |
| 8     | 1'    | <i>c</i> -C <sub>6</sub> H <sub>11</sub> OH                 | CH <sub>2</sub> Cl <sub>2</sub> | 15      | 76       | 80 : 20          |



- Glycosidation of 2,3,4,6-tetra-O-benzyl-D-glucopyranosyl tosylate with cyclohexylmethanol or cyclohexanol in toluene was found to result in the  $\alpha : \beta$  ratio of 52:48 and 54:46, respectively.
- A. J. Kirby, *The Anomeric Effect and Related Stereoelectronic Effects at Oxygen*, Springer-Verlag, New York, 1983, pp. 15-17.
- The glycosidation is thought to proceed via the glycopyranosyl iodide generated *in situ*. For the limited use of glycopyranosyl iodide derivatives as glycosyl donors, see: F. K. Kronzer and C. Schuerch, *Carbohydr. Res.*, **34**, 71 (1974).

(Received in Japan 2 May 1990)