

2-Bromo-1-ethyl Pyridinium Tetrafluoroborate (BEP): A Powerful Coupling Reagent for *N*-Methylated Peptide Synthesis

Peng Li and Jie Cheng Xu*


Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, 354 Feng Lin Road, Shanghai 200032, P. R. China

(Received November 4, 1999; CL-990945)

2-Bromo-1-ethyl pyridinium tetrafluoroborate (BEP) was firstly utilized to the synthesis of peptides containing *N*-methyl amino acid residues both in solution and solid phase, and demonstrated high reactivity, low racemization and excellent yields, which were approved by the successful synthesis of a series of oligopeptides and hindered peptide fragments, such as the 8-11 segment of Cyclosporine A and the pentapeptide moiety of Dolastatin 15.

Since the first phosphonium salt BOP¹ was exploited and used to peptide synthesis, many onium type coupling reagents have been designed, synthesized and widely used in peptide chemistry, such as HOBt-derived onium salts: HBTU,² PyBOP,³ HBPYU,⁴ HBPIU,⁵ HBMDU⁶ and the corresponding HOAt-based onium salts⁷: AOP, PyAOP, HATU, HAPYU, HAMDU. To further enhance the coupling efficiency during peptide synthesis, we have designed and synthesized immonium and thiazolium type coupling reagents, such as BOMI,^{8a} BDMP,^{8b} BPMP, AOMP and BEMT^{8c} based upon our previous studies.^{8d-8g} We now introduce pyridinium salt,^{9a,9b} 2-bromo-1-ethyl pyridinium tetrafluoroborate (BEP), of which analogues 1-methyl-2-halopyridinium iodides were shown to be efficient for the syntheses of esters,^{9c} lactones^{9d} and carboxamides,^{9e} into this field and demonstrate its high efficiency in peptide synthesis, especially in the incorporation of hindered *N*-methyl amino acids into peptides, such as cyclosporins,¹⁰ didemins¹¹ and dolastatins.¹²

Comparing to other halogenated coupling reagents PyCIU,^{13a} CIP,^{13b} BOP-Cl^{13c} and PyBroP,^{13d} Pyridinium salt BEP can be more readily synthesized by the *N*-alkylation of 2-bromo pyridine using triethyloxonium tetrafluoroborate with nearly quantitative yield as colorless shelf-stable crystals.¹⁴



HPLC monitoring the model reactions (*Z*-Me-Val-OH + Me-Val-OMe-HCl → *Z*-Me-Val-Me-Val-OMe and *Z*-Gly-Phe-OH + Val-OCH₃·HCl → *Z*-Gly-Phe-Val-OCH₃) showed that reagent BEP behaved higher reactivity and lower racemization during coupling comparing to CBMI, PyBroP, PyCIU and BTFFH et al. (Table 1 and Figure 1).

To further demonstrate the efficiency of BEP during peptide coupling, a series of oligopeptides were synthesized as shown in Table 2. We also synthesized successfully two extensively *N*-alkylated peptide segments. During the synthesis of 8-11 tetrapeptide fragment of cyclosporine A, the *C*-terminal was

Table 1. Comparison of racemization and reactivity of BEP with other coupling reagents^a

Coupling reagent	Yield/% (t = 2 min)	D-isomer content/%
PyBroP	6.10	22.3
PyCIU	5.64	33.2
BTFFH	9.35	25.9
BOP-Cl	5.34	4.13
CMBI	19.7	16.8
BEP	50.8	4.59
BEP + HOAt	75.9	1.42

^aModel reaction: *Z*-Gly-Phe-OH + Val-OCH₃·HCl → *Z*-Gly-Phe-Val-OCH₃. Reaction conditions: T: -10 °C, Base: DIEA, Sol: CH₂Cl₂, substrate ratio: *N*-protected amino acid : amino acid ester hydrochloride : coupling reagent : Base = 1: 1.1: 1.1: 3.2 (mol).




Figure 1. Comparison of reactivity of BEP with other coupling reagents^a.


^aModel reaction: *Z*-MeVal-OH + MeVal-OCH₃·HCl → *Z*-MeVal-MeVal-OCH₃. Reaction conditions: T: 25 °C; Base: DIEA; Solvent: CH₂Cl₂ (0.1 M); Substrate ratio: *N*-protected amino acid : amino acid ester hydrochloride : Coupling reagent : Base = 1: 1.1: 1.1: 3.2 (mol).

Table 2. Preparation of peptides using BEP in solution^a


Peptide ^b	yield/% ^c	mp/°C	[α] _D (conc., solv.)
Fmoc-Nva ^Φ -Sar-OBzl	97.6	38-39	-9.3 (0.3, CHCl ₃)
<i>Z</i> -MeVal ^Φ -MeVal-OMe	95.4	oil	-206 (1, MeOH)
Fmoc-Val ^Φ -MeVal-OCH ₃	88.4	oil	-86.0 (0.3, CHCl ₃)
Boc-Pro ^Φ -Pro-OBzl	90.2	oil	-109.4 (1, CHCl ₃)
<i>Z</i> -Aib ^Φ -Aib-OCH ₃	95.8	108-109	—
Fmoc-MeLeu ^Φ -MeVal-OBzl	91.4	oil	-122 (0.2, CHCl ₃)
Fmoc-MeLeu ^Φ -MeLeu-MeVal-OBzl	48.1	oil	-114.8 (1, MeOH)
Fmoc-D-Ala ^Φ -MeLeu-MeLeu-MeVal-OBzl	94.3	36-37	-145.4 (0.5, CHCl ₃)
Boc-Val-Val-MeVal ^Φ -Pro-OBzl ^d	87.5	89-90	-181 (0.3, CHCl ₃)

^aThe reactions were carried out as for PyBroP. In a typical experimental procedure, DIEA (3.2 equiv.) was added to a cold mixture (-10 °C) of *N*-protected amino acid (1 equiv.), amino acid ester hydrochloride (1.1 equiv.), and BEP (1.1 equiv.) in CH₂Cl₂ (2-4 mL/mmol), stirred for 1 minute cold and for an hour at room temperature; the reaction time should be properly prolonged for the coupling between *N*-methyl amino acids. ^bThe CO-NH bond formed in the peptide is indicated by Φ, all products were confirmed by ¹H NMR, EIMS and other characterizations. ^cIsolated yields based on *N*-protected amino acid. ^dFurther confirmed by COSY and ESI-MS.

intensively protected as benzyl ester to challenge the spontaneous diketopiperazine formation. A 48% yield was still obtained during the synthesis of Fmoc-MeLeu-MeLeu-MeVal-OBzl by sequential deprotection and coupling of Fmoc-MeLeu-MeVal-OBzl with Fmoc-MeLeu-OH. According to the report by Wenger during the synthesis of cyclosporine A using modified mixed pivalic anhydride method, the same desired tripeptide was not obtained at all due to the spontaneous formation of diketopiperazine.¹⁵ To further examine the performance of reagent BEP in SPPS, The hindered 8-11tetrapeptide of CsA and the linear undecapeptide of CsO were also synthesized in solid phase using BEP from Fmoc-MeVal-Wang resin. The purities of the obtained crude peptides were 95% and ~5%¹⁶ respectively.



Cyclosporin A



Dolastatin 15

We also synthesized the hindered pentapeptide moiety of dolastatin 15 which was a pseudopeptide bearing promising antineoplastic activity. The Boc group was used for the N_α -protection, thus a 57.7% overall yield of the protected pentapeptide was obtained via seven coupling steps. No or little N -carboxyanhydride formed during peptide coupling using BEP, although Boc-protected amino acids were more readily converted into NCA than Cbz- or Fmoc-protected derivatives and resulted in relatively low yield using halogenerated reagents.^{13d}

In general, the pyridinium type coupling reagent BEP was shown to be a very efficient coupling reagent for the synthesis of the hindered peptide containing N -methyl amino acid residues with fast reaction speed, low racemization and good yields.

References and Notes

- 1 B. Castro, J. R. Dromoy, G. Evin, and C. Selve, *Tetrahedron Lett.*, **1975**, 1219.
- 2 V. Dourtoglou, J. C. Ziegler, and B. Gross, *Tetrahedron Lett.*, **1978**, 1269.
- 3 J. Coste, D. Le-Nguyen, and B. Castro, *Tetrahedron Lett.*, **31**, 205 (1990).
- 4 a) S. Q. Chen and J. C. Xu, *Tetrahedron Lett.*, **33**, 647 (1992). b) J. Coste, E. Frerot, and P. Jouin, *Tetrahedron Lett.*, **32**, 1967 (1991).
- 5 P. Henklein, M. Beyer mann, M. Bienert, and R. Knorr, "Proceeding of the Twenty-First European Peptide Symposium" ed. by E. Giralt and D. Andreu, ESCOM, Science, Leiden, (1991), p. 67.
- 6 Y. Kiso, Y. Fujiwara, T. Kimura, A. Nishitani, and K. Akaji, *Int. J. Pept. Protein Res.*, **40**, 308 (1992).
- 7 a) L. A. Carpino, A. El-Faham, C. A. Minor, and F. Albericio, *J. Chem. Soc., Chem. Commun.*, **1994**, 201. b) F. Albericio, M. Cases, J. Alsina, S. A. Triolo, L. A. Carpino, and S. A. Kates, *Tetrahedron Lett.*, **38**, 4853 (1997). c) L. A. Carpino, *J. Am. Chem. Soc.*, **115**, 4397 (1993). d) L. A. Carpino and A. El-Faham, *J. Org. Chem.*, **59**, 696 (1994).
- 8 a) P. Li and J. C. Xu, *Tetrahedron Lett.*, **40**, 3605 (1999). b) P. Li and J. C. Xu, *Chem. Lett.*, **1999**, in press. c) P. Li and J. C. Xu, *Tetrahedron Lett.*, **40**, 8301 (1999). d) S. Q. Chen and J. C. Xu, *Tetrahedron Lett.*, **32**, 6711 (1991). e) S. Q. Chen and J. C. Xu, *Tetrahedron Lett.*, **33**, 647 (1992). f) S. Q. Chen and J. C. Xu, *Chin. Chem. Lett.*, **4**, 847 (1993). g) S. Q. Chen and J. C. Xu, *Chin. J. Chem.*, **13**, 175 (1995).
- 9 a) H. Balli and F. Kersting, *Liebigs Ann. Chem.*, **647**, 1 (1961). b) K. Saigo, M. Usui, and K. Kikuchi, *Bull. Chem. Soc. Jpn.*, **50**, 1863 (1977). c) T. Mukaiyama, M. Usui, and E. Shimada, *Chem. Lett.*, **1975**, 1045. d) T. Mukaiyama, M. Usui and K. Saigo, *Chem. Lett.*, **1976**, 49. e) E. Bald, K. Saigo and T. Mukaiyama, *Chem. Lett.*, **1975**, 1163.
- 10 R. M. Wenger, *Angew. Chem., Int. Ed. Engl.*, **24**, 77 (1985).
- 11 P. Jouin, T. Poncet, M-N. Dufour, A. Pantaloni, and B. Castro, *J. Org. Chem.*, **54**, 617 (1989) and references cited therein.
- 12 G. R. Pettit, S. B. Singh, D. L. Herald, P. Lloyd-Williams, D. Kantoci, D. D. Burkett, J. Barkoczy, F. Hogan, and T. R. Wardlaw, *J. Org. Chem.*, **59**, 6287 (1994) and references cited therein.
- 13 a) J. Coste, E. Frerot, and P. Jouin, *Tetrahedron Lett.*, **32**, 1967 (1991). b) K. Akaji, N. Kuriyama, and Y. Kiso, *Tetrahedron Lett.*, **35**, 3315 (1994). c) C. Van der Auwera and M. J. O. Anteunis, *Bull. Soc. Chim. Belg.*, **95**, 203 (1986). d) J. Coste, E. Frerot, and P. Jouin, *J. Org. Chem.*, **59**, 2437 (1994).
- 14 Coupling reagent BEP was prepared according to literature^{9a} in 95.2% yield. mp 103-104 °C. ¹H NMR (300 MHz, d_6 -acetone) δ 1.69 (t, J = 7.3 Hz, 3H), 5.01 (q, J = 7.3 Hz, 2H), 8.24 (m, 1H), 8.53-8.57 (m, 2H), 9.32 (d, J = 6.9 Hz, 1H) ppm. IR (KBr): ν = 3106, 1617, 1571, 1500, 1467, 1296, 1050, 786, 718, 521 cm^{-1} . FAB-MS m/z : 186, 188.
- 15 a) J. M. Humphrey and A. R. Chamberlin, *Chem. Rev.*, **97**, 2243 (1997). b) R. M. Wenger, *Helv. Chim. Acta.*, **66**, 2672 (1983).
- 16 UV analysis of the Fmoc deprotection step indicated that the coupling yield of each step was above 90%. The low purity of the crude product was most likely that the amide bonds between N -methyl amino acids in long peptide fragment were prone to undergo hydrolysis under strong acidic medium.^{7b}