

Full Paper

Taming troublesome Suzuki-Miyaura reactions in water solution of surfactants by the use of lecithin: a step beyond the micellar model

Chiara Ceriani, Erika Ghiglietti, Mauro Sassi, Sara Mattiello, and Luca Beverina

Org. Process Res. Dev., **Just Accepted Manuscript** • DOI: 10.1021/acs.oprd.0c00285 • Publication Date (Web): 04 Sep 2020

Downloaded from pubs.acs.org on September 4, 2020

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.

Taming troublesome Suzuki-Miyaura reactions in water solution of surfactants by the use of lecithin: a step beyond the micellar model

Chiara Ceriani,[†] Erika Ghiglietti,[†] Mauro Sassi,[†] Sara Mattiello,^{*,†} and Luca Beverina^{*,‡}

[†]*Department of Materials Science, University of Milano-Bicocca, Via R. Cozzi, 55, Milano, Italy.*

[‡]*Department of Materials Science, University of Milano-Bicocca and INSTM, Via R. Cozzi, 55, Milano, Italy.*

E-mail: sara.mattiello@unimib.it; luca.beverina@unimib.it

Abstract

The use of water solutions of industrial and designer surfactants enables performing a wide variety of chemical transformation on hydrophobic precursors. Mostly reactions are clean, fast and efficient with vast benefits for overall sustainability. The widespread adoption of such methods is somewhat hampered by a lack of generality in the case of troublesome substrates and scaling up. The common literature solution for such issues is the use of small amounts of organic solvents. We here show that the use of a mixture of L- α -lecithin and Tween 80 is a preferable solution enabling the taming of particularly troublesome reactions, where even the cosolvent approach fails. The strong reduction of all interface tensions characterizing such complex multiphase systems is key to the results achieved. The protocol is general and scalable.

Introduction

The development of improved synthetic methods offering selectivity, high yield and short reaction times while providing moderate or no heating is a cornerstone of modern sustainable chemistry.^{1,2} Unfortunately, the need for organic solvents remains the elephant in the room.^{3,4} Despite the obvious benefits, their replacement with water in homogeneous phase reactions is troublesome as the vast majority of fine chemicals and/or precursors are water insoluble. Literature shows that the addition of surfactants to water can drastically change such a scenario.⁵⁻⁹ Several different chemical processes can be profitably performed in water with high yield, irrespective to the solubility of reagents and products.¹⁰⁻¹⁴ Gathered under the banner of “micellar catalysis” (MC) such methods witnessed an enormous boost in scope, generality and synthetic usefulness in the last 10 years, even at the industrial scale.¹⁵⁻¹⁸

Strictly speaking, MC refers to the catalytic effect on reactions performed in an homogeneous single-phase nanodispersed medium obtained in the presence of a water based surfactant solution above the critical micellar concentration (CMC).¹⁹ Under such conditions, surfactant molecules assemble into micelles, thermodynamically stable association colloids having a hierarchical structure consisting in a lipophilic core and a hydrophilic periphery. Micelles formation is accompanied by a steep increase in solubility of lipophilic reagents/catalysts, mostly localizing in the micelle core. Higher reagents concentration, co-localization and hydrophobic effects play a central role in the observed catalysis.^{19,20}

However, such a prototypical micellar system is limited to low reagents loading. Concentrations lower than 10^{-4} M, far below the saturation concentration which maintains a single isotropic solution (maximum additive concentration, MAC), are typical of earlier kinetic studies.^{21,22} The scope of surfactant enhanced reactions was later extended towards synthetically useful applications. Such shift was accompanied by substantial deviations from the aforescribed micellar model. Formal concentrations of reactants were substantially increased exceeding MAC and thus leading to complex multi-phase systems. As an example, typical experimental conditions involve the use of a 1 wt% to 3 wt% water solution of the

surfactant and 0.5 M to 1 M formal concentrations of poorly water soluble reagents resulting in inhomogeneous mixtures as those depicted in Fig. 1.




Figure 1: Pictures of relevant micellar reactions reported in the literature. A) Direct arylation between N-(m-hydroxyphenyl)-N',N'-dimethylurea and 4-iodoanisole. Copyright 2017 American Chemical Society.¹⁶ B) Suzuki-Miyaura coupling between BTBT-Br and phenylboronic acid. Reproduced from Ref.¹⁴ with permission from the Royal Society of Chemistry. C) Amide bond formation between (S)-N-(benzyloxycarbonyl)valine and tert-Butyl (3R,4S,5S)-3-methoxy-4-(N-methylamino)-5-methylheptanoate hydrochloride. Copyright 2016 American Chemical Society.²³ D) Cross-metathesis reaction between 1-tert-butyltrimethylsilyloxy-2-(2-propenyl)benzene and methylvinylketone. Copyright 2014 American Chemical Society.²⁴

Despite the extensive literature from the group of prof. Lipshutz and prof. Handa amongst others²³ clearly demonstrated that dealing with a complex multiphase system does not preclude effective surfactant enhanced catalysis, unpredictable behavior is often encountered.²⁵ These limits often manifest themselves as lack of reproducibility when changing the experimental set-up, particularly regarding to the mixing method.^{11,23} The use of limited amounts of organic solvents along with mild heating mitigates the problem but does not represent a completely satisfactory solution: processes still rely on organic solvent and no insight on the underlying mechanism is given.^{16,17,23,26} In such complex multi-phase systems the surfactant can still form micelles or other aggregates but also lies at the phase boundary, lowering the interfacial tension and allowing the formation of very small, kinetically stable, droplets/particles.²⁷ Alongside with the *bona fide* micellar catalysis, other surfactant

mediated processes are also at work. Boyer *et al.* showed that under biphasic conditions, the catalytic effect of the surfactant can result from a combination of interfacial catalysis (IC) and an hybrid inverse phase transfer catalysis (IPTC).^{28–30} The latter involves a dual role played by the surfactant: enhancing mass transport of reagents to the water phase and therein promoting the micellar catalysis of the chemical step (Fig. 2). This observation




Figure 2: Principles of an inverse phase transfer catalysis (IPTC) and interfacial catalysis (IC) with a surfactant as catalyst (S, substrate; R, reactant; P, product) as described by Boyer *et al.*²⁹


highlights the fact that in complex multi-phase systems normal and inverse phase transfer catalysis (PTC, IPTC), micellar catalysis (MC) and interface catalysis (IC) become intertwined phenomena. So far, the capability to form micelles of a certain dimension and aspect ratio guided the selection of the most suitable surfactant for reactions in water environment. Such features are critical but insufficient: the capability to maintain efficient mass transfer through different phases during the whole chemical process is at least as important.

We here describe a different approach capable of taming capricious reactions without the use of any organic solvent and/or heating. For this purpose, we sampled some particular examples of troublesome coupling reactions carried out on liquid reagents having issues of reproducibility when performed in aqueous surfactant solutions. Once recognized that the true nature of reactions carried out with liquid reagents and limited amount of surfactant is that of an emulsion, we selected a surfactant mixture specifically devised to better coun-

1
2
3 terbalance the surface tension. In particular, we selected a mixture of two commercially
4 available, food-grade amphiphiles, Tween 80 and L- α -lecithin, known as efficient emulsifiers
5 for oil spills. Under such conditions we demonstrate that it is possible to successfully address
6 all reproducibility issues in terms of mixing method in a Suzuki-Miyaura (SM) coupling so
7 tenaciously irreproducible that not even the cosolvent method afforded satisfactory results.
8 We also show that such approach is general, providing excellent results for other reagents
9 featuring similar behavior.
10
11
12
13
14
15
16
17
18

19 Results and discussion

20
21
22 We identified several SM coupling reactions showing reproducibility issues, some of which
23 are shown in Scheme 1 (additional examples and reactions details are listed in Table S1
24 of the ESI). Differences in the reaction outcome were observed upon modifications in: ves-
25 sel, stirrer/mixing apparatus and stirring speed. Among them, the SM coupling of (4-
26 bromophenyl)methyl acetate (**1**) with 2-thienylboronic acid (shown in heading of Table 1)
27 displays a particularly erratic behavior. Data in the first part of Table 1 (entries 1-8) summa-
28 rize the results obtained upon systematically varying the mixing setup using either a 2 wt%
29 solution of Kolliphor EL (K-EL) or a 2 wt% solution of the state of the art designer surfactant
30 TPGS-750M in water as the reaction medium. The choice of such surfactants was dictated
31 by their literature established performance for micellar coupling reactions.³¹⁻³³ Comparison
32 among entries 1 to 5 clearly shows that conversion is strikingly influenced by the chosen
33 mixing setup, going from 15% to 80%. However, when the reaction was performed twice
34 using the same setup and stirring speed, consistent results were obtained (entry 4-5). In all
35 of the cases, leaving reactions stirring for more than 3 hours did not improve conversions.
36 Not even performing the reaction using the well-known TPGS-750M surfactant allowed to
37 obtain reproducible results (entry 6-7). Such a dependency of conversions from the mixing
38 method suggests that interphase mass transfer may become rate limiting once massive phase
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Scheme 1: Examples of variable results obtained for SM coupling reactions of **1**, **2** and **3** with 2-thienylboronic acid. Reaction conditions: 1 M of bromide, 1 M of boronic acid, 1.5 M of NEt_3 , 0.02 M of Pd(dppf)Cl_2 in 2 wt% aqueous solution of Kolliphor EL, room temperature, 3 hours. Couplings are performed on 2 mmol of bromide, and scale is unvaried in all reactions. Setup a: 10 mL roundbottom test tube (inner diameter: 1.2 cm, height: 9 cm), cylindrical stirrer (2.5 cm length), see Figure S1-A. Setup c: 4 mL vial (inner diameter: 1.3 cm, height: 4.4 cm), two piled round stirrers (diameter: 1.0 cm, height: 0.8 cm), see Figure S1-C. Setup d: 4 mL vial (inner diameter: 1.3 cm, height: 4.4 cm). Reaction was firstly homogenized using an UltraTurrax high shear mixer for 10 minutes at 5000 rpm (see Figure S1-D), and subsequently mixed using two piled round stirrers (diameter: 1.0 cm, height: 0.8 cm).


separation occurs during the evolution of reagents into products. This phenomena appear as the ultimate result of the kinetic instability of the emulsion towards flocculation, coalescence and creaming.³⁴ In support of this hypothesis, when we carried out the coupling using an Ultra-Turrax homogenizer that guaranteed thorough mixing, we consistently observed essentially quantitative conversion (entry 8).

As already mentioned, literature acknowledges that the use of moderate amount of a cosolvent might be an efficient strategy to tackle such mixing issues in micellar catalysis.^{23,26,35,36}

Acetone and THF in particular are recognized as efficient cosolvents for the SM coupling.¹⁶

However, such an approach - which remains a valid and resourceful protocol to clear away

Table 1: SM coupling of **1** with 2-thienylboronic performed with different setups and aqueous media (reaction conditions: 1.0 M of bromide, 1.0 M of boronic acid, 1.5 M of NEt₃, 0.02 M of Pd(dppf)Cl₂, room temperature, 3 hours). Setup *a*, *c* and *d* are described in caption of Scheme 1.



Entry	Setup	Stir. speed (rpm)	Reaction medium	Conv.
1	<i>a</i>	800	2 wt% K-EL in H ₂ O	80%
2	<i>a</i>	1000	2 wt% K-EL in H ₂ O	43%
3	<i>b</i>	800	2 wt% K-EL in H ₂ O	15%
4	<i>c</i>	1000	2 wt% K-EL in H ₂ O	65%
5	<i>c</i>	1000	2 wt% K-EL in H ₂ O	68%
6*	<i>a</i>	800	2 wt% TPGS-750M in H ₂ O	50%
7*	<i>a</i>	1000	2 wt% TPGS-750M in H ₂ O	67%
8	<i>d</i>	1000	2 wt% K-EL in H ₂ O	94%
9	<i>a</i>	800	(2 wt% K-EL in H ₂ O)/THF 9:1	25%
10	<i>a</i>	1000	(2 wt% K-EL in H ₂ O)/THF 9:1	72%
11	<i>a</i>	800	(2 wt% K-EL in H ₂ O)/acetone 9:1	70%
12	<i>a</i>	1000	(2 wt% K-EL in H ₂ O)/acetone 9:1	61%

Setup *b*: 100 mL roundbottom test tube (inner diameter: 2.7 cm, height: 20 cm), cylindrical stirrer (2 cm length), see Figure S1-B.

* Reaction was performed under N₂ atmosphere.

most of the toxic organic solvents - in our case improved only modestly the behavior of the reaction (entry 9-12) and, at the same time, does not fully honor the commitment to clear away organic solvents.

Once recognized that the reproducibility was dependent on the kinetic stability of obtained emulsion, we decided to approach the problem using an improved surfactant mixture able to tackle this issue. In particular, taking inspiration from the literature pertaining self-emulsifying systems (i.e. systems requiring very little energy to form fine emulsions),^{37,38} we chose to test the Tween 80/L- α -lecithin surfactants mixture. Thanks to their synergistic

action, this food-grade amphiphiles combination already proved to be an effective and safer alternative to toxic solvent-based dispersants for marine oil spills^{39–41} and also found application in drug-delivery.⁴² We thus decided to test an aqueous 2 wt% Tween 80/lecithin 8:2 mixture (TL82), suitable to obtain oil-in-water emulsions, as the medium for our troublesome SM coupling. This particular ratio was selected in order to obtain an HLB (Hydrophilic Lipophilic Balance) value similar to that of both Kolliphor EL (with which we already had a vast experience) and TPGS-750-M, the standard in micellar catalysis. Also, lecithin is not water soluble and using mixtures at a higher lecithin content would have led to poor dispersibility in water.

The first striking difference with prior experiments was observed during sample preparation: a simple visual comparison, Fig. 3, highlights the superior stability of the emulsion of **1** in TL82 *vs.* that obtained in K-EL. While both mixtures were prepared identically and




Figure 3: Coalescence of the emulsions of **1** (1 mmol) prepared in 1 mL of a 2 wt% aqueous solution of K-EL (left) and TL82 (right).

thoroughly mixed with the aid of an ultrasound bath, in the case of K-EL rapid coalescence occurs and two layers separate. Conversely, a fine and stabler emulsion forms when using TL82, meaning that the surface tension of this system is lower and droplets of organic phase are stabilized more effectively.

Table 2 summarizes results obtained performing the coupling reaction in both aqueous K-EL and TL82 surfactants mixture with two different setups.

Table 2: Conversions for the SM coupling shown in Scheme 1 performed in 2 wt% aqueous solution of K-EL or TL82. The stirring setup (10 mL roundbottom test-tube equipped with a cylindrical stirring bar) is unvaried in all reactions.


Entry	Surfactant	Atm.	Stirring speed (rpm)	Conversion (Time)
1	K-EL	air	800	80% (3 h)
2	K-EL	air	1000	43% (3 h)
3	TL82	air	800	90% (1 h)
4	TL82	air	1000	92% (1 h)
5	TL82	N ₂	800	97% (1 h)

Reactions performed in TL82 do not show the reproducibility issues previously observed with K-EL and TPGS-750M, with conversion rising to 90% in 1 hour when working under air unregarding to the mixing setup used to perform the reaction (entry 3-4). Employing such self-emulsifying system allows obtaining an emulsion whose features are essentially independent from the mixing method due to the reduced surface tension. The observed higher catalytic activity is also believed to be a consequence of the formation of a finer and stable emulsion and unhindered interphase mass transport.

Performing the reaction under inert atmosphere allowed reaching an even higher 97% conversion (entry 5). This last result was expected, as environmental oxygen is a known poisoning agent for the Pd(0) catalyst involved in the reaction.

To exclude an effect of ionic interactions between the 2-thienyl boronate and lecithin (due to its zwitterionic nature) in the observed reaction behavior, we also performed the coupling using a K-EL/lecithin 8:2 mixture, which does not have self-emulsifying capabilities. With this surfactants mixture the reaction gave again unreliable results (See Table S2 in the ESI). As the emulsifying capabilities of the TL82 mixture proved suitable to both overcome the reproducibility issues of the chosen test coupling and to improve the reaction conversion, we decided to test its performance on other substrates featuring similar behavior. We therefore performed the coupling reaction of 2-thienylboronic acid with other aryl bromides: 4-

bromoanisole (**2**), 3-bromoquinoline (**3**) and 3-bromothiophene (**4**). Scheme 2 summarizes the obtained results.



Entry	Product	Surfactant	Conversion (Time)
1	2b	K-EL	58% (3 h)
2	2b	TL82	100% (1 h)
3	3b	K-EL	63% (3 h)
4	3b	TL82	94% (3 h)
5	4b	K-EL	60% (3 h)
6	4b	TL82	100% (1 h)

Scheme 2: Conversions obtained for the coupling reaction between **2**, **3** and **4** with 2-thienylboronic acid (1 M of bromide, 1 M of boronic acid, 1.5 M of NEt_3 , 0.02 M of Pd(dppf)Cl_2 , room temperature) in 2 wt% surfactant solution of K-EL or TL82. Reactions performed with TL82 surfactant were run under N_2 atmosphere. The stirring setup (10 mL roundbottom test-tube equipped with a cylindrical stirring bar, stirring speed of 800 rpm) is unvaried in all reactions.

In all of the cases, reactions performed using the K-EL surfactant did not exceed 65% conversion after 3 hours. Using the TL82 surfactants mixture we obtained quantitative conversion in the case of substrates **2** and **4** in just one hour, and 94% conversion for substrate **3** in 3 hours. As observed for substrate **1**, the use of TL82 allows to obtain higher conversions in a shorter time. Preliminary results show that the TL82 can be successfully applied in the Sonogashira and an Heck couplings as well (see Scheme S1 of the ESI). In terms of stability under the reaction conditions required by the various cross couplings we tested, the TL82

mixture appears to be suitably robust (see Section 5 of the ESI). Indeed, lecithin itself is reported to be particularly stable at high pH levels and in the absence of oxygen. Such conditions are very similar to those we employed in our reactions.^{43–45}

Finally, we tested the robustness towards scaling up of our protocol by performing the coupling reaction of **1** with 2-thienylboronic acid on 10 g of reagents (a 14th fold increase in the scale) in a round-bottom flask and using a mechanic stirrer, under otherwise identical conditions to Table 2, entry 5 test. Upon the addition of NEt_3 , the initial biphasic mixture turned into an homogeneous emulsion (Figure 4), and the reaction spontaneously warmed up to 35 °C over ten minutes. We achieved complete conversion within 1 h and isolated the




Figure 4: SM coupling of **1** with 2-thienylboronic on a 10 g scale. a) Reaction mixture appearance before the addition of NEt_3 b) Reaction mixture appearance one minute after the addition of NEt_3 c) Reaction mixture appearance one hour after the addition of NEt_3 .

target compound in 97% yield. Lowering the catalyst amount to 1.5 mol% and 1 mol% of $\text{Pd}(\text{dppf})\text{Cl}_2$ gave respectively a 83% and 78% conversion after 3 hours. Literature reports examples of catalysts that are far more effective than the commercially available one we selected. In particular, Liphshutz and Handa demonstrated the possibility to carry out SM and Sonogashira reactions at ppm levels of Pd.^{46,47} Such a solution also represents the ideal strategy to reduce possible palladium contamination in the product, a problem we intentionally did not address in this paper. We here focus only on demonstrating that TL82 mixture enables running efficient SM coupling on otherwise troublesome substrates even in the large scale.

Concerned about the possible hazards connected with the exothermic behavior we observed, we repeated the reaction three more times on the same scale, with the same setup and at 2 mol% catalyst loading. The first repetition was performed working identically, but with a thermometer inserted in the reaction mixture (previously we monitored the temperature of the reactor's walls by means of an IR scanner). The maximum temperature reached was 50 °C, and conversion was again quantitative after 1 h. In the second repetition we added the base dropwise over a period of 10 minutes, while keeping the reactor immersed in a water bath. In this case, we did not measure any appreciable raise in the temperature, yet we noted that such a slow addition is not helpful for the homogeneity of the reaction mixture. Lumps of sticky material formed, which were difficult to redisperse. The conversion was 72% and the isolated yield was 68%. In the third repetition we studied the effect of the slow addition of the catalyst. While setting up the reaction, we added the base all at once to the emulsion of bromide and boronic acid. We noticed that just after the addition of the base and prior to the introduction of any catalyst the reaction warmed up to almost 40 °C, meaning that most of the exothermic effect is connected with the acid-base reaction of the boronic acid with the base. This observation reduces the concerns for possible decomposition reactions leading to runaway thermal effects. Once back at room temperature, we added the catalyst over a period of 10 minutes. Again, we observed some demixing while doing so. The conversion was 84% and the isolated yield was 82%. Puzzled by the partial demixing effect observed in both cases, we heated samples taken from the second and third repetition at 50 °C by external heating, thereby restoring a homogeneous, freely stirring emulsion. Our observations suggest that reaching a temperature around 50 °C in the early stage of the reaction is helpful in maintaining a homogeneous emulsion (well in agreement with the fact that emulsions are easier to form at high temperature).

Conclusions

Micellar reactions carried out at formal concentrations vastly exceeding MAC are strongly impacting the sustainability of a vast number of chemical transformations including cross-couplings. We have faced the known issue of lack of reproducibility, particularly when changing experimental set up and reaction scale, from a different perspective with respect to the existing literature. Whereas the general consensus to tame problematic micellar reactions requires the use of an organic cosolvent, we focused our attention on the details of the formulation state of the reaction mixture and carefully tuned the tensides cocktail accordingly. Above MAC, micellar reactions are complex heterogeneous mixtures where micellar catalysis coexists with other catalytic mechanisms like IC and IPTC. The mass transfer between emulsion droplets, dispersed particles and micelles dominates the reaction behavior to the point that the capability of a given surfactant to assemble into micelles of a certain shape and/or volume has only a limited impact on the overall conversion. On the other hand, the capability of the tensides to efficiently and rapidly emulsify the oil phase maximizing the contact area of all interphases becomes the limiting step. Indeed, we show that the use of a Tween 80/L- α -lecithin surfactants mixture, a system well known to readily provide finely dispersed emulsion of the most hostile hydrophobes, leads to satisfactory results in a number of S-M couplings where other commercial and designer surfactants failed, even in the presence of cosolvents. Even more importantly, all reactions proved to be insensitive to setup parameters like the stirring speed and reaction scale, generally heavily impacting on standard micellar protocols. Our findings clearly show that the capability of a given surfactant to self-assemble into micelles with given characteristics does not necessarily grant success in all reactions. Above MAC, the true nature of the reactive formulation at hand requires balancing of interface tension between complex pseudo phases, thus favoring the selection of high surface pressure tensides. Our approach is general, does not require the use of any organic solvent and is based on industrially scaled, food grade surfactants.

Experimental details

General Information. All reagents, Pd(dppf)Cl₂ and Pd(dtbpf)Cl₂ catalysts were bought from Fluorochem. [Pd(π -cinnamyl)Cl]₂, surfactants and solvents were bought from Sigma-Aldrich. L- α -lecithin (95%) was bought from Nutritional Biochemical Corporation. For reactions performed under anaerobic environment, water was carefully deoxygenated before use by bubbling N₂ under reflux for three hours, and NEt₃ was carefully deoxygenated by freeze-pump-thaw degassing.

Reaction conversions were quantified by gas chromatography-mass spectroscopy (GC-MS) or ¹H NMR. For reactions performed on substrates **1**, **2** and **3**, composition of reaction mixtures was quantified by quantitative GC-MS through peak integration based on response factor method (see ESI for details). For reactions performed on other substrates, composition of reaction mixtures was quantified by ¹H NMR.

General procedure for small scale reactions. Unless when otherwise stated, reactions were performed in a roundbottom test tube (10 mL volume, available from CEM Corporation) equipped with a cylindrical stirring bar and under standard laboratory atmosphere (setup *a* shown in figure S1A of the ESI). In a typical procedure, bromide (2 mmol), boronic acid (2 mmol) and Pd(dppf)Cl₂ (0.04 mmol) were weighted in the vessel and 2 mL of the chosen surfactants solution was added under stirring. After 5 minutes N(Et)₃ (3 mmol) was added. The reaction mixture was stirred for 3 hours, and subsequently diluted with CH₂Cl₂, filtered through a pad of silica and submitted to GC-MS characterization.

Scaled-up synthesis of 1b. Reaction was performed in a 250 mL 3-necked roundbottom flask equipped with a mechanical stirrer and a thermometer. 2-thienyl boronic acid (3.583 g, 28.00 mmol) and Pd(dppf)Cl₂ (410 mg, 0.560 mmol) were weighted in the flask, which was subsequently put under N₂ atmosphere. Methyl-4-bromophenylacetate (6.414 g, 28.00 mmol) and degassed TL82 surfactants mixture (28 mL) were subsequently added. The mixture was

1
2
3 stirred at 300 rpm for 5 minutes before addition of degassed $N(Et)_3$ (5.9 mL, 42 mmol). The
4
5 reaction spontaneously warmed up to 50 °C over 10 minutes. The reaction was stirred for 1
6
7 hour. Excess bromide was steam distilled from the reaction mixture by means of a Claisen
8
9 distillation apparatus. During the distillation, the volume of the reservoir was kept constant
10
11 at 100 mL by constant addition of water. The composition of both the distillate and the
12
13 mixture was monitored constantly by GC-MS. When no bromide was left in the reservoir, the
14
15 mixture was cooled at 4 °C and the precipitate thus formed was isolated by suction filtration,
16
17 washed thoroughly on the filter with water and dried till constant weight. 6.302 g, 27.1 mmol,
18
19 96.9% (m.p. 50-51 °C). The raw material was submitted to NMR characterization. Further
20
21 purification was performed by evaporative distillation (water bath, 90 °C, 0.02 mbar) using a
22
23 standard sublimation apparatus (m.p. 54-55 °C). The NMR spectra of the raw and sublimed
24
25 materials (Figure S2) are nearly identical. 1H NMR (400 Hz, $CDCl_3$): δ [ppm] 7.57 (d, $J=8.3$
26
27 Hz, 2H), 7.30-7.26 (m, 4H), 7.07 (dd, $J=5.1, 3.6$ Hz, 1H), 3.71 (s, 3H), 3.64 (s, 2H).
28
29
30

31 Acknowledgement

32
33
34 Financial support from the Italian Ministry of University and Research (MIUR) through
35
36 grant “Dipartimenti di Eccellenza 2017 Materials For Energy” and PRIN-2017 research pro-
37
38 gramme (project “BOOSTER”) is gratefully acknowledged.
39
40
41

42 Supporting Information Available

43
44
45 Pictures of mixing setups. Additional reactions. E-factor of the scaled-up synthesis of **1b**.
46
47 Assessment of TL82 stability under working conditions. Micrographs of the emulsion of **1**
48
49 in aqueous TL82.
50
51
52
53
54
55
56
57
58
59
60

References

- (1) Sheldon, R. A. Fundamentals of green chemistry: Efficiency in reaction design. *Chem. Soc. Rev.* **2012**, *41*, 1437–1451.
- (2) Sheldon, R. A. The E factor 25 years on: the rise of green chemistry and sustainability. *Green Chem.* **2017**, *19*, 18–43.
- (3) Clarke, C. J.; Tu, W.-C.; Levers, O.; Bröhl, A.; Hallett, J. P. Green and Sustainable Solvents in Chemical Processes. *Chem. Rev.* **2018**, *118*, 747–800.
- (4) Alder, C. M.; Hayler, J. D.; Henderson, R. K.; Redman, A. M.; Shukla, L.; Shuster, L. E.; Sneddon, H. F. Updating and further expanding GSK's solvent sustainability guide. *Green Chem.* **2016**, *18*, 3879–3890.
- (5) Sorella, G. L.; Strukul, G.; Scarso, A. Recent advances in catalysis in micellar media. *Green Chem.* **2015**, *17*, 644–683.
- (6) Gallou, F.; Ostaszewski, R.; Walde, P.; Serrano-Luginbühl, S.; Ruiz-Mirazo, K. Soft and dispersed interface-rich aqueous systems that promote and guide chemical reactions. *Nat. Rev. Chem.* **2018**, *2*, 306–327.
- (7) Romney, D. K.; Arnold, F. H.; Lipshutz, B. H.; Li, C.-J. Chemistry Takes a Bath: Reactions in Aqueous Media. *J. Org. Chem.* **2018**, *83*, 7319–7322.
- (8) Kitanosono, T.; Masuda, K.; Xu, P.; Kobayashi, S. Catalytic Organic Reactions in Water toward Sustainable Society. *Chem. Rev.* **2018**, *118*, 679–746.
- (9) Sassi, M.; Mattiello, S.; Beverina, L. Syntheses of Organic Semiconductors in Water. Recent Advancement in the Surfactants Enhanced Green Access to Polyconjugated Molecules. *Eur. J. Org. Chem.* **2020**, *2020*, 3942–3953.
- (10) Lipshutz, B. H.; Ghorai, S. “Designer”-surfactant-enabled cross-couplings in water at room temperature. *Aldrichimica Acta* **2012**, *45*, 3–16.

- (11) Lipshutz, B. H. When Does Organic Chemistry Follow Nature's Lead and "Make the Switch"? *J. Org. Chem.* **2017**, *82*, 2806–2816.
- (12) Lempke, L.; Ernst, A.; Kahl, F.; Weberskirch, R.; Krause, N. Sustainable Micellar Gold Catalysis – Poly(2-oxazolines) as Versatile Amphiphiles. *Adv. Synth. Catal.* **2016**, *358*, 1491–1499.
- (13) Yetra, S. R.; Rogge, T.; Warratz, S.; Struwe, J.; Peng, W.; Vana, P.; Ackermann, L. Micellar Catalysis for Ruthenium(II)-Catalyzed C-H Arylation: Weak-Coordination-Enabled C-H Activation in H₂O. *Angew. Chem. Int. Ed.* **2019**, *58*, 7490–7494.
- (14) Sanzone, A.; Mattiello, S.; Garavaglia, G. M.; Calascibetta, A. M.; Ceriani, C.; Sassi, M.; Beverina, L. Efficient synthesis of organic semiconductors by Suzuki–Miyaura coupling in an aromatic micellar medium. *Green Chem.* **2019**, *21*, 4400–4405.
- (15) Gallou, F.; Isley, N. A.; Ganic, A.; Onken, U.; Parmentier, M. Surfactant technology applied toward an active pharmaceutical ingredient: more than a simple green chemistry advance. *Green Chem.* **2016**, *18*, 14–19.
- (16) Gabriel, C. M.; Lee, N. R.; Bigorne, F.; Klumphu, P.; Parmentier, M.; Gallou, F.; Lipshutz, B. H. Effects of Co-solvents on Reactions Run under Micellar Catalysis Conditions. *Org. Lett.* **2017**, *19*, 194–197.
- (17) Lee, N. R.; Gallou, F.; Lipshutz, B. H. S_NAr Reactions in Aqueous Nanomicelles: From Milligrams to Grams with No Dipolar Aprotic Solvents Needed. *Org. Process Res. Dev.* **2017**, *21*, 218–221.
- (18) Sharma, S.; Das, J.; Braje, W. M.; Dash, A. K.; Handa, S. A Glimpse into Green Chemistry Practices in the Pharmaceutical Industry. *ChemSusChem* **2020**, *13*, 2859–2875.
- (19) Khan, M. N. *Micellar catalysis*; CRC Press, 2006.

- (20) Mancin, F.; Scrimin, P.; Tecilla, P.; Tonellato, U. Amphiphilic metalloaggregates: Catalysis, transport, and sensing. *Coord. Chem. Rev.* **2009**, *253*, 2150 – 2165.
- (21) Fendler, E. J.; Fendler, J. H. *Advances in Physical Organic Chemistry*; Elsevier, 1970; Vol. 8; pp 271–406.
- (22) Dwars, T.; Paetzold, E.; Oehme, G. Reactions in Micellar Systems. *Angew. Chem. Int. Ed.* **2005**, *44*, 7174–7199.
- (23) Gallou, F.; Guo, P.; Parmentier, M.; Zhou, J. A General and Practical Alternative to Polar Aprotic Solvents Exemplified on an Amide Bond Formation. *Org. Process Res. Dev.* **2016**, *20*, 1388–1391.
- (24) Klumphu, P.; Lipshutz, B. H. “Nok”: A Phytosterol-Based Amphiphile Enabling Transition-Metal-Catalyzed Couplings in Water at Room Temperature. *J. Org. Chem.* **2014**, *79*, 888–900.
- (25) Lipshutz, B. H.; Petersen, T. B.; Abela, A. R. Room-Temperature Suzuki-Miyaura Couplings in Water Facilitated by Nonionic Amphiphiles. *Org. Lett.* **2008**, *10*, 1333–1336.
- (26) Brocklehurst, C. E.; Gallou, F.; Hartwig, J. C. D.; Palmieri, M.; Ruffe, D. Microtiter Plate (MTP) Reaction Screening and Optimization of Surfactant Chemistry: Examples of Suzuki-Miyaura and Buchwald-Hartwig Cross-Couplings in Water. *Org. Process Res. Dev.* **2018**, *22*, 1453–1457.
- (27) Mollet, H.; Grubenmann, A. *Formulation Technology: emulsions, suspensions, solid forms*; John Wiley & Sons, 2008.
- (28) Boyer, B.; Hambardzoumian, A.; Roque, J.-P.; Beylerian, N. Reaction in Biphasic Water/Organic Solvent System in the Presence of Surfactant: Inverse Phase Transfer Catalysis versus Interfacial Catalysis. *Tetrahedron* **2000**, *56*, 303–307.

- (29) Boyer, B.; Hambardzoumian, A.; Roque, J.-P.; Beylerian, N. Inverse phase transfer catalysis versus interfacial catalysis. Effect of medium stirring in the epoxidation reaction of chalcone by hydrogen peroxide. *J. Chem. Soc., Perkin Trans. 2* **2002**, 1689–1691.
- (30) Piradashvili, K.; Alexandrino, E. M.; Wurm, F. R.; Landfester, K. Reactions and Polymerizations at the Liquid–Liquid Interface. *Chem. Rev.* **2015**, *116*, 2141–2169.
- (31) Mattiello, S.; Rooney, M.; Sanzone, A.; Brazzo, P.; Sassi, M.; Beverina, L. Suzuki-Miyaura Micellar Cross-Coupling in Water, at Room Temperature, and under Aerobic Atmosphere. *Org. Lett.* **2017**, *19*, 654–657.
- (32) Lipshutz, B. H.; Ghorai, S.; Abela, A. R.; Moser, R.; Nishikata, T.; Duplais, C.; Gaston, R. D.; Gadwood, R. C. TPGS-750-M: A Second-Generation Amphiphile for Metal-Catalyzed Cross-Couplings in Water at Room Temperature. *J. Org. Chem.* **2011**, *76*, 4379–4391.
- (33) Andersson, M. P.; Gallou, F.; Klumphu, P.; Takale, B. S.; Lipshutz, B. H. Structure of nanoparticles derived from designer surfactant TPGS-750-M in water, as used in organic synthesis. *Chem. Eur. J.* **2018**, *24*, 6778–6786.
- (34) Capek, I. Degradation of kinetically-stable o/w emulsions. *Adv. Colloid Interface Sci.* **2004**, *107*, 125–155.
- (35) Rooney, M.; Mattiello, S.; Stara, R.; Sanzone, A.; Brazzo, P.; Sassi, M.; Beverina, L. Suzuki-Miyaura cross-coupling of latent pigments in water/toluene emulsion under aerobic atmosphere. *Dyes Pigm.* **2018**, *149*, 893–901.
- (36) Vaghi, L.; Sanzone, A.; Sassi, M.; Pagani, S.; Papagni, A.; Beverina, L. Synthesis of Fluorinated Acridines via Sequential Micellar Buchwald-Hartwig Amination/Cyclization of Aryl Bromides. *Synthesis* **2018**, *50*, 1621–1628.

- (37) Lawrence, M.; Rees, G. D. Microemulsion-based media as novel drug delivery systems. *Adv. Drug Deliv. Rev.* **2000**, *45*, 89–121.
- (38) Solans, C.; Morales, D.; Homs, M. Spontaneous emulsification. *Curr. Opin. Colloid Interface Sci.* **2016**, *22*, 88–93.
- (39) Riehm, D. A.; Neilsen, J. E.; Bothun, G. D.; John, V. T.; Raghavan, S. R.; McCormick, A. V. Efficient dispersion of crude oil by blends of food-grade surfactants: Toward greener oil-spill treatments. *Mar. Pollut. Bull.* **2015**, *101*, 92–97.
- (40) Riehm, D. A.; Rokke, D. J.; Paul, P. G.; Lee, H. S.; Vizanko, B. S.; McCormick, A. V. Dispersion of oil into water using lecithin-Tween 80 blends: The role of spontaneous emulsification. *J. Colloid Interface Sci.* **2017**, *487*, 52–59.
- (41) Athas, J. C.; Jun, K.; McCafferty, C.; Owoseni, O.; John, V. T.; Raghavan, S. R. An Effective Dispersant for Oil Spills Based on Food-Grade Amphiphiles. *Langmuir* **2014**, *30*, 9285–9294.
- (42) Kelmann, R. G.; Kuminek, G.; Teixeira, H. F.; Koester, L. S. Carbamazepine parenteral nanoemulsions prepared by spontaneous emulsification process. *Int. J. Pharm.* **2007**, *342*, 231–239.
- (43) Haidar, I.; Harding, I. H.; Bowater, I. C.; Eldridge, D. S.; Charman, W. N. The role of lecithin degradation on the pH dependent stability of halofantrine encapsulated fat nano-emulsions. *Int. J. Pharm.* **2017**, *528*, 524–535.
- (44) Long, J. H. OBSERVATIONS ON THE STABILITY OF LECITHIN. *J. Am. Chem. Soc.* **1908**, *30*, 881–895.
- (45) Elworthy, P. H.; Saunders, L. 58. Surface force and stability of lecithin sols. *J. Chem. Soc.* **1957**, 330–335.

- (46) Handa, S.; Smith, J. D.; Zhang, Y.; Takale, B. S.; Gallou, F.; Lipshutz, B. H. Sustainable HandaPhos-ppm Palladium Technology for Copper-Free Sonogashira Couplings in Water under Mild Conditions. *Org. Lett.* **2018**, *20*, 542–545.
- (47) Takale, B. S.; Thakore, R. R.; Handa, S.; Gallou, F.; Reilly, J.; Lipshutz, B. H. A new, substituted palladacycle for ppm level Pd-catalyzed Suzuki–Miyaura cross couplings in water. *Chem. Sci.* **2019**, *10*, 8825–8831.

Graphical TOC Entry

