

Synthesis of (*cis*-6-Methyltetrahydropyran-2-yl)acetic Acid Involving the Use of an Organoselenium-mediated Cyclization Reaction

Steven V. Ley,* Barry Lygo, Huguetta Molines, and Judith A. Morton

Department of Chemistry, Imperial College, London SW7 2AY, U.K.

A short stereospecific synthesis of (*cis*-6-methyltetrahydropyran-2-yl)acetic acid (**1**) has been achieved from readily available starting materials using a novel organoselenium-mediated cyclization of alkenyl-substituted β -oxoesters.


The use of organoselenium-mediated cyclizations to construct both hetero¹- and carbo²-cyclic ring systems is now well recognised. Here we show how methodology developed in our laboratories³ can be used efficiently to synthesise (*cis*-6-methyltetrahydropyran-2-yl)acetic acid (**1**), a natural product recently isolated from *Viverra civetta*.⁴

The readily available alkenyl-substituted β -oxoester⁵ (**2**) reacts at room temperature with *N*-phenylselenophthalimide (NPSP),⁶ and trace amounts (0.01 equiv.) of tin tetrachloride to give a 9:1 mixture of compounds (**3**, R=Me) and (**4**, R=Me) in 84% combined yield. This mixture can be used directly in the next reaction; however, if conventional flash chromatography is used to purify the products, (**4**) is seen to rearrange to (**3**) which can be isolated in a pure form.[†] Treatment of (**2**) with NPSP and greater quantities of SnCl₄ (1 equiv.) results in the rapid formation of the carbocyclic product (**5**) in 83% yield.

Reduction of (**3**) and (**4**) with tri-*n*-butyltin hydride⁷ proceeds well, without ring opening, to give (**6**, R=Me) and (**7**, R=Me) in 64% yield. Upon similar treatment, pure (**3**) gave (**6**) in comparable yield.

Hydrogenation of (**6**) and (**7**) using Raney-nickel as a catalyst gave a single compound shown to be methyl (*cis*-6-methyltetrahydropyran-2-yl)acetate (**8**) in 71% yield. This ester was identical with the previously synthesised material.⁴ Hydrolysis of (**8**) gave a 92% yield of (*cis*-6-methyltetrahydropyran-2-yl)acetic acid (**1**).^{4,8} [m.p. 51–52 °C; ¹H n.m.r. δ 10.1 (1H, br.s, CO₂H), 3.84–3.72 (1H, m, H_X), 3.53 (1H, m, H_C), 2.9 (1H, q, H_B, *J*_{AB} 15, *J*_{BX} 7.5 Hz), 2.48 (1H, q, H_A, *J*_{AB} 15 Hz), 1.9–1.15 (6H, m), and 1.19 (3H, d, *J* 6.3 Hz)].

In an effort to simplify this route further we chose to combine the two reduction steps with a final deprotection reaction.



Scheme 1. Reagents and conditions: i, NPSP–SnCl₄ (1.1:0.01), room temp., CH₂Cl₂, 2 h; ii, NPSP–SnCl₄ (1.1:1.0), room temp., CH₂Cl₂, 30 min; iii, Bu₃SnH, azoisobutyronitrile, toluene, heat, 1 h; iv, H₂–Raney-nickel, 100 atm, EtOH, 60 °C, 20 h; v, 10% NaOH, MeOH–H₂O, heat, 15 min.

Thus, when the benzyl-substituted compound (**9**) was treated with NPSP–SnCl₄ (1.1:0.01 equiv.) as before, a 4:1 mixture of (**3**, R=CH₂Ph) and (**4**, R=CH₂Ph) was obtained in 57%

[†] All new compounds were fully characterised by spectroscopic and microanalytical methods.

yield. This mixture, on reduction with H_2 -Raney-nickel, afforded (1) in one step in an unoptimised 47% yield.

We thank the S.E.R.C. for research studentships (to B.L. and J.A.M.), N.A.T.O. for a study grant (to H. M.), and the R.S.C. for the Hickinbottom Research Award (to S.V.L.).

Received, 20th August 1982; Com. 1009

References

- 1 For example see M. de M. Campos and N. Petragnani, *Chem. Ber.*, 1960, **93**, 317; K. C. Nicolaou, *Tetrahedron*, 1981, **37**, 4097, and references therein; D. L. J. Clive, C. G. Russell, G. Chittattu, and A. Singh, *ibid.*, 1980, **36**, 1399; Z. Lysenko, F. Ricciardi, J. E. Semple, P. C. Wang, and M. M. Joullie, *Tetrahedron Lett.*, 1978, 2679; P. L. Beaulieu, V. M. Morisset, and D. G. Garratt, *ibid.*, 1980, 129; D. L. J. Clive, V. Farina, A. Singh, C. K. Wong, W. A. Kiel, and S. M. Menchen, *J. Org. Chem.*, 1980, **45**, 2120; T. R. Hoye and A. J. Caruso, *ibid.*, 1981, **46**, 1198; A. Toshimitsu, T. Aoi, S. Uemura, and M. Okano, *ibid.*, 1981, **46**, 3021; K. C. Nicolaou, R. L. Magolda, W. J. Sipio, W. E. Barnette, Z. Lysenko, and M. M. Joullie, *J. Am. Chem. Soc.*, 1980, **102**, 3784; K. C. Nicolaou, S. P. Seitz, W. J. Sipio, and J. F. Blount, *ibid.*, 1979, **101**, 3884; M. Petrzilka, *Helv. Chim. Acta*, 1978, **61**, 3075; G. Jaurand, J.-M. Beau, and P. Sinay, *J. Chem. Soc., Chem. Commun.*, 1981, 572.
- 2 For example see D. L. J. Clive, G. Chittattu, and C. K. Wong, *J. Chem. Soc., Chem. Commun.*, 1978, 441; A. Toshimitsu, S. Uemura, and M. Okano, *ibid.*, 1982, 87; T. Kametani, K. Suzuki, H. Kurobe, and H. Nemoto, *Chem. Pharm. Bull.*, 1981, **29**, 105, and previous papers in the series; F. Rouessac and H. Zamarlik, *Tetrahedron Lett.*, 1981, 2643; M. Alderdice and L. Weiler, *Can. J. Chem.*, 1981, **59**, 2239; D. G. Garratt, M. D. Ryan, and A. Kabo, *ibid.*, 1980, **58**, 2329.
- 3 W. P. Jackson, S. V. Ley, and J. A. Morton, *J. Chem. Soc., Chem. Commun.*, 1980, 1028; W. P. Jackson, S. V. Ley, and A. J. Whittle, *ibid.*, 1980, 1173; W. P. Jackson, S. V. Ley, and J. A. Morton, *Tetrahedron Lett.*, 1981, 2601.
- 4 B. Maurer, A. Grieder, and W. Thommen, *Helv. Chim. Acta*, 1979, **62**, 44; B. Maurer and W. Thommen, *ibid.*, 1979, **62**, 1096.
- 5 P.-E. Sum and L. Weiler, *Can. J. Chem.*, 1979, **57**, 1475.
- 6 K. C. Nicolaou, D. A. Claremon, W. E. Barnette, and S. P. Seitz, *J. Am. Chem. Soc.*, 1979, **101**, 3704.
- 7 D. L. J. Clive, G. J. Chittattu, V. Farina, W. A. Kiel, S. M. Menchen, C. G. Russell, A. Singh, C. K. Wong, and N. J. Curtis, *J. Am. Chem. Soc.*, 1980, **102**, 4438.
- 8 R. S. Mali, M. Pohmakotr, B. Weidmann, and D. Seebach, *Liebigs Ann. Chem.*, 1981, 2272; D. Seebach, M. Pohmakotr, C. Schregenberger, B. Weidmann, R. S. Mali, and S. Pohmakotr, *Helv. Chim. Acta*, 1982, **65**, 419.