

Aerobic Photooxidation of Phosphite Esters Using Diorganotelluride Catalysts


Makoto Oba,[†] Yasunori Okada,[†] Kozaburo Nishiyama,^{*,†} and Wataru Ando^{*,‡}

Department of Materials Chemistry, Tokai University, 317 Nishino, Numazu, Shizuoka 410-0395, Japan, and National Institute of Advanced Industrial Science and Technology (AIST), 1-1-1 Higashi, Tsukuba, Ibaraki 305-8565, Japan

nishiyam@wing.ncc.u-tokai.ac.jp; wataru.ando@aist.go.jp

Received February 4, 2009

ABSTRACT



Diorganotellurides containing bulky aromatic substituents are found to catalyze the photooxidation of phosphite esters using aerobic oxygen as a terminal oxidant. A Hammett plot with substituted triaryl phosphites yielding $\rho = 2.88$ agrees with a nucleophilic oxygen transfer from telluroxide to phosphite.

Oxidation reactions are a significant class of important and essential chemical processes. In general, such reactions are conducted using traditional inorganic and organic oxidants, such as dichromates, permanganates, organic peroxides, etc., stoichiometrically.¹ Recently, increasingly severe environmental constraints have generated greater interest in the use of environmentally benign molecular oxygen or hydrogen peroxide as oxidants.² Although oxidation of organic molecules by the ground-state triplet oxygen or hydrogen peroxide is thermodynamically favored, they are not kinetically competent to serve as an oxidant. Therefore, a large number of catalytic strategies have been proposed to achieve rate enhancement in the oxidation processes that have been developed.³

Direct oxygenation of an organic substrate through singlet oxygen via [2 + 2], [4 + 2], and ene additions is one of the most often used pathways in organic synthesis,⁴ because the reaction rate of singlet oxygen is much greater than that of

triplet oxygen as a result of its low activation energy. Furthermore, photosensitized oxygenations of organosulfur⁵ and phosphorus⁶ compounds have been extensively studied with respect to their synthetic and mechanistic aspects. However, reports explaining the use of singlet oxygen in catalytic systems are still scarce. It was demonstrated independently by Abatjoglou and Krief that diorganoselenides promoted the reoxidation of osmium(VI) to osmium(VIII) using singlet oxygen in the olefin dihydroxylation

(3) For recent reviews, see: (a) Matsumoto, T.; Ueno, M.; Wang, N.; Kobayashi, S. *Chem. Asian J.* **2008**, *3*, 196–214. (b) Imada, Y.; Naota, T. *Chem. Rec.* **2007**, *7*, 354–361. (c) Schultz, M. J.; Sigman, M. S. *Tetrahedron* **2006**, *62*, 8227–8241. (d) Stahl, S. S. *Angew. Chem., Int. Ed.* **2004**, *43*, 3400–3420. (e) Sheldon, R. A.; Arends, I. W. C. E.; ten Brink, G.-J.; Dijkstra, A. *Acc. Chem. Res.* **2002**, *35*, 774–779.

(4) (a) Clennan, E. L. *Tetrahedron* **2000**, *56*, 9151–9179. (b) Stratakis, M.; Orfanopoulos, M. *Tetrahedron* **2000**, *56*, 1595–1615. (c) Wasserman, H. H.; Ives, J. L. *Tetrahedron* **1981**, *37*, 1825–1852. (d) Prein, M.; Adam, W. *Angew. Chem., Int. Ed.* **1996**, *35*, 477–494. (e) Adam, W.; Prein, M. *Acc. Chem. Res.* **1996**, *29*, 275–283. (f) Schenck, G. O.; Eggert, H.; Denk, W. *Liebigs Ann. Chem.* **1953**, *584*, 177–198. (g) Foote, C. S.; Wexler, S. *J. Am. Chem. Soc.* **1964**, *86*, 3979–3981.

(5) (a) Ando, W.; Takata, T. *Photooxidation of Sulfur Compounds Singlet Oxygen*; Frimer, A. A., Ed.; CRC Press: Boca Raton, FL, 1985; Vol. III, pp 1–117. (b) Clennan, E. L. *Acc. Chem. Res.* **2001**, *34*, 875–884.

(6) (a) Ho, D. G.; Gao, R.; Celaje, J.; Chung, H.-Y.; Selke, M. *Science* **2003**, *302*, 259–262. (b) Gao, R.; Ho, D. G.; Dong, T.; Khuu, D.; Franco, N.; Sezer, O.; Selke, M. *Org. Lett.* **2001**, *3*, 3719–3722. (c) Tsuji, S.; Kondo, M.; Ishiguro, K.; Sawaki, Y. *J. Org. Chem.* **1993**, *58*, 5055–5059.

[†] Tokai University.

[‡] National Institute of Advanced Industrial Science and Technology (AIST).

(1) (a) Haines, A. H. *Methods for the Oxidation of Organic Compounds*; Academic Press: London, 1988. (b) Sheldon, R. A.; Kochi, J. K. *Metal Catalyzed Oxidation of Organic Compounds*; Academic Press: New York, 1981.

(2) (a) Bäckvall, J.-E. *Modern Oxidation Methods*; Wiley-VCH: Weinheim, 2004. (b) Eissen, M.; Metzger, J. O.; Schmidt, E.; Schneidewind, U. *Angew. Chem., Int. Ed.* **2002**, *41*, 414–436.

reactions.⁷ In this work, we focused on diorganotelluride catalysts. Detty and co-workers reported that singlet oxygen can efficiently oxidize diorganotellurides to give the corresponding telluroxides or their hydrates;⁸ however, there have been no reports concerning their application to organic synthesis.

We have reported recently that photosensitized oxygenation of bulky diaryl tellurides produced corresponding telluroxides.⁹ The diaryl telluroxides obtained were found to oxidize some organic substrates, and in some cases, the starting telluride was recovered quantitatively. These findings helped us to investigate the catalytic ability of diaryl tellurides in photooxygenation reactions. In this work, the catalytic photooxidation of phosphites in the presence of organotellurides is selected as an example to elucidate environmentally benign novel processes.

Initially, we investigated the catalytic photooxidation of triphenyl phosphite under various reaction conditions; the results are compiled in Table 1. When a 0.1 M acetonitrile

Table 1. Aerobic Photooxidation of Triphenyl Phosphite Catalyzed by Diaryl Telluride

$$(\text{PhO})_3\text{P} \xrightarrow[\text{hv, sensitizer, air}]{\text{catalyst (1 mol \%)}} (\text{PhO})_3\text{PO}$$

run	catalyst	solvent (sens) ^a	time (h)	temp (°C)	yield (%) ^b
1	Tip ₂ Te	CH ₃ CN (RB)	4	10–15	quant
2	Tip ₂ Te	CH ₃ CN (RB)	2.5	40–45	quant
3	Tip ₂ Te ^c	CH ₃ CN (RB)	4	40–45	28
4	Tip ₂ Te	C ₅ H ₅ N (HP)	4	10–15	quant
5	Tip ₂ Te	CH ₂ Cl ₂ (TPP)	4	10–15	43
6	Tip ₂ Te	CH ₃ OH (RB)	4	10–15	15
7	Mes ₂ Te	CH ₃ CN (RB)	4	10–15	67
8	An ₂ Te	CH ₃ CN (RB)	4	10–15	14
9	Ph ₂ Te	CH ₃ CN (RB)	4	10–15	24

^a RB, HP, and TPP denote rose bengal, hematoporphyrin, and tetraphenylporphyrin, respectively. ^b Determined by ³¹P NMR spectroscopy. ^c 0.1 mol %.

solution of triphenyl phosphite containing bis(2,4,6-triisopropylphenyl) telluride (Tip₂Te, 1 mol %) was irradiated under aerobic conditions with a 500 W halogen lamp employing rose bengal (RB, 10⁻⁴ M) as a photosensitizer at 10–15 °C for 4 h, analysis of the reaction mixture by ³¹P NMR spectroscopy showed a quantitative formation of triphenyl phosphate (run 1). Similar treatment in the dark, without a sensitizer, or in the absence of Tip₂Te resulted in only a trace amount of phosphate formation or no reaction, indicating that the reaction involves singlet oxygen and that triaryl phosphite itself is completely unreactive toward singlet oxygen.

(7) (a) Krief, A.; Colaux-Castillo, C. *Pure Appl. Chem.* **2002**, *74*, 107–113. (b) Krief, A.; Colaux, C. *Tetrahedron Lett.* **1999**, *40*, 4189–4192. (c) Abatjoglou, A. G.; Bryant, R. D. *Tetrahedron Lett.* **1981**, *22*, 2051–2054.

(8) (a) Sergueievski, P.; Detty, M. R. *Organometallics* **1997**, *16*, 4386–4391. (b) Detty, M. R.; Merkel, P. B. *J. Am. Chem. Soc.* **1990**, *112*, 3845–55.

(9) Oba, M.; Endo, M.; Nishiyama, K.; Ouchi, A.; Ando, W. *Chem. Commun.* **2004**, 1672–1673.

Other diaryl tellurides, such as bis(2,4,6-trimethylphenyl) telluride (Mes₂Te), bis(4-methoxyphenyl) telluride (An₂Te), and diphenyl telluride, were also employed as a catalyst. The catalytic activity of Mes₂Te was moderate, producing triphenyl phosphate in 67% yield (run 7), but An₂Te and Ph₂Te afforded the phosphate only in 14% and 24% yields, respectively (runs 8 and 9). The bulky aryl substituents on the tellurium atom are apparently responsible for increasing the catalytic activity.

A noticeable solvent effect was observed in this reaction. Although the use of pyridine as a solvent produced similar result affording triphenyl phosphate in quantitative yield (run 4), changing the solvent to dichloromethane decreased the reaction rate producing the phosphate in 43% yield after 4 h (run 5). In methanol solvent, the reaction retarded significantly producing triphenyl phosphate in 15% yield (run 6). In all solvents used, the isolated Tip₂TeO, the possible active species in this reaction (vide infra), could efficiently oxidize triphenyl phosphite. Therefore, the observed solvent effect is probably attributed to the efficiency of photosensitized formation of Tip₂TeO. At the present time, we cannot provide a clear explanation of the solvent effect because the mechanism of the singlet oxygen oxidation of Tip₂Te would be very complex. Studies that address these issues are under investigation.

We found finally that carrying out the oxidation at an elevated temperature (40–45 °C) in acetonitrile solvent medium slightly improves the efficiency of the catalyst, and the reaction reached completion within 2.5 h (run 2). Efforts have been made to reduce the amount of catalyst; however, we could not obtain acceptable yields using 0.1 mol % of Tip₂Te (run 3).

To explore the scope and limitation of the present method, the oxidation of a variety of phosphite esters was studied; the representative results are shown in Table 2. In the

Table 2. Tip₂Te-Catalyzed Photooxidation of Phosphite Esters

$$(\text{RO})_3\text{P} \xrightarrow[\text{CH}_3\text{CN, 40-50 }^\circ\text{C, air}]{\text{Tip}_2\text{Te (1 mol \%), hv, rose bengal}} (\text{RO})_3\text{PO}$$

run	(RO) ₃ P	time (h)	yield (%) ^a
1	(PhO) ₃ P	2.5	quant (92) ^b
2	(4-ClC ₆ H ₄ O) ₃ P	1	quant
3	(4-CF ₃ C ₆ H ₄ O) ₃ P	2	quant (94) ^b
4	(4-MeC ₆ H ₄ O) ₃ P	8	quant
5	(4-MeOC ₆ H ₄ O) ₃ P	8	quant (93) ^b
6	(PhO) ₂ (MeO)P	4	91
7	(PhO)(BuO) ₂ P	4	quant


^a Determined by ³¹P NMR spectroscopy. ^b Isolated yield.

oxidation of triaryl phosphites, the presence of an electron-withdrawing *para* substituent, such as a chloro or trifluoromethyl group, resulted in an increase in reactivity producing corresponding phosphates in quantitative yields after 1 or 2 h, respectively (runs 2 and 3). However, an electron-donating methyl or methoxy group required somewhat longer

reaction time (8 h) to attain completion (runs 4 and 5). Mixed alkyl/aryl phosphite esters were also employed as substrates. Although (PhO)₂POMe and (BuO)₂POPh are moderately reactive toward singlet oxygen, Tip₂Te addition improved the reaction efficiency affording the corresponding mixed phosphate esters in almost quantitative yields (runs 6 and 7). In the absence of Tip₂Te, (PhO)₂(MeO)PO and (PhO)(BuO)₂PO were obtained in 30% and 89% yields, respectively.

Various possible active species that may be involved in the transfer of an oxygen atom from singlet oxygen to phosphites are postulated. In particular, peroxytelluroxide (Tip₂Te⁺OO⁻), the initial intermediate perceived to be generated by the oxidation of Tip₂Te through singlet oxygen, may oxidize phosphites. Its sulfur analog, R₂S⁺OO⁻,¹⁰ is known to react with triphenyl phosphite.¹¹ In our case, we could not observe the reaction between Tip₂Te⁺OO⁻ and triphenyl phosphite individually because Tip₂TeO also oxidizes the phosphite. Therefore, we tried to capture Tip₂Te⁺OO⁻ using Ph₂SeO, which is reported to be more reactive toward R₂S⁺OO⁻ than triphenyl phosphite.¹² However, we could not obtain any trapping product such as Ph₂SeO₂, indicating a rather low reactivity of Tip₂Te⁺OO⁻ intermediate. At the present state of this research, we assume that the Tip₂TeO, generated by the photosensitized oxygenation of Tip₂Te, acts as a main oxygen carrier (Scheme 1).

Scheme 1. Catalytic Cycle for Diorganotelluride-Catalyzed Oxidation of Phosphites with Singlet Oxygen



In fact, reaction of triphenyl phosphite with 1 equiv of the isolated Tip₂TeO in acetonitrile at room temperature for 30 min afforded quantitative yields of triphenyl phosphate and Tip₂Te. The regeneration of Tip₂Te is indispensable to complete the catalytic cycle in this process.

To obtain further insights into the reaction mechanism between phosphite and telluroxide, we invoked a conventional Hammett methodology.¹³ Relative reactivities of substituted triaryl phosphites toward Tip₂TeO were determined by competitive reaction using unsubstituted triphenyl phosphite and ³¹P NMR spectroscopy. The plot of the logarithm of relative rate constants, log(k_X/k_H), against Hammett's σ-values exhibits a linear relationship (correlation





Figure 1. Hammett plot for the reaction of substituted triaryl phosphites with Tip₂TeO.

coefficient $R^2 = 0.94$) with a reaction constant $\rho = 2.88$. The large positive slope indicates nucleophilic transfer of oxygen atom from telluroxide to phosphite and significant negative charge accumulation on the phosphorus atom in the transition state of the rate-determining step. Although a rational interpretation of this Hammett correlation in terms of charge distribution in the transition state awaits theoretical verification, we postulate the mechanism for the oxidation of phosphite ester by Tip₂TeO, as shown in Scheme 2. We

Scheme 2. Nucleophilic Oxygen Atom Transfer from Ar₂TeO to (RO)₃P



propose that the attack of a negatively charged oxygen atom on the phosphorus atom generating a zwitterionic intermediate is rate-determining. Similar mechanisms are also proposed by Foote and Stratakis.¹⁴

In summary, we have elucidated the first instance of organotelluride-catalyzed photooxidation of phosphites using molecular oxygen as a terminal oxidant. This catalytic aerobic oxidation is very efficient, and further application of this catalytic system is under investigation.

Acknowledgment. Financial support from Tokai University is gratefully acknowledged.

Supporting Information Available: Experimental procedures and copies of NMR spectra for selected compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL900240S

(14) (a) Stratakis, M.; Rabalagos, C.; Sofikiti, N. *Tetrahedron Lett.* **2003**, *44*, 349–351. (b) Stratakis, M.; Orfanopoulos, M.; Foote, C. S. *Tetrahedron Lett.* **1991**, *32*, 863–866.

(10) (a) Liang, J.-J.; Gu, C.-L.; Kacher, M. L.; Foote, C. S. *J. Am. Chem. Soc.* **1983**, *105*, 4717–4721. (b) Jensen, F.; Greer, A.; Clennan, E. L. *J. Am. Chem. Soc.* **1998**, *120*, 4439–4449.

(11) (a) Clennan, E. L.; Stensaas, K. L.; Rupert, S. D. *Heteroat. Chem.* **1998**, *9*, 51–56. (b) Nahm, K.; Foote, C. S. *J. Am. Chem. Soc.* **1989**, *111*, 1909–1910.

(12) Sofikiti, N.; Rabalagos, C.; Stratakis, M. *Tetrahedron Lett.* **2004**, *45*, 1335–1337.

(13) Hansch, C.; Leo, A.; Taft, R. W. *Chem. Rev.* **1991**, *91*, 165–195.