

Organic & Biomolecular Chemistry

Accepted Manuscript



This article can be cited before page numbers have been issued, to do this please use: T. Hatanaka, R. Yuki, R. Saito and K. SASAKI, *Org. Biomol. Chem.*, 2016, DOI: 10.1039/C6OB02256A.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this *Accepted Manuscript* with the edited and formatted *Advance Article* as soon as it is available.

You can find more information about *Accepted Manuscripts* in the [Information for Authors](#).

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard [Terms & Conditions](#) and the [Ethical guidelines](#) still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.

Journal Name

COMMUNICATION

 α -Methylphenacyl thioesters as convenient thioacid precursorsToru Hatanaka,^a Ryosuke Yuki,^a Ryota Saito^{a,b} and Kaname Sasaki^{*a}Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x


www.rsc.org/

α -Methylphenacyl (Mpa) thioesters are described as precursors of thioacids. Mpa thioesters are accessible via the condensation of carboxylic acids and phenacyl thiol, which is easily prepared without column chromatography. The Mpa thioesters are selectively deprotected by reduction with zinc dust in the presence of conventional thioacid protecting groups. In addition, the Mpa group exhibits orthogonal reactivity to the Boc group. These features are expected to facilitate the preparation of complex thioacids, including those in peptides.

The thioacid functional group, due to its unique reactivity, can be an expedient handle by which to convergently and chemoselectively install modification modules, particularly in peptides.¹ For example, thioacid-based amide ligation reactions relying on unprotected aziridines,² electron-deficient azides,³ electron-deficient arylsulfonamides,⁴ isonitriles,⁵ and isocyanates⁶ have been reported. Furthermore, condensation reactions between thioacids and amines have also been described recently, employing the Sanger and Mukaiyama reagents,⁷ HOBT,⁸ organonitrite,⁹ Cu(II),¹⁰ HOBT-Cu(OAc)₂,¹¹ and *N,O*-bis(trimethylsilyl)acetamide as activators.¹²

A promising approach to preparing peptide thioacids is to carry thioesters through the peptide elongation sequence and then release the thioacids by reliable and simple reactions. Currently, several thioesters are available as thioacid precursors, including 2,4,6-trimethoxybenzyl (Tmob),³ 9-fluorenylmethyl (Fm),¹³ trityl (Trt),¹⁴ and 2-cyanoethyl,¹⁵ which release thioacids under either basic or acidic conditions. As for C-terminal peptide thioacids, another approaches were reported, which utilize *N-S* acyl transfer¹⁶⁻¹⁸ or acyl hydrazines.¹⁹ We were interested in a different type of precursor that would permit a wider range of applications for thioacid chemistry, and herein, report a novel thioacid

precursor that has, notably, reactivity orthogonal to the Boc group. As thioesters are not generally stable to nucleophilic amines such as piperidine, Boc-based elongation would be the first choice for preparing thioacid-bearing peptides. Therefore, thioacid precursors with reactivity orthogonal to the Boc group could be useful for the preparation of peptide thioacids. In this context, we focused on phenacyl groups. As a protective group for carboxylic acids, phenacyl esters have been used due to their stability under acidic conditions and removal by one-electron reduction in mildly acidic media.²⁰ However, phenacyl esters are known to be relatively electrophilic and susceptible to undesired transesterification. Therefore, we prepared phenacyl thiols **3a–c**, which vary in bulkiness (Scheme 1), to examine the stabilities of their thioesters. Each variant was prepared from the commercially available phenacyl bromides **1a–c** by a two-step sequence of substitution with thioacetate to give **2a–c**, followed by deacetylation. These are all easily scalable spot-to-spot reactions, and afford phenacylthiol **3a**, α -methylphenacylthiol **3b**, and α,α -dimethylphenacylthiol **3c** quantitatively, without the need for column chromatography.

Scheme 1. Preparation of thiols **3a–c** and thioesters **4a–c**.

^aDepartment of Chemistry, Toho University, 2-2-1 Miyama, Funabashi 274-8510 Japan. ^bResearch Center for Materials with Integrated Properties, Toho University, 2-2-1 Miyama, Funabashi, Chiba 274-8510, Japan. *Email: kaname.sasaki@sci.toho-u.ac.jp


Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x

COMMUNICATION

Journal Name

Each thiol was efficiently coupled with Boc-protected tryptophan using standard coupling procedures in the presence of DMAP.²¹ The lower yield in the preparation of **4a** is due to the partial loss of **3a** observed during silica gel column chromatography. Thioesters **4a–c** were subjected to reducing conditions using zinc dust in aqueous acetic acid as a benchmark (Table 1). The efficiency in liberating the thioacid was evaluated after conversion to the corresponding benzyl thioester **5**, because thioacids generally have two tautomeric forms (*O*- and *S*-acids) that complicate spectroscopic analysis.²² All three substrates were deprotected to give the corresponding thioacid. The reaction rate clearly depended on the bulkiness of the phenacyl moiety. The bulkier thioester **4c** required a longer reaction time than **4a** and **4b**, even though the intermediate phenacyl radical of **4c** would presumably be more stable than those of the others. Elevating the reaction temperature shortened the reaction time and improved the yield. We considered the α -methylphenacyl group, abbreviated as Mpa, to be a good entry as a precursor for thioacids.²³

Table 1. Deprotection of phenacyl thioesters **4a–c**.

				
Entry	Thioester 4a–c	Temp./ °C	Time	Yield of 5 / % ^a
1	4a	25	1 h	92
2	4a	40	20 min	88
3	4b	25	3 h	86
4	4b	40	1 h	86
5	4c	25	26 h	50
6	4c	40	6 h	83


^aIsolated yields.

Table 2 summarizes the scope of this thioacid preparation procedure. Each of the aromatic, aliphatic, and amino acids **6a–i** was converted to the Mpa thioester **7a–i** using EDCI and DMAP in excellent isolated yields. These precursors afforded the corresponding thioacids smoothly when treated with zinc at ambient temperature. Excess zinc dust was removed by filtration through a silica gel pad, and then the solvent was evaporated to effectively give the thioacid, according to the yield after benzylation. It is noteworthy that no diastereomeric benzyl thioester of **8f** was observed, which excludes the possibility of racemization in the reaction sequence. The present transformation was also applicable to Fmoc-protected amino acid **6g**. Furthermore, carboxylic acids having additional thioester groups such as **6h** and **6i** could be converted to the bithioesters, and the Mpa group could be removed in the presence of either an Fm- or Tmob-thioester. The thioacids could be also captured by 3-bromopropionitrile instead of benzyl bromide to give the 2-cyanoethyl thioester **8i'**.

View Article Online


DOI: 10.1039/C6OB02256A

Table 2. Substrate scope of thioacid preparation via Mpa thioesters.

			
Entry	R ² -COOH	Yield of 7a–i / % ^a	Yield of 8a–i / % ^a
1	PhCOOH (6a)	90	80
2	ⁿ C ₉ H ₁₉ COOH (6b)	92	91
3	Boc-Phe-OH (6c)	94	83
4	Boc-Gly-OH (6d)	95	83
5	Boc-Ser(OBn)-OH (6e)	98	78
6	Boc-Thr(OBn)-OH (6f)	94	73
7	Fmoc-Trp-OH (6g)	98	85
8	Boc-Glu(SFm)-OH (6h)	74	78
9	Fmoc-Glu(STmob)-OH (6i)	77	81
10	6i	-	77 ^b


^aIsolated yields. ^b3-Bromopropionitrile was used instead of BnBr, which afforded 2-cyanoethyl thioester **8i'**.

Next, we examined the orthogonality of the Mpa thioesters and Boc groups. As shown in Scheme 2, Boc-protected thioglycinate **7d** was employed as the C-terminus, and peptide elongation based on Boc-TFA chemistry was conducted. In the presence of the Mpa thioester, the Boc group was successfully deprotected in 40% TFA/CH₂Cl₂. The subsequent coupling reaction afforded dipeptidyl thioester **9**, where NMM performed better than DIPEA as a base. Another deprotection-coupling sequence was performed with lysine having a Boc group at the ω -amine to give tripeptide **10** without any problems. In the presence of the ω -Boc group, Mpa was cleanly removed using zinc at 40 °C to afford the peptidyl thioacid in good yield, which was confirmed by derivatization to the benzyl thioester **11**.



Scheme 2. Synthesis of oligopeptide α-thioacid.

Finally, we prepared ω-Mpa thioglutamate **16** from the known glutamate **12** by protecting group manipulation, *i.e.*, allylation of the α-carboxylic acid, deprotection of the ω-9-fluorenylmethyl ester, thioesterification using MpaSH, and deallylation using a Pd catalyst. This glutamate residue could be incorporated into a peptide by Boc-TFA chemistry without loss of the Mpa group to afford tripeptide **18**. Again, the Mpa group could be removed in the presence of the Boc group, and benzyl thioester **19** was obtained after treatment with benzyl bromide. This orthogonality permits the preparation of peptidyl thioacids at either the C-terminus or the side-chain.



Scheme 3. Synthesis of oligopeptide having thioacid on its side-chain.

Conclusions

α-Methylphenacyl (Mpa) thioesters were developed as precursors for thioacids. MpaSH is easily accessible from commercially available, low-cost materials and readily introduced to carboxylic acids. Mpa thioesters can facilitate the preparation of peptides bearing a thioacid either at the C-terminus or on the side chain. The orthogonal reactivity to the Boc-group and chemoselectivity over Tmob- and Fm-thioesters are expected to enable the preparation of complex thioacids such as peptide thioacids. We are now applying this chemistry to the preparation of oligo- and polypeptides.

Acknowledgments

This work was financially in part supported by a JSPS Grant-in-Aid for Young Scientists B (JP26810062) to KS and the MEXT-

Supported Program for the Strategic Research Foundation at Private Universities (2012-2016) to RS. We also thank Prof. Hidemasa Hikawa, Toho University, for help with FABHRMS experiments.

View Article Online
DOI: 10.1039/C6OB02256A

Notes and references

1. K. Sasaki, S. Aubry and D. Crich, *Phosphorus, Sulfur Silicon Relat. Elem.*, 2011, **186**, 1005-1018.
2. N. Assem, A. Natarajan and A. K. Yudin, *J. Am. Chem. Soc.*, 2010, **132**, 10986-10987.
3. N. Shangguan, S. Katukojvala, R. Greenberg and L. J. Williams, *J. Am. Chem. Soc.*, 2003, **125**, 7754-7755.
4. D. Crich, K. Sasaki, M. Y. Rahaman and A. A. Bowers, *J. Org. Chem.*, 2009, **74**, 3886-3893.
5. Y. Rao, X. Li and S. J. Danishefsky, *J. Am. Chem. Soc.*, 2009, **131**, 12924-12926.
6. D. Crich and K. Sasaki, *Org. Lett.*, 2009, **11**, 3514-3517.
7. D. Crich and I. Sharma, *Angew. Chem. Int. Ed.*, 2009, **48**, 2355-2358.
8. P. Wang and S. J. Danishefsky, *J. Am. Chem. Soc.*, 2010, **132**, 17045-17051.
9. J. Pan, N. O. Devarie-Baez and M. Xian, *Org. Lett.*, 2011, **13**, 1092-1094.
10. S. M. Mali, S. V. Jadhav and H. N. Gopi, *Chem. Commun.*, 2012, **48**, 7085-7087.
11. R. Joseph, F. B. Dyer and P. Garner, *Org. Lett.*, 2013, **15**, 732-735.
12. W. Wu, Z. Zhang and L. S. Liebeskind, *J. Am. Chem. Soc.*, 2011, **133**, 14256-14259.
13. D. Crich, K. Sana and S. Guo, *Org. Lett.*, 2007, **9**, 4423-4426.
14. D. Crich and I. Sharma, *Angew. Chem. Int. Ed.*, 2009, **48**, 7591-7594.
15. R. Raz and J. Rademann, *Org. Lett.*, 2012, **14**, 5038-5041.
16. A. Shigenaga, Y. Sumikawa, S. Tsuda, K. Sato and A. Otaka, *Tetrahedron*, 2010, **66**, 3290-3296.
17. S. L. Pira, E. Boll and O. Melnyk, *Org. Lett.*, 2013, **15**, 5346-5349.
18. T. Shimizu, R. Miyajima, K. Sato, K. Sakamoto, N. Naruse, M. Kita, A. Shigenaga and A. Otaka, *Tetrahedron*, 2016, **72**, 992-998.
19. C. Chen, Y. Huang, L. Xu, Y. Zheng, H. Xu, Q. Guo, C. Tian, Y. Li and J. Shi, *Org. Biomol. Chem.*, 2014, **12**, 9413-9418.
20. P. G. M. Wuts and T. W. Greene, in *Protective groups in organic synthesis*, John Wiley & Sons, Hoboken, New Jersey, 4th edition edn., 2007, pp. 568-569.
21. B. Neises and W. Steglich, *Angew. Chem. Int. Ed. Engl.*, 1978, **17**, 522-524.
22. S. Kato, Y. Kawahara, H. Kageyama, R. Yamada, O. Niyomura, T. Murai and T. Kanda, *J. Am. Chem. Soc.*, 1996, **118**, 1262-1267.
23. In the reaction of **4b**, conversion to the corresponding thioacid was varified by ESIHRMS; calcd. for C₁₆H₁₉N₂O₃S⁻ [M - H]⁻ 319.1122, found 319.1103.