

C(10)	0.571 (1)	0.0999 (4)	0.8924 (3)	0.0651
C(11)	0.884 (1)	0.1618 (4)	0.7123 (3)	0.0540
C(12)	0.548 (1)	0.1673 (4)	1.0241 (3)	0.0629
C(13)	0.879 (1)	0.0753 (4)	0.9941 (3)	0.0656
C(14)	0.839 (1)	0.1431 (5)	1.1203 (3)	0.0736
C(15)	0.885 (1)	0.1744 (5)	0.5705 (3)	0.0702
C(16)	1.056 (2)	0.1004 (5)	0.5955 (4)	0.0799
C(17)	0.967 (1)	0.0719 (4)	0.6738 (3)	0.0714
C(18)	0.965 (1)	0.0898 (5)	1.0687 (4)	0.0743
C(19)	0.633 (1)	0.1811 (4)	1.0985 (3)	0.0719

Larson, A. C. (1970). *Crystallographic Computing*, edited by F. R. Ahmed, S. R. Hall & C. P. Huber, pp. 291–294. Copenhagen: Munksgaard.

Paquette, L. A., Lau, C. J. & Rogers, R. D. (1988). *J. Am. Chem. Soc.* **110**, 2592–2600.

Walker, N. & Stuart, D. (1983). *Acta Cryst.* **A39**, 158–166.

Watkin, D. J., Carruthers, J. R. & Betteridge, P. W. (1985). *CRYSTALS Users Guide*. Chemical Crystallography Laboratory, Univ. of Oxford, England.

Table 2. Selected geometric parameters (Å, °)

S(1)—Cl(2)	2.046 (2)	C(8)—C(12)	1.368 (7)
S(1)—N(4)	1.581 (4)	C(8)—C(13)	1.378 (8)
S(1)—O(5)	1.409 (4)	C(9)—C(11)	1.514 (8)
S(1)—O(7)	1.415 (4)	C(11)—C(17)	1.529 (8)
O(3)—C(9)	1.318 (6)	C(12)—C(19)	1.383 (8)
O(3)—C(10)	1.462 (6)	C(13)—C(18)	1.389 (8)
N(4)—C(11)	1.497 (6)	C(14)—C(18)	1.373 (9)
N(4)—C(15)	1.488 (6)	C(14)—C(19)	1.35 (1)
O(6)—C(9)	1.206 (7)	C(15)—C(16)	1.51 (1)
C(8)—C(10)	1.489 (7)	C(16)—C(17)	1.498 (9)
N(4)—S(1)—Cl(2)	104.9 (2)	C(11)—C(9)—O(3)	109.6 (5)
O(5)—S(1)—Cl(2)	105.1 (2)	C(11)—C(9)—O(6)	125.3 (5)
O(5)—S(1)—N(4)	109.2 (2)	C(8)—C(10)—O(3)	107.9 (5)
O(7)—S(1)—Cl(2)	105.1 (2)	C(9)—C(11)—N(4)	110.4 (4)
O(7)—S(1)—N(4)	109.6 (2)	C(17)—C(11)—N(4)	102.7 (4)
O(7)—S(1)—O(5)	121.6 (3)	C(17)—C(11)—C(9)	111.7 (5)
C(10)—O(3)—C(9)	116.7 (5)	C(19)—C(12)—C(8)	121.0 (6)
C(11)—N(4)—S(1)	118.0 (3)	C(18)—C(13)—C(8)	120.3 (5)
C(15)—N(4)—S(1)	120.1 (4)	C(19)—C(14)—C(18)	119.9 (6)
C(15)—N(4)—C(11)	110.1 (4)	C(16)—C(15)—N(4)	104.3 (4)
C(12)—C(8)—C(10)	119.6 (5)	C(17)—C(16)—C(15)	103.3 (6)
C(13)—C(8)—C(10)	121.9 (5)	C(16)—C(17)—C(11)	105.3 (5)
C(13)—C(8)—C(12)	118.6 (5)	C(14)—C(18)—C(13)	119.9 (6)
O(6)—C(9)—O(3)	125.0 (5)	C(14)—C(19)—C(12)	120.3 (6)

Analysis of the intensity data revealed that the systematic absences were consistent with the space group $P2_12_12_1$. Corrections were applied for Lorentz and polarization effects. The structure was solved using the direct methods program *SIR88* (Burla, Camalli, Cascarano, Giacovazzo, Polidori, Spagna & Viterbo, 1989), which revealed the positions of all the non-H atoms. All other computations were performed using the *CRYSTALS* program (Watkin, Carruthers & Betteridge, 1985). The H atoms were subsequently placed geometrically during the refinement process. Full-matrix least-squares refinement was used for the positions and isotropic temperature factors of all the non H-atoms; a parameter to allow for the effects of secondary extinction was included.

Lists of structure factors, anisotropic displacement parameters and H-atom coordinates have been deposited with the IUCr (Reference: HE1006). Copies may be obtained through The Managing Editor, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

References

- Barton, T. J., Rogido, R. & Clardy, J. C. (1970). *Tetrahedron Lett.* pp. 2081–2084.
- Burla, M. C., Camalli, M., Cascarano, G., Giacovazzo, C., Polidori, G., Spagna, R. & Viterbo, D. (1989). *J. Appl. Cryst.* **22**, 389–393.
- Furst, G. T., Wachsman, M. A., Pieroni, J., White, J. G. & Moriconi, E. J. (1973). *Tetrahedron*, **29**, 1675–1677.

Acta Cryst. (1994). **C50**, 1284–1286

2,6-Bis(*p*-nitrophenylthiomethyl)pyridine

R. SILLANPÄÄ

Department of Chemistry, University of Turku,
FIN-20500 Turku, Finland

R. KIVEKÄS

Department of Chemistry, University of Helsinki,
PO Box 6, FIN-00014, Finland

L. ESCRICHE

Departament de Química, Universitat Autònoma
de Barcelona, Campus de Bellaterra,
08193 Bellaterra, Barcelona, Spain

G. SÁNCHEZ-CASTELLÓ AND F. TEIXIDOR

Institut de Ciència de Materials, CSIC, Campus de
Bellaterra, 08193 Bellaterra, Barcelona, Spain

(Received 17 June 1993; accepted 2 December 1993)


Abstract

In the title compound, C₁₉H₁₅N₃O₄S₂, the pyridine ring, one thiomethyl group and the adjoining nitrobenzene group are approximately in the same plane. The C—C—S—C torsion angle is $-179.9(3)^\circ$. The C atom of the second thiomethyl group is also in this plane, but the S(2) atom is *anti* to the plane resulting in a C—C—S—C torsion angle of $67.7(4)^\circ$.

Comment

As a part of our study on metal complexes of *S,S'*-alkyl and *S,S'*-aryl substituted 2,6-bis(thiomethyl)pyridine derivatives (Teixidor, Sánchez-Castelló, Lucena, Escriche, Kivekäs, Sundberg & Casabó, 1991), we now report the crystal structure of 2,6-bis(*p*-nitrophenylthiomethyl)pyridine, (I). This type of ligand is a source of an NS₂-coordinating

moiety. A drawing of the molecule is shown in Fig. 1.



The solid-state conformation of the title ligand is not very suitable for tridentate coordination as a result of the axial orientation of the S(2) atom with respect to the pyridine ring. The nitro group N(2)O(1)O(2) is only slightly (3.5°) twisted with respect to the neighbouring benzene group, but the corresponding angle for the other nitro group is 9.1° .




Fig. 1. ORTEP (Johnson, 1976) plot of the title compound. Displacement ellipsoids are shown at 30% probability levels, except for H atoms which are drawn with isotropic temperature factors of $1.0 \text{ \AA}^2$.

Experimental

NaOH 98% (1.80 g, 45.0 mmol) in ethanol (200 cm^3) was stirred under reflux for 30 min with *p*-nitrothiophenol (7.00 g, 45.0 mmol). This solution was added to a solution of 2,6-bis(bromomethyl)pyridine (5.38 g, 20.0 mmol) in ethanol (100 cm^3) and was stirred for 1 h at 273 K. A deep yellow precipitate appeared which was filtered off and washed twice with water (30 cm^3). The resulting residue was dissolved in thf (100 cm^3), dried (Na_2SO_4) and evaporated under vacuum to obtain an orange solid. Yield: 56% (4.86 g). Analysis: calculated for $\text{C}_{19}\text{H}_{15}\text{N}_3\text{O}_4\text{S}_2$, C 55.2, H 3.6, N 10.2, S 15.5%; found, C 55.3, H 3.8, N 9.9, S 15.3%. ^1H NMR (CDCl_3): δ 4.52 (s, 4, py— CH_2 —S), δ 7.77 (m, 11, aromatic). Slow evaporation of a methanolic solution of this compound

afforded slightly yellow crystals suitable for X-ray diffraction analysis. Dehydrated and deoxygenated ethanol was used in the synthesis. 2,6-Bis(bromomethyl)pyridine was prepared according to a reported procedure (Offerman & Vögtle, 1977). *p*-Nitrothiophenol is commercially available (Aldrich) and was used as received. Microanalyses were performed on a Perkin–Elmer 240-B instrument. Proton NMR spectra were recorded on a Bruker 400 MHz AC instrument.

Crystal data

$\text{C}_{19}\text{H}_{15}\text{N}_3\text{O}_4\text{S}_2$

$M_r = 413.47$

Triclinic

$P\bar{1}$

$a = 10.779 (2) \text{ \AA}$

$b = 12.241 (2) \text{ \AA}$

$c = 7.540 (2) \text{ \AA}$

$\alpha = 98.50 (2)^\circ$

$\beta = 103.95 (2)^\circ$

$\gamma = 74.24 (1)^\circ$

$V = 925.6 (7) \text{ \AA}^3$

$Z = 2$

$D_x = 1.483 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation

$\lambda = 0.71069 \text{ \AA}$

Cell parameters from 25 reflections

$\theta = 16\text{--}20^\circ$

$\mu = 0.305 \text{ mm}^{-1}$

$T = 296 \text{ K}$

Needle

$0.20 \times 0.20 \times 0.15 \text{ mm}$

Yellow

Data collection

Rigaku AFC-5S diffractometer

ω – 2θ scans

Absorption correction: none

3443 measured reflections

3254 independent reflections

1963 observed reflections

$[I > \sigma(I)]$

$R_{\text{int}} = 0.039$

$\theta_{\text{max}} = 25^\circ$

$h = 0 \rightarrow 14$

$k = -16 \rightarrow 16$

$l = -10 \rightarrow 10$

3 standard reflections

monitored every 150

reflections

intensity variation: none

Refinement

Refinement on F

$R = 0.051$

$wR = 0.046$

$S = 1.25$

1963 reflections

298 parameters

Only coordinates of H atoms refined

$w = 1/\sigma^2(F)$

$(\Delta/\sigma)_{\text{max}} = 0.03$

$\Delta\rho_{\text{max}} = 0.23 \text{ e \AA}^{-3}$

$\Delta\rho_{\text{min}} = -0.29 \text{ e \AA}^{-3}$

Atomic scattering factors

from *International Tables*

for X-ray Crystallography

(1974, Vol. IV)

Table 1. Fractional atomic coordinates and equivalent isotropic displacement parameters ($\text{\AA}^2$)

	$B_{\text{eq}} = (4/3)\sum_i \sum_j \beta_{ij} a_i \cdot a_j$		
	<i>x</i>	<i>y</i>	<i>z</i>
S(1)	0.4396 (1)	0.42936 (9)	0.6825 (2)
S(2)	0.8119 (1)	0.02513 (9)	0.9996 (2)
O(1)	−0.0517 (3)	0.8816 (3)	0.4391 (6)
O(2)	−0.1363 (3)	0.7469 (3)	0.2914 (5)
O(3)	0.2253 (3)	0.2726 (3)	1.2039 (5)
O(4)	0.3084 (4)	0.1573 (4)	1.4140 (6)
N(1)	0.6938 (3)	0.3101 (3)	0.8209 (4)
N(2)	−0.0470 (3)	0.7810 (4)	0.3957 (6)
N(3)	0.3146 (4)	0.1988 (4)	1.2794 (7)
C(1)	0.6904 (4)	0.4204 (3)	0.8705 (5)
C(2)	0.7972 (4)	0.4568 (4)	0.9788 (6)
C(3)	0.9118 (4)	0.3770 (4)	1.0386 (6)
C(4)	0.9165 (4)	0.2625 (4)	0.9885 (6)
C(5)	0.8065 (4)	0.2326 (3)	0.8800 (5)

C(6)	0.5635 (4)	0.5047 (3)	0.7990 (6)	3.5 (2)
C(7)	0.3012 (3)	0.5409 (3)	0.6098 (5)	3.0 (2)
C(8)	0.1924 (4)	0.5060 (4)	0.5007 (6)	3.9 (2)
C(9)	0.0795 (4)	0.5840 (4)	0.4312 (6)	4.2 (2)
C(10)	0.0725 (4)	0.6980 (3)	0.4733 (5)	3.4 (2)
C(11)	0.1774 (4)	0.7345 (3)	0.5831 (6)	3.9 (2)
C(12)	0.2918 (4)	0.6561 (3)	0.6504 (6)	3.5 (2)
C(13)	0.8038 (4)	0.1100 (4)	0.8184 (7)	4.4 (2)
C(14)	0.6616 (4)	0.0815 (3)	1.0709 (6)	3.6 (2)
C(15)	0.6553 (4)	0.0481 (4)	1.2361 (7)	4.2 (2)
C(16)	0.5433 (5)	0.0860 (4)	1.3059 (7)	4.6 (3)
C(17)	0.4350 (4)	0.1575 (3)	1.2068 (7)	4.0 (2)
C(18)	0.4365 (4)	0.1907 (3)	1.0414 (7)	4.1 (2)
C(19)	0.5499 (4)	0.1530 (3)	0.9718 (6)	3.8 (2)

Table 2. Selected geometric parameters (Å, °)

S(1)—C(6)	1.809 (5)	C(3)—C(4)	1.386 (6)
S(1)—C(7)	1.764 (3)	C(4)—C(5)	1.374 (6)
S(2)—C(13)	1.808 (6)	C(5)—C(13)	1.510 (6)
S(2)—C(14)	1.759 (4)	C(7)—C(8)	1.392 (6)
O(1)—N(2)	1.218 (6)	C(7)—C(12)	1.379 (6)
O(2)—N(2)	1.215 (5)	C(8)—C(9)	1.370 (5)
O(3)—N(3)	1.217 (5)	C(9)—C(10)	1.370 (6)
O(4)—N(3)	1.223 (8)	C(10)—C(11)	1.370 (6)
N(1)—C(1)	1.340 (5)	C(11)—C(12)	1.378 (5)
N(1)—C(5)	1.346 (4)	C(14)—C(15)	1.389 (7)
N(2)—C(10)	1.462 (5)	C(14)—C(19)	1.401 (5)
N(3)—C(17)	1.464 (7)	C(15)—C(16)	1.371 (7)
C(1)—C(2)	1.381 (6)	C(16)—C(17)	1.380 (6)
C(1)—C(6)	1.507 (5)	C(17)—C(18)	1.373 (8)
C(2)—C(3)	1.375 (5)	C(18)—C(19)	1.384 (7)
C(6)—S(1)—C(7)	102.7 (2)	S(1)—C(7)—C(12)	126.6 (3)
C(13)—S(2)—C(14)	104.9 (2)	C(8)—C(7)—C(12)	118.5 (3)
C(1)—N(1)—C(5)	117.9 (3)	C(7)—C(8)—C(9)	121.0 (4)
O(1)—N(2)—O(2)	123.1 (4)	C(8)—C(9)—C(10)	119.5 (4)
O(1)—N(2)—C(10)	117.9 (4)	N(2)—C(10)—C(9)	119.3 (4)
O(2)—N(2)—C(10)	119.0 (4)	N(2)—C(10)—C(11)	120.0 (4)
O(3)—N(3)—O(4)	122.9 (5)	C(9)—C(10)—C(11)	120.7 (3)
O(3)—N(3)—C(17)	118.9 (5)	C(10)—C(11)—C(12)	119.9 (4)
O(4)—N(3)—C(17)	118.2 (4)	C(7)—C(12)—C(11)	120.5 (4)
N(1)—C(1)—C(2)	122.7 (3)	S(2)—C(13)—C(5)	114.8 (3)
N(1)—C(1)—C(6)	116.4 (3)	S(2)—C(14)—C(15)	115.9 (3)
C(2)—C(1)—C(6)	120.9 (4)	S(2)—C(14)—C(19)	125.0 (4)
C(1)—C(2)—C(3)	118.9 (4)	C(15)—C(14)—C(19)	119.1 (4)
C(2)—C(3)—C(4)	119.0 (4)	C(14)—C(15)—C(16)	121.4 (4)
C(3)—C(4)—C(5)	118.9 (3)	C(15)—C(16)—C(17)	118.5 (5)
N(1)—C(5)—C(4)	122.6 (4)	N(3)—C(17)—C(16)	119.2 (5)
N(1)—C(5)—C(13)	115.2 (3)	N(3)—C(17)—C(18)	118.9 (4)
C(4)—C(5)—C(13)	122.1 (3)	C(16)—C(17)—C(18)	121.9 (4)
S(1)—C(6)—C(1)	109.6 (3)	C(17)—C(18)—C(19)	119.6 (4)
S(1)—C(7)—C(8)	114.9 (3)	C(14)—C(19)—C(18)	119.6 (5)
S(1)—C(6)—C(1)—N(1)	6.8 (5)		
S(1)—C(6)—C(1)—C(2)	−174.3 (3)		
S(2)—C(13)—C(5)—N(1)	−112.0 (4)		
S(2)—C(13)—C(5)—C(4)	68.1 (5)		
O(1)—N(2)—C(10)—C(9)	178.2 (4)		
O(1)—N(2)—C(10)—C(11)	−3.4 (6)		
O(2)—N(2)—C(10)—C(9)	−2.0 (6)		
O(2)—N(2)—C(10)—C(11)	176.4 (4)		
O(3)—N(3)—C(17)—C(16)	170.9 (4)		
O(3)—N(3)—C(17)—C(18)	−9.3 (6)		
O(4)—N(3)—C(17)—C(16)	−9.0 (7)		
O(4)—N(3)—C(17)—C(18)	170.7 (5)		
C(1)—C(6)—S(1)—C(7)	−179.9 (3)		
C(5)—C(13)—S(2)—C(14)	67.7 (4)		
C(6)—S(1)—C(7)—C(8)	174.8 (3)		
C(6)—S(1)—C(7)—C(12)	−5.2 (4)		
C(13)—S(2)—C(14)—C(15)	−166.0 (3)		
C(13)—S(2)—C(14)—C(19)	16.0 (4)		

($1.2 \times B_{\text{eq}}$ of the carrying atom). The calculations were performed with *TEXSAN* (Molecular Structure Corporation, 1989) software using a VAXstation 3520 computer. The figures were drawn using *ORTEPII* (Johnson, 1976).

RK thanks SUOMEN KULTTUURIRAHASTO for the grant.

Lists of structure factors, anisotropic displacement parameters, H-atom coordinates and bond distances involving H atoms have been deposited with the IUCr (Reference: AB1100). Copies may be obtained through The Managing Editor, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

References

- Gilmore, C. J. (1984). *J. Appl. Cryst.* **17**, 42–46.
 Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
 Molecular Structure Corporation (1989). *TEXSAN. TEXRAY Single Crystal Structure Analysis Software*. Version 5.0. MSC, 3200 Research Forest Drive, The Woodlands, TX 77381, USA.
 Offerman, W. & Vögtle, F. (1977). *Synthesis*, pp. 272–273.
 Teixidor, F., Sánchez-Castelló, G., Lucena, N., Escriche, L., Kivekäs, R., Sundberg, M. & Casabó, J. (1991). *Inorg. Chem.* **30**, 4931–4935.

Acta Cryst. (1994). **C50**, 1286–1288

2-(*o*-Methoxyphenoxy)- 1-methylbenzimidazole, C₁₅H₁₄N₂O₂

K. T. VASUDEVAN AND PUTTARAJA

*Department of Physics, Bangalore University,
Bangalore 560056, India*

AND M. V. KULKARNI

*Department of Chemistry, Central College,
Bangalore University, Bangalore 560001, India*

(Received 8 March 1993; accepted 5 October 1993)

Abstract

The benzimidazole ring is planar and makes a dihedral angle of 79.02 (2)° with the aryloxy ring. The N—CH₃ and O—CH₃ groups are *anti* to each other. The molecules are held together in the crystal by van der Waals interactions.

Comment

Several 1-dialkylaminoalkyl-2-aryloxybenzimidazoles have been found to exhibit excellent muscle-relaxant

The structure was solved by direct methods (Gilmore, 1984) and successive Fourier syntheses. Refinement was performed with full-matrix least-squares methods, with non-H atoms anisotropic and H atoms with fixed displacement parameters