

The Coordination of Small Molecules by Manganese(II) Phosphine Complexes. Part 13*. The Synthesis and Characterisation of some Novel Manganese(II) Complexes of Tertiary Butylisocyanide, $\text{MnX}_2(\text{CNBu}^t)$ ($\text{X} = \text{Cl}, \text{Br}, \text{I}, \text{NCS}$), and their Reaction with Tri(*n*-butyl)phosphine and Molecular Oxygen

CHRISTOPHER G. BENSON[†], CHARLES A. McAULIFFE^{**}, ANTHONY G. MACKIE and S. PETER TANNER[†]

Department of Chemistry, University of Manchester Institute of Science and Technology, Manchester M60 1QD, U.K.

(Received September 11, 1986)

Abstract

Novel tertiary butylisocyanide complexes of manganese(II) with unusual stoichiometry, $\text{MnX}_2(\text{CNBu}^t)$ ($\text{X} = \text{Cl}, \text{Br}, \text{I}, \text{NCS}$), have been prepared and characterised by magnetic susceptibility measurements and infrared and electron spin resonance spectroscopy. For $\text{X} = \text{Cl}, \text{Br}, \text{I}$ the complexes have been assigned a polymeric structure similar to that found for $\text{MnI}_2(\text{PPhMe}_2)$, whilst $\text{Mn}(\text{NCS})_2(\text{CNBu}^t)$ is probably dimeric as the infrared spectrum exhibits bands assignable to both bridging and terminal thiocyanate groups. The $\text{MnX}_2(\text{CNBu}^t)$ show no ability to reversibly bind dioxygen, but in solution in tetrahydrofuran PBu_3^n displaces the isocyanide ligand and for $\text{X} = \text{Cl}, \text{Br}, \text{I}$ these solutions absorb dioxygen in a 1:1 Mn:O₂ ratio.

Introduction

We have reported our investigations of manganese(II) phosphine complexes, $\text{MnX}_2(\text{PR}_3)$ ($\text{X} = \text{Cl}, \text{Br}, \text{I}, \text{NCS}$), which can reversibly coordinate small molecules such as dioxygen [1–4], carbon monoxide [5], nitric oxide [6], and ethylene [7] and irreversibly coordinate carbon disulphide [8], sulphur dioxide [9] and tetracyanoethylene [10].

We have extended these studies to manganese(II) complexes containing ligands other than tertiary phosphines. For example, we have shown that the $\text{Mn}(\text{OPPh}_3)_4\text{I}_2$ complex can undergo an insertion reaction with sulphur dioxide to form $[\text{Mn}(\text{OPPh}_3)_4(\text{OSOI})_2]$, and half of the sulphur dioxide may be desorbed by heating [11]. We now wish to report our studies of manganese(II)/isocyanide complexes.

The isocyanide ligand in transition metal complexes can be formally regarded as an analogue of the CO ligand. Manganese(II) complexes of isocyanides of general formula $[\text{Mn}(\text{CNR})_6]\text{X}_2$ ($\text{R} = \text{Ph}, \text{Et}, \text{Bu}^t, \text{CH}=\text{CH}_2$) have previously been prepared by nitric acid oxidation of the corresponding manganese(I) complex, followed by precipitation with a suitable counterion (e.g. CdBr_4^{2-} , PF_6^- , NO_3^-) [12–15]. All of these complexes have the low-spin (t_{2g}^5) ground state and not the high-spin ($t_{2g}^3e_g^2$) configuration, which is more usual for manganese(II) complexes. Cyclic voltammetry of $[\text{Mn}(\text{CNR})_6]^{2+}$ ($z = 1, 2$) species result in complexes of manganese in the 0, +1, +2, and +3 oxidation states [15–18]. A number of other manganese(I) complexes are known, e.g. $\text{Mn}_2(\text{CO})_{10-n}(\text{CNR})_n$ ($n = 1–3$), $\text{Mn}_2(\text{CO})_7(\text{CNR})\{\text{P}(\text{O}^i\text{Pr})_3\}_2$ and $\text{Mn}_2(\text{CO})_n(\text{diphos})(\text{CNR})$ ($n = 4, 5$), formed by the reaction of RNC with $\text{Mn}_2(\text{CO})_{10}$ [19, 20], $\text{R}'\text{Mn}(\text{CO})_5$ [21], $\text{Mn}_2(\text{CO})_8\{\text{P}(\text{O}^i\text{Pr})_3\}_2$ [19] or $\text{Mn}_2(\text{CO})_5(\text{diphos})$ [22, 23]. In the case of manganese(I) complexes $\nu(\text{CN})$ of the isocyanide ligand decreases in energy upon coordination, indicating $d\pi-\pi_{\text{CN}}^*$ bonding in the complexes, whereas for manganese(II) the $\nu(\text{CN})$ increases on coordination [15, 24].

It can thus be seen that manganese(II) complexes of isocyanides are rare, but those which are known are of the $[\text{Mn}(\text{CNR})_6]^{2+}$ type. Here we wish to report some novel $[\text{MnX}_2(\text{CNBu}^t)]$ ($\text{X} = \text{Cl}, \text{Br}, \text{I}, \text{NCS}$) complexes and our observations of their reaction with PBu_3^n and molecular oxygen.

Results and Discussion

The method used to prepare the $\text{MnX}_2(\text{CNBu}^t)$ complexes is essentially the same as that used to prepare the analogous $\text{MnX}_2(\text{PR}_3)$ complexes (see 'Experimental'). It thus differs greatly from the methods used to prepare the $[\text{Mn}(\text{CNR})_6]^{2+}$ complexes inasmuch as the method reported here is a direct one-step process employing anhydrous condi-

*For Parts 11 and 12, see refs. 1 and 2, respectively.

**Author to whom correspondence should be addressed.

[†]Permanent address: Department of Chemistry, University of West Florida, Pensacola, Fla. 32514, U.S.A.

TABLE I. Some Physical and Analytical (%)^a Data for the Manganese(II) Complexes

Complex	Colour	Melting point (°C)	μ_{eff} (μ_{B})	C	H	N	X	Mn
MnCl ₂ (CNBu ^t)	White	>280	6.0	30.3(28.7)	4.3(4.9)		35.3(33.9)	26.5(26.3)
MnBr ₂ (CNBu ^t)	White	>280	5.6	21.2(20.1)	3.3(3.0)		52.7(53.7)	17.9(18.5)
MnI ₂ (CNBu ^t)	yellow	126(d)	5.1	15.0(15.3)	2.9(2.3)	3.5(3.6)	62.2(64.8)	15.0(14.0)
Mn(NCS) ₂ (CNBu ^t)	pale green	>280	6.0	32.8(33.1)	3.7(3.5)	16.4(16.5)		21.2(21.6)

^aFound(calc.).TABLE II. Some Significant Infrared Bands for the Manganese(II) Complexes (cm⁻¹)

Compound	$\nu(\text{CN})$	$\nu(\text{Mn}-\text{C})$	$\nu(\text{Mn}-\text{X}-\text{Mn})$	Other
MnCl ₂ (CNBu ^t)	2220(s)	213(sh)	234(sh)	
MnBr ₂ (CNBu ^t)	2105(s)	213(m)	188(br)	
MnI ₂ (CNBu ^t)	2179(s)	212(m)	120(br)	
Mn(NCS) ₂ (CNBu ^t)	2200(s)	214(sh)		2120(s) ^a 2098(s) ^a 264(s) ^b

^a $\nu(\text{CN})$ of NCS group. ^b $\nu(\text{Mn}-\text{N})$.

tions, whereas the latter require the pre-preparation of the manganese(I) complex followed by subsequent oxidation [12, 13]. Some physical properties and analytical data for the complexes are listed in Table I.

Some significant infrared spectral bands of the complexes as mulls are listed in Table II. A strong $\nu(\text{CN})$ band due to the isocyanide ligand is present in the 2220–2179 cm⁻¹ region, which may be compared to $\nu(\text{CN})$ in the free ligand at 2122 cm⁻¹. Isocyanide ligands are capable of acting as either good σ -donors or π -acceptors [15, 24] and, in general, CNBu^t behaves as a π -acceptor in zerovalent complexes, $\nu(\text{CN})$ decreases on coordination and as a σ -donor, $\nu(\text{CN})$ increases in energy on coordination, towards metals in higher oxidation states. It is clearly acting as a σ -donor ligand in these MnX₂(CNBu^t) complexes. The magnitude of the shift to higher energy, Cl(78 cm⁻¹) > Br(60 cm⁻¹) > I(57 cm⁻¹), is in accord with the electronegativity of the coordinated halogen, causing an overall shift in electron density away from the CNBu^t ligand towards the halide, X ← Mn ← CNBu^t.

For the MnX₂(CNBu^t) (X = Cl, Br, I) complexes only one $\nu(\text{Mn}-\text{X})$ band, assignable to bridging Mn–X–Mn moieties, Table II, is observable, and it is tempting, in the absence of terminal $\nu(\text{Mn}-\text{X})$ bands, to assign the complexes the structure found for the analogous MnI₂(PPhMe₂) (I) [25]. For Mn(NCS)₂(CNBu^t), however, there are two $\nu(\text{CN})$ bands assignable to thiocyanate at 2120 and 2097 cm⁻¹. Using accepted criteria [26], these can be assigned to bridging and terminal thiocyanate, respectively, suggesting a dimeric structure II. All of the complexes




exhibit an infrared band between 214–212 cm⁻¹, tentatively assigned to $\nu(\text{Mn}-\text{C})$.

The room temperature magnetic moments are listed in Table I. The values, 6.0–5.1 μ_{B} , are, for the most part, consistent with a degree of spin-pairing via Mn–X–Mn bridges and appear to mirror the magnetic behaviour of the analogous phosphine complexes. They also differ fundamentally from the low-spin hexakis(isocyanide)manganese(II) complexes [12]. The variable room temperature magnetic susceptibility measurements for MnBr₂(CNBu^t) show, Fig. 1, that this complex obeys the Curie–Weiss law with $\theta = 28$ K, suggesting antiferromagnetism, Fig. 1.

The room temperature X-band powder ESR spectra of the MnX₂(CNBu^t) show only single intense peaks at $g = 2$, providing no unambiguous structural information. However, the complexes dissolved in tetrahydrofuran (THF) and the X-band spectra at 93 K, Figs. 2–5, did yield some relevant structural data.




Fig. 1. Graph of reciprocal magnetic susceptibility vs. temperature for $\text{MnBr}_2(\text{CNBu}^t)$.




Fig. 2. X-band ESR spectra of (a) $\text{MnCl}_2(\text{CNBu}^t)$ and (b) $\text{MnCl}_2(\text{PPhBu}_2^t)$ in frozen THF, 93 K.




Fig. 3. X-band ESR spectra of (a) $\text{MnBr}(\text{CNBu}^t)$ and (b) $\text{MnBr}_2(\text{PPhBu}_2^t)$ in frozen THF, 93 K.




Fig. 4. X-band ESR spectra of (a) $\text{MnI}_2(\text{CNBu}^t)$ and (b) $\text{MnI}_2(\text{PPhBu}_2^t)$ in frozen THF, 93 K.

For the $\text{MnX}_2(\text{CNBu}^t)$ ($X = \text{Br}, \text{I}$), Figs. 3 and 4, peaks at $g = 6$ and $g = 2$ suggest axial symmetry and structure **III** would seem likely in tetrahydrofuran solution. Whilst the ESR spectra of the MnX_2 -





Fig. 5. X-band ESR spectra of (a) $\text{Mn}(\text{NCS})_2(\text{CNBu}^t)$ and (b) $\text{Mn}(\text{NCS})\{\text{P}(\text{C}_5\text{H}_{11})_3\}$ in frozen THF, 93 K.



(CNBu^t) ($X = \text{Cl}, \text{NCS}$) are more complex, Figs. 2 and 5, they are also indicative of axial symmetry. Further evidence for structure **III** is available from the solution infrared spectrum which exhibits only a single band at 2097 cm^{-1} assignable to terminal $\text{Mn}-\text{NCS}$ moieties.

Whilst the stoichiometry and magnetic and ESR properties of the $\text{MnX}_2(\text{CNBu}^t)$ complexes mirror those of the $\text{MnX}_2(\text{PR}_3)$ complexes no activity towards the reversible coordination of dioxygen has been observed in the former. Figures 2–5 contain a



comparison of the ESR spectra of the two types of complexes (for $X = \text{Cl}, \text{I}, \text{NCS}$) and it can be seen that there are strong similarities in THF solution. The exception is the bromide, for, whilst $\text{MnBr}_2(\text{CNBu}^t)$ does exhibit peaks at $g = 2$ and $g = 6$, Fig. 3, the $g = 6$ peak is of a different character from that of, for example, $\text{MnBr}_2(\text{PPhBu}^t)$.

Phosphine/Isocyanide Exchange Reactions

Although no activity towards dioxygen binding was observed for the $\text{MnX}_2(\text{CNBu}^t)$ complexes, tetrahydrofuran solutions of some $\text{MnX}_2(\text{CNBu}^t)$ complexes became active on addition of a stoichiometric (1:1) quantity of PBu_3^n , Table III. All measurements were made at 193 K using our standard gas burette apparatus [3].

The data given in Table III indicate that all the halide complexes, $\text{MnX}_2(\text{CNBu}^t)$ ($X = \text{Cl}, \text{Br}, \text{I}$), on addition of PBu_3^n (1:1), are active towards dioxygen coordination. The pseudohalide thiocyanate complex does not, however, form an active complex with PBu_3^n in tetrahydrofuran. This is consistent with data from observations of $\text{Mn}(\text{NCS})_2(\text{PBu}_3^n)$ in THF solution [27].

On exposure of THF solutions of a 1:1 mixture of $\text{MnX}_2(\text{CNBu}^t)$ ($X = \text{Cl}, \text{Br}, \text{I}$) and PBu_3^n at 0°C to an atmosphere of dry dioxygen a rapid and vivid colour change occurs. The visible spectra of the $\text{MnCl}_2(\text{CNBu}^t)/\text{PBu}_3^n/\text{O}_2$ and $\text{MnBr}_2/\text{PBu}_3^n/\text{O}_2$ systems exhibited absorption bands at 529 and 395 nm ($X = \text{Cl}$) and 574 and 418 nm ($X = \text{Br}$), the spectra being identical to those found for $\text{MnX}_2(\text{PBu}_3^n)(\text{O}_2)$ ($X = \text{Cl}, \text{Br}$) complexes in tetrahydrofuran [28]. The infrared spectra of these solutions exhibit bands at *ca.* 2120 cm^{-1} , assignable to free CNBu^t . The reactions may thus be expressed



The $\text{MnI}_2(\text{CNBu}^t)/\text{PBu}_3^n$ system also undergoes a vivid colour change in THF at 195 K when exposed to dioxygen. On warming to room temperature,

TABLE III. Dioxygen Uptake for the $\text{MnX}_2(\text{CNBu}^t)/\text{PBu}_3^n$ System^a

Complex	Concentration ($\times 10^{-3}\text{ mol dm}^{-3}$)	Total O_2 uptake (cm^3)	Less blank (cm^3)	Mn: O_2 ratio
$\text{MnCl}_2(\text{CNBu}^t)/\text{PBu}_3^n$	4.44	32.0	10.0	1.01
$\text{MnBr}_2(\text{CNBu}^t)/\text{PBu}_3^n$	6.19	35.2	13.2	0.95
$\text{MnI}_2(\text{CNBu}^t)/\text{PBu}_3^n$	9.14	42.6	20.6	1.01
$\text{Mn}(\text{NCS})_2(\text{CNBu}^t)/\text{PBu}_3^n$	6.23	22.4	0	0

^a100 cm^3 aliquots; blank THF absorbs 22.0 cm^3 dioxygen at 193 K.




Fig. 6. X-band ESR spectra of (a) $\text{MnCl}_2(\text{CNBu}^t) + \text{PBu}_3^n$ and (b) $\text{MnCl}_2(\text{PBu}_3^n)$ in frozen THF, 93 K.




Fig. 8. X-band ESR spectra of (a) $\text{MnI}_2(\text{CNBu}^t) + \text{PBu}_3^n$ and (b) $\text{MnI}_2(\text{PBu}_3^n)$ in frozen THF, 93 K.




Fig. 7. X-band ESR spectra of (a) $\text{MnBr}_2(\text{CNBu}^t) + \text{PBu}_3^n$ and (b) $\text{MnBr}_2(\text{PBu}_3^n)$ in frozen THF, 93 K.




Fig. 9. X-band ESR spectra of (a) $\text{Mn}(\text{NCS})_2(\text{CNBu}^t) + \text{PBu}_3^n$ and (b) $\text{Mn}(\text{NCS})_2\{\text{P}(\text{C}_5\text{H}_{11})_3\}$ in frozen THF, 93 K.

however, the green colour rapidly dispersed, consistent with observations on the $\text{MnI}_2(\text{PBu}_3^n)/\text{O}_2$ system [28].

Further evidence for a ligand-exchange process

taking place on addition of PBu_3^n to $\text{MnX}_2(\text{CNBu}^t)$ comes from ESR spectra which show subtle differences between the $\text{MnX}_2(\text{CNBu}^t)$ and the $\text{MnX}_2(\text{CNBu}^t)/\text{PR}_3$ systems, Figs. 6–9.

Experimental

Anhydrous manganese(II) salts were prepared as previously reported [29]. Physical measurements were performed as published earlier [30]. Gas uptakes were carried out as described in ref. 3.

Preparation of $MnX_2(CNBut)^t$ Complexes

The method was the same for each complex and will be exemplified for $Mn(NCS)_2(CNBut)^t$ here: anhydrous manganese(II) thiocyanate (1.02 g, 6.0 mmol) was added to freshly distilled dry toluene (70 cm³) in a 250 cm³ side-arm flask which had previously been flame dried *in vacuo*. Tertiary butylisocyanide (0.67 cm³) (Fluka Chemicals) was added and the mixture stirred for 72 h. The resulting yellow–white complex was filtered using a Schlenk apparatus and dried *in vacuo*.

Acknowledgements

We are grateful to The Science Research Council and The Central Electricity Generating Board for the award of a CASE Studentship (C.G.B.) and The Board of Regents of The University of West Florida for Sabbatical Leave (S.P.T.).

References

- 1 D. S. Barratt and C. A. McAuliffe, *J. Chem. Soc., Dalton Trans.*, submitted for publication.
- 2 D. S. Barratt and C. A. McAuliffe, *Inorg. Chim. Acta*, submitted for publication.
- 3 C. A. McAuliffe, H. F. Al-Khateeb, D. S. Barratt, J. C. Briggs, A. Challita, A. Hosseiny, M. G. Little, A. G. Mackie and K. Minten, *J. Chem. Soc., Dalton Trans.*, 2148 (1983).
- 4 M. G. Little, C. A. McAuliffe and J. B. Raynor, *J. Chem. Soc., Chem. Commun.*, 68 (1982).
- 5 C. A. McAuliffe, D. S. Barratt, C. G. Benson, A. Hosseiny, M. G. Little and K. Minten, *J. Organomet. Chem.*, 258, 23 (1983).
- 6 D. S. Barratt and C. A. McAuliffe, *J. Chem. Soc., Chem. Commun.*, 594 (1984).
- 7 G. A. Gott and C. A. McAuliffe, *J. Chem. Soc., Dalton Trans.*, accepted for publication (Paper 6/1561).
- 8 D. S. Barratt and C. A. McAuliffe, *Inorg. Chim. Acta*, 97, 37 (1985).
- 9 D. S. Barratt, C. G. Benson, G. A. Gott, C. A. McAuliffe and S. P. Tanner, *J. Chem. Soc., Dalton Trans.*, 2661 (1985).
- 10 G. A. Gott and C. A. McAuliffe, *J. Chem. Soc., Dalton Trans.*, (1987) in press (Paper 6/987).
- 11 G. A. Gott, J. Fawcett, C. A. McAuliffe and D. R. Russell, *J. Chem. Soc., Chem. Commun.*, 1283 (1984).
- 12 L. Malatesta and F. Bonati, 'Isocyanide Complexes of Metals', Wiley, New York, 1971.
- 13 L. Naldini, *Gazz. Chim. Ital.*, 90, 871 (1960).
- 14 D. S. Matteson and R. A. Bailey, *J. Am. Chem. Soc.*, 90, 3761 (1968).
- 15 D. T. Plummer and R. J. Angelici, *Inorg. Chem.*, 22, 4063 (1983).
- 16 P. M. Treichel and G. E. Dirreen, *J. Organomet. Chem.*, 39, C20 (1972).
- 17 P. M. Treichel, G. E. Dirreen and H. J. Much, *J. Organomet. Chem.*, 44, 339 (1972).
- 18 P. M. Treichel, D. W. Firsich and G. P. Essenmacher, *Inorg. Chem.*, 18, 2405 (1979).
- 19 S. Grant, J. Newman and A. R. Manning, *J. Organomet. Chem.*, 96, C11 (1975).
- 20 R. D. Adams and D. E. Chodosh, *J. Organomet. Chem.*, 87, C48 (1975).
- 21 K. K. Joshi, P. L. Pauson and W. H. Stubbs, *J. Organomet. Chem.*, 1, 51 (1963).
- 22 A. L. Balch and L. S. Benner, *J. Organomet. Chem.*, 135, 339 (1977).
- 23 L. S. Benner, M. M. Olmstead and A. L. Balch, *J. Organomet. Chem.*, 159, 289 (1978).
- 24 F. A. Cotton and F. Zingales, *J. Am. Chem. Soc.*, 83, 351 (1961).
- 25 T. J. King, B. Beagley, J. C. Briggs, A. Hosseiny, C. A. McAuliffe, K. L. Minten and W. E. Hill, *J. Chem. Soc., Chem. Commun.*, 1305 (1984).
- 26 A. H. Norbury, *Adv. Inorg. Chem. Radiochem.*, 17, 231 (1975).
- 27 A. Hosseiny, *Ph.D. Thesis*, University of Manchester, 1980.
- 28 H. F. Al-Khateeb, *Ph.D. Thesis*, University of Manchester, 1980.
- 29 A. Hosseiny, A. G. Mackie, C. A. McAuliffe and K. Minten, *Inorg. Chim. Acta*, 49, 99 (1981).
- 30 A. Hosseiny, C. A. McAuliffe, K. Minten, M. J. Parrot, R. Pritchard and J. Tames, *Inorg. Chim. Acta*, 39, 227 (1980).