

[Chem. Pharm. Bull.]
28(6)1971—1972(1980)

Syntheses of Fomecins A and B, Antibiotics produced by the Basidiomycete *Fomes juniperinus*

Fomecins A and B, phenolic antibiotics produced by the basidiomycete *Fomes juniperinus* SCHRENK, were synthesized starting from gallic acid.

Keywords—synthesis; fomecin A; fomecin B; antibiotic; phenol; *Fomes juniperinus*; basidiomycete

Fomecins A and B, isolated from the culture liquid of the basidiomycete *Fomes juniperinus* SCHRENK, have been characterized as 6-hydroxymethyl-2,3,4-trihydroxybenzaldehyde (**1**) and 3,4,5-trihydroxyphthalaldehyde (**2**), respectively.¹⁾ They show weak but rather broad antibacterial activities and antiviral activities.^{1a)} Chemical characteristics of such polyhydric phenols, combined with their biological activities, led us to undertake the syntheses of these two antibiotics.

3,4,5-Trimethoxybenzylalcohol (**3**), prepared from gallic acid (**4**) by a known procedure,²⁾ reacted with dichloromethyl methyl ether in CH_2Cl_2 at 0° (1 hr) with TiCl_4 as a catalyst to give 6-chloromethyl-2,3,4-trimethoxybenzaldehyde (**5**) (58%); mp 86° ; MS m/e : 246 ($M+2$), 244 (M^+), 209 ($M-35$); IR (Nujol) 1673 cm^{-1} (CHO); NMR (CDCl_3 , 60 MHz) δ : 10.45 (s, CHO), 6.95 (s, Ar), 5.05 (s, CH_2Cl), 4.02, 3.97, 3.90 (s, OCH_3). SeO_2 oxidation of **5** in dioxane (reflux, 2 hr) afforded 3,4,5-trimethoxyphthalaldehyde (**6**), mp $97-99^\circ$, in 56% yield, which was treated with BBr_3 ($-70-20^\circ$, 18 hr) in CH_2Cl_2 to give **2**. The conversion of **5** into **2** was also accomplished as follows. Reaction of **5** with AgNO_3 in MeOH gave methyl ether (**7**), mp 77° , (92%). Treatment of **7** with BBr_3 in CH_2Cl_2 provided 6-bromomethyl-2,3,4-trihydroxybenzaldehyde (**8**), mp ca. 150° (dec.), in 85% yield: MS m/e : 248 ($M+2$), 246 (M^+), 167 ($M-79$), 166 ($M-80$); IR (KBr) 1630 cm^{-1} (CHO); NMR ($\text{DMSO}-d_6$) δ : 10.28 (s, CHO), 6.60 (s, Ar), 4.98 (s, CH_2Br). Oxidation of **8** with SeO_2 in dioxane gave **2** in 27% yield; mp ca. 230° (dec.); MS m/e : 182 (M^+), 154 ($M-28$), 153 ($M-29$); UV (EtOH) 264, 287, 343 nm; IR (KBr) $1681, 1627\text{ cm}^{-1}$ (CHO); NMR (acetone- d_6) δ : 12.65 (s, OH), 10.82 (s, CHO), 10.07 (s, CHO), 9.27, 8.87 (s, OH), 7.20 (s, Ar). Melting points and the UV, IR, and NMR spectral data of **2** were consistent with those of fomecin B reported by McMorris and Anchel.^{1b,3)}




Chart 1

Several attempts to hydrolyze **8** were unsuccessful. However, catalytic hydrogenation of **2** in ethanol under 3 atm hydrogen in the presence of platinum oxide afforded **1** (8%) which was separated from the starting material by countercurrent distribution; mp 160° (dec.);

- 1) a) M. Anchel, A. Hervey, and W.J. Robbins, *Proc. Nat. Acad. Sci.*, **38**, 655 (1952); b) T.C. McMorris and M. Anchel, *Can. J. Chem.*, **42**, 1595 (1964).
- 2) S. Goodwin and B. Witkop, *J. Am. Chem. Soc.*, **79**, 179 (1957).
- 3) A total synthesis of fomecin B has just been reported: S.M. Al-Mousawi, R.J.S. Green, and J.F.W. McOmie, *Bull. Soc. Chim. Belg.*, **88**, 883 (1979).

MS m/e : 184 (M^+), 167 ($M-17$), 166 ($M-18$); UV (EtOH) 242, 305 nm; IR (KBr) 1637 cm^{-1} (CHO); NMR ($\text{DMSO}-d_6$) δ : 10.11 (s, CHO), 6.50 (s, Ar), 4.65 (s, CH_2O). **1** was identified by mp, UV, IR, NMR, and MS with an authentic sample of fomecin A. On acetylation, **1** gave the tetraacetate (**9**), mp 134° , which was identical to that derived from natural fomecin A.

Acknowledgement We are grateful to Dr. M.S.R. Nair, The New York Botanical Garden, for the authentic sample of fomecin A.

Shionogi Research Laboratories
Shionogi and Co., Ltd.
Fukushima-ku, Osaka, 553 Japan

KUNIO HAYASHI
KATSUYA TOKURA
KEI OKABE

Received April 25, 1980