


GENERATION AND REACTIONS OF THE ALLYLIC CARBANION SPECIES  
FROM 2-METHYL-3-(PHENYLTHIO)PROPENOIC ACID DERIVATIVES


Masaki KITAOKA, Yasuhiro TAKAHASHI, Hiroshi KOSUGI, and Hisashi UDA\*  
Chemical Research Institute of Non-Aqueous Solutions,  
Tohoku University, Katahira-2, Sendai 980

(Z)-2-Methyl-3-(phenylthio)propenoic acid and (E)-N-t-butyl-2-methyl-3-(phenylthio)propenamide produce selectively the corresponding allylic carbanion species, which react with aldehydes and ketones. The same carbanion species are also generated from 2-(phenylthiomethyl)propenoic acid and its N-t-butyl amide.

In the previous papers, we reported that (E)-2-methyl-3-(phenylthio)propenoic acid (1)<sup>1)</sup> and the methyl ester (2)<sup>2)</sup> produced selectively, upon treatment with lithium diisopropylamide (LDA) at -80 °C, the vinyl carbanion species (3) and (4), which reacted with carbonyl compounds to afford synthetically useful functionalized butenolides and cyclopentenones.<sup>1-3)</sup> If the allylic carbanion species could be generated selectively from the appropriate derivatives of 1, 1 would be used more widely as a versatile intermediate in construction of complex organic molecules. We report here the results of the lithiation of the isomeric (Z)-acid (5) and the (E)-amide derivative (6), in contrast to the cases of 1 and 2, leading selectively to the allylic carbanion species.



Upon treatment with 2.4 molequiv. of lithium 2,2,6,6-tetramethylpiperidide (LTMP) in tetrahydrofuran (THF) at -97 °C for 0.5 h the (Z)-acid (5)<sup>4,5)</sup> produced the allylic carbanion species (7), not the vinylic one, as indicated by a quenching experiment with deuterium oxide<sup>6)</sup> and by the reaction with aldehydes and ketones as described below.<sup>7)</sup> The allylic carbanion species (7) thus generated was treated with a slight excess of aldehydes or ketones at -97~-80 °C for 1~2 h to give the hydroxy carboxylic acid (8). In all of the reactions the separation of 8 was so difficult that the total crude product was subjected to lactonization by treating with *p*-toluenesulfonic acid in refluxing benzene to afford the α-phenylthio-methylene lactone (9) as the sole isolable product. The results are summarized in Table 1. The (E)-configuration of the phenylthiomethylene group in each 9 was established on the basis of the following evidence: appearance of the olefinic

Table 1. Reaction of the allylic carbanion species (**7**) with aldehydes and ketones

Starting acid	Aldehyde or ketone	Product ( <b>9</b> ) R <sup>1</sup> R <sup>2</sup>	Yield, % <sup>a</sup>
<b>5</b>	CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> CHO	H CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub>	43
"	(CH <sub>3</sub> ) <sub>2</sub> CHCHO	H (CH <sub>3</sub> ) <sub>2</sub> CH	46
"	CH <sub>3</sub> CO(CH <sub>2</sub> ) <sub>4</sub> CH <sub>3</sub>	CH <sub>3</sub> CH <sub>3</sub> (CH <sub>2</sub> ) <sub>4</sub>	34
"	cyclohexanone	-(CH <sub>2</sub> ) <sub>5</sub> -	49
<b>10</b>	(CH <sub>3</sub> ) <sub>2</sub> CHCHO	H (CH <sub>3</sub> ) <sub>2</sub> CH	40
"	cyclohexanone	-(CH <sub>2</sub> ) <sub>5</sub> -	40

<sup>a</sup> Yields are for the isolated pure products. The remaining in each reaction was mostly polymeric acid material.

proton signal at  $\delta$  7.20–7.60 overlapped with the aromatic proton signals. Such a down field chemical shift is acceptable for the (E)-geometry in referring the values of the (E)-acid (**1**) and -ester (**2**),  $\delta$  7.70 and 7.51, respectively.<sup>8)</sup> Furthermore, it was found that, in attempts to separate compound (**8**) as the methyl ester, sometimes the starting material was recovered as the (E)-ester (**2**), not the (Z)-ester. Thus, these results clearly indicate that the (Z)-configuration in the starting acid (**5**) was no longer retained in the allylic carbanion species and that the adduct arising from the reaction at the  $\beta'$  site in **7** and having sterically more favorable (E)-configuration was the only isolable product.

From the result on the deuteration experiment of **7**<sup>6)</sup> the isomeric acrylic acid derivative (**10**)<sup>9)</sup> would be expected to generate the same dianion species (**7**). Actually, under the identical conditions compound (**10**) produced **7** and reacted with isobutyraldehyde or cyclohexanone to yield, after lactonization, the same product (**9**) (Table 1).

For generation of an allylic carbanion species and for synthetic utility, we found that the (E)-amide derivative (**6**) as well as the isomer (**11**) was much more superior to the (Z)-acid (**5**). Thus, treatment of **6** or **11**<sup>10)</sup> with 2.2 mol equiv. of LDA in THF at -80~70 °C for 1.5 h produced selectively the relatively stable allylic carbanion species (**12**).<sup>11)</sup> The reaction of the dianion (**12**) with aldehydes

and ketones proceeded cleanly at  $-45 \sim -25$  °C for 1.5 h to give the single adduct ( $13$ ) in high yield.<sup>12)</sup> The results are summarized in Table 2. It was found that, although an example, the reaction with an  $\alpha,\beta$ -unsaturated ketone, cyclohexenone, gave only the 1,2-addition product (Table 2) and that the double bond in all  $13$  has (*E*)-configuration. Compound ( $13$ ) could be transformed into the  $\alpha$ -phenylthio-methylene lactone ( $9$ ) in good yield by a simple operation, heating in refluxing xylene.

The dianion species ( $12$ ), in contrast to the dianion ( $7$ ), also reacted with other electrophiles. Thus, addition of methyl acrylate ( $-55 \sim -30$  °C, 1 h), allyl bromide ( $-40 \sim -25$  °C, 40 min), or propylene oxide ( $-40$  °C/room temp., 1 day) to ( $12$ ) led to the corresponding adducts ( $14$ ), ( $15$ ), and ( $16$ ) in 48, 59, and 42% yields, respectively.

From the above results, the selective generation of the allylic carbanion species ( $7$ ) and ( $12$ ) from  $5$  and  $6$  would be accounted for by decrease of the acidity




Table 2. Reaction of the allylic dianion species ( $12$ ) with aldehydes and ketones


Aldehyde or ketone	Product ( $13$ ) R <sup>1</sup> R <sup>2</sup>	Yield, % <sup>a</sup>
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>2</sub> CHO	H CH <sub>3</sub> (CH <sub>2</sub> ) <sub>2</sub>	97
(CH <sub>3</sub> ) <sub>2</sub> CHCHO	H (CH <sub>3</sub> ) <sub>2</sub> CH	95
p-CH <sub>3</sub> OC <sub>6</sub> H <sub>4</sub> CHO	H p-CH <sub>3</sub> OC <sub>6</sub> H <sub>4</sub>	86
CH <sub>3</sub> COCH <sub>3</sub>	CH <sub>3</sub> CH <sub>3</sub>	80
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>4</sub> COCH <sub>3</sub>	CH <sub>3</sub> CH <sub>3</sub> (CH <sub>2</sub> ) <sub>4</sub>	87
cyclohexanone	-(CH <sub>2</sub> ) <sub>5</sub> -	73
cyclohex-2-en-1-one	-(CH <sub>2</sub> ) <sub>3</sub> CH=CH-	83

<sup>a</sup> Yields are for the isolated pure products, on the basis of recovered **6** (ca. 10%).


of the olefinic hydrogen atom due to the (Z)-geometry in **5** and the amide functionality<sup>13)</sup> in **6**.

We thank the Ministry of Education, Science, and Culture of Japan for financial support (a Grant-in-Aid for Scientific Research, No. 56540318).

#### References

- 1) Y. Takahashi, H. Hagiwara, H. Uda, and H. Kosugi, *Heterocycles*, **15**, 225 (1981).
  - 2) K. Isobe, M. Fuse, H. Kosugi, H. Hagiwara, and H. Uda, *Chem. Lett.*, **1979**, 785.
  - 3) Y. Takahashi, K. Isobe, H. Hagiwara, H. Kosugi, and H. Uda, *J. Chem. Soc., Chem. Commun.*, **1981**, 714; Y. Takahashi, H. Kosugi, and H. Uda, *ibid.*, **1982**, 496; *Chem. Lett.*, **1982**, 815.
  - 4) The pure (Z)-acid (**5**), mp 88~89 °C (from ether-petroleum ether) was prepared from the (E)-ester (**2**) by sequential reactions [photoisomerization, lithiation of the resulting isomeric mixture and reaction with *n*-butyraldehyde (twice), saponification of the (Z)-ester recovered unchanged (cf. Ref. 7)].
  - 5) All new compounds were characterized by combustion analysis as well as by IR and <sup>1</sup>H NMR spectroscopy.
  - 6) The <sup>1</sup>H NMR spectrum of the recovered material revealed clearly the signals due to the deuterated species (**1**) and (**1**). However, the yield and the proportion of **1** and **1** could not be determined.
- 

**1**



**1**
- 7) Lithiation with LDA at -90~-70 °C could not be detected. Also, no lithiation was observed upon treatment of the (Z)-ester with either LDA or LTMP at -90~-70 °C. After quenching with deuterium oxide, the starting materials were completely recovered undeuterated.
  - 8) The (Z)-isomer of **9** ( $R^1 = H$ ,  $R^2 = n-C_4H_9$ ), prepared by photoisomerization, revealed the olefinic proton signal at  $\delta$  6.88, being in good agreement with those of the (Z)-acid (**5**) and the ester,  $\delta$  7.10 and 6.83, respectively.
  - 9) G. K. Pajagopalan and S. Swaminathan, *Synthesis*, **1976**, 409.
  - 10) The amides (**6**) and (**1**) were easily prepared by the reaction of the corresponding acid chloride with *t*-butylamine.
  - 11) Deuteration of **12** resulted in the 90% recovery of the ca. 80%  $\beta'$ -deuterated **6**.
  - 12) We also found that the *N,N*-diisopropyl amide derivative produced the allylic monoanion species. In this case, however, the monoanion species was considerably unstable, decomposed above -50 °C and the reaction with acetone gave two adducts (29 and 44%) arising from the addition of acetone on both  $\beta$  and  $\beta'$  reactive sites.
  - 13) K. Tanaka, Y. Nozaki, N. Tamura, R. Tanikaga, and A. Kaji, *Chem. Lett.*, **1980**, 1567; D. J. Kempf, K. D. Wilson, and P. Beak, *J. Org. Chem.*, **47**, 1610 (1982).

(Received May 2, 1983)