

## $\beta$ Gal(1 $\rightarrow$ S-4) $\beta$ GlcNAc-OR: A GALACTOSIDASE-STABLE SUBSTRATE FOR $\alpha$ (1 $\rightarrow$ 3)FUCOSYLTRANSFERASE

Yili Ding, Ole Hindsgaul,\* Hong Li and Monica M. Palcic

Department of Chemistry, University of Alberta, Edmonton, Alberta, T6G 2G2, Canada


Received 7 August 1998; accepted 1 October 1998

**Abstract:** 1'-4 Thio-*N*-acetylglucosamine was chemically synthesized as a galactosidase-stable substrate for  $\alpha$ (1 $\rightarrow$ 3)fucosyltransferase. The product of enzymatic fucose addition was confirmed to be the thio-Le X analog.


© 1998 Elsevier Science Ltd. All rights reserved.

*N*-Acetylglucosamine is one of the most frequently occurring constituents of glycosphingolipids and glycoproteins in which the carbohydrate moieties often represent epitopes for cellular recognition, signal transduction, and adhesion phenomena.<sup>1</sup> For instance, *N*-Acetylglucosamine (LacNAc) can serve as a recognition site in bacterial adhesion<sup>2</sup> and as a tumor-associated antigen.<sup>3</sup> LacNAc also serves as a substrate for fucosyltransferases, sialyltransferases, and *N*-acetylglucosaminyltransferases involved in the biosynthesis of complex oligosaccharides.

We have previously reported<sup>4</sup> that fluorescently labeled LacNAc derivatives are useful for assaying the biosynthetic activities of glycosyltransferases in crude cell extracts where product formation can be quantitated by capillary electrophoresis with laser-induced fluorescence detection. One of the difficulties in such experiments, however, was that products of degradation initiated by  $\beta$ -galactosidase seriously depleted the LacNAc substrate. We therefore considered two possible analogs (**2** and **3**) of LacNAc (**1**) that might overcome this limitation. Disaccharide **2**, where the ring oxygen of the  $\beta$ -Gal residue was replaced with a sulfur, was available from previous work<sup>5</sup> where it was found to be resistant to  $\beta$ -galactosidase. However, enzymatic assays revealed **2** to be neither an acceptor nor an inhibitor for an  $\alpha$ (1 $\rightarrow$ 3/4) fucosyltransferase partially purified from human milk. We therefore turned to the synthesis and enzymatic evaluation of the thio-disaccharide **3** where the oxygen in the glycosidic linkage is replaced by sulfur. Such 1-thio-glycosides are known to be resistant to glycosidases.<sup>6</sup>



3,4,6-Tri-*O*-acetyl-2-phthalimido- $\alpha$ -D-galactopyranosyl bromide (**4**) was coupled with 8-methoxycarbonyloctanol in dichloromethane in the presence of AgOTf, affording the  $\beta$ -linked glycoside **5** in 71% yield. *O*-Deacetylation of **5** with NaOMe/MeOH, followed by treatment with hydrazine acetate in ethanol and subsequent *N*-acetylation, yielded 8-methoxycarbonyl-2-acetamido-2-deoxy- $\beta$ -D-galactopyranoside (**6**) in 58% yield (three steps). Selective benzylation of **6** with benzoyl chloride (2.2 equiv at  $-40^\circ\text{C}$ ), gave 3,6-dibenzoate derivative **7** in 63% yield. Reaction with triflic anhydride in pyridine converted **7** into the 4-*O* triflate **8**, which was then coupled with 2,3,4,6-tetra-*O*-acetyl-1-thio- $\beta$ -D-galactopyranose (**9**) in DMF containing NaH at  $-20^\circ\text{C}$ . The resulting sulfur-linked disaccharide was deacetylated with NaOMe/MeOH, providing 1'-4 thio-*N*-acetylactosamine (**3**) in 28% isolated yield (three steps). The  $^1\text{H}$  NMR signals at 4.43 ppm (d, 1H,  $J = 9.6$  Hz), 4.39 ppm (d, 1H,  $J = 8.4$  Hz) and  $^{13}\text{C}$  NMR signals at 102.35 ppm, 86.26 ppm confirmed the structure.



**Scheme 1** *a*: 1,  $\text{HO}(\text{CH}_2)_8\text{CO}_2\text{Me}$ , AgOTf, 4 Å MS, 71%. *b*: 1, NaOMe/MeOH; 2,  $\text{NH}_2\text{NH}_2\cdot\text{HOAc}$ , EtOH,  $70^\circ\text{C}$ ; 3, MeOH, Ac<sub>2</sub>O, 58% (1-3, three steps). *c*: Benzoyl chloride (2.2 equiv),  $-40^\circ\text{C}$ , 63%. *d*:  $\text{Tf}_2\text{O}$ , pyridine,  $-10^\circ\text{C}$ . *e*: 1, **9** (1.5 equiv), DMF, NaH; 2, NaOMe/MeOH; 38% (three steps).

The sulfur-linked disaccharide **3** was evaluated as a potential acceptor for a partially-purified human milk  $\alpha(1\rightarrow3/4)$  fucosyltransferase that has been extensively characterized and used in synthesis.<sup>7</sup> By using an

established radioactive “Sep-Pak assay”,<sup>8</sup> disaccharide **3** was found to be a kinetically competent acceptor displaying a  $K_m$  value of 230  $\mu\text{M}$  and  $V_{\text{max}}$  (20 pmol/min for the enzyme preparation used) was nearly the same (~90%) as that for the parent LacNAc disaccharide **1**. Under the same conditions, **2** was inactive.

To confirm that the expected product was formed, a preparative incubation was performed using 1.2 mg of **3**.<sup>9</sup> The produced trisaccharide **10** showed  $^1\text{H}$  NMR signals at 5.03 ppm (d, 1H,  $J = 3.8$  Hz), 4.47 ppm (d, 1H,  $J = 8.4$  Hz), 4.46 ppm (d, 1H,  $J = 8.4$  Hz) and 4.62 ppm (q, 1H) confirming the 1'-4-thio-Le X structure.<sup>10</sup>

Finally, the resistance of **3** to  $\beta$ -galactosidase was experimentally verified. No hydrolysis of thiodisaccharide **3** was evident after incubation with 590 mU of  $\beta$ -galactosidase from E.Coli for 30 min under conditions previously reported.<sup>5</sup> Galactose release was monitored using a kit from Boehringer-Mannheim containing 88 mU of galactose dehydrogenase and NAD.<sup>5</sup> Under these conditions, the parent LacNAc **1** (510  $\mu\text{M}$ ) was completely hydrolyzed.

**Acknowledgment.** This work was supported by the Natural Sciences and Engineering Research Council of Canada.

## References and notes

1. Schmidt, R. R. *Adv. Carbohydr. Chem. Biochem.* **1994**, 50, 21.
2. Karlsson, K. A. *Adv. Rev. Biochem.* **1989**, 59, 309.
3. Hakomori, S. *Adv. Cancer Res.* **1989**, 52, 257.
4. Zhang, Y.; Le, X.; Dovichi, N. J.; Compston, C. A.; Palcic, M. M.; Diedrich, P.; Hindsgaul, O. *Anal. Biochem.* **1994**, 227, 368.
5. Yuasa, H.; Hindsgaul, O.; Palcic, M. M. *J. Amer. Chem. Soc.* **1992**, 114, 5891.
6. Defaye, J.; Gelas, J. *In Study in Natural Products Chemistry*, Atta-ur-Rahman Ed. Elsevier: Amsterdam, **1991**, 8, 315.
7. Qian, X.; Hindsgaul, O.; Li, H.; Palcic, M. M. *J. Amer. Chem. Soc.* **1998**, 120, 2184 and references cited therein.
8. Palcic, M. M.; Heerze, L. D.; Pierce, M.; Hindsgaul, O. *Glycoconjugate J.* **1988**, 5, 49.
9. The reaction mixture contained **3** (1.2 mg), GDP-Fuc (1.7 mg), 30 mU of milk fucosyltransferase in 400  $\mu\text{L}$  of 25 mM sodium cacodylate buffer, pH 6.5, containing 25 mM  $\text{MnCl}_2$ , 0.2% BSA and 25% glycerol

was incubated at 37 °C for 3 days. Additional GDP-Fuc (1.0 mg) was added after 24 h and again after 48 h. The reaction was stopped by the addition of 8 mL of water and the sample was isolated by loading the mixture onto two sequential C-18 Sep-Pak cartridges (Waters). The cartridges were washed with water to remove enzyme and unreacted nucleotide donor and the product was eluted with methanol to give trisaccharide **10**.

10. Spectral data for selected new compounds: **3**.  $^1\text{H}$  NMR (360 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  4.43 (d, 1H,  $J = 9.6$  Hz, H'-1), 4.39 (d, 1H,  $J = 8.0$  Hz, H-1), 3.64 (s, 3H,  $\text{OCH}_3$ ), 2.83 (m, 1H, H-2), 1.96 (s, 3H,  $\text{NHCOCH}_3$ );  $^{13}\text{C}$  NMR (300 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  102.35 (C-1), 86.26 (C-1'), 81.01, 78.20, 75.95, 73.59, 70.67, 70.48 (C-3, C-5, C-2', C-3', C-4' and C-5'), 70.07 ( $\text{OCH}_2$ ), 63.31, 62.84 (C-6 and C-6'), 58.57 (C-2), 51.95 ( $\text{OCH}_3$ ), 23.02 ( $\text{NHCOCH}_3$ ); HRMS for  $\text{C}_{24}\text{H}_{43}\text{NO}_{12}\text{SNa}$   $[\text{M}+\text{Na}]^+$  592.2404, found: 592.2408. **6**.  $^1\text{H}$  NMR (360 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  4.35 (d, 1H,  $J = 8.4$  Hz, H-1), 3.64 (s, 3H,  $\text{OCH}_3$ ), 1.95 (s, 3H,  $\text{NHCOCH}_3$ ). **7**.  $^1\text{H}$  NMR (360 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  5.86 (d, 1H,  $J = 8.9$  Hz, NH), 5.44 (dd, 1H, H-3), 4.76 (d, 1H,  $J = 8.3$  Hz, H-1), 3.66 (s, 3H,  $\text{OCH}_3$ ), 3.52 (m, 1H, H-2), 1.89 (s, 3H,  $\text{NHCOCH}_3$ );  $^{13}\text{C}$  NMR (300 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  101.05 (C-1), 74.24, 72.16, 66.74 (C-3, C-4 and C-5), 69.48 ( $\text{OCH}_2$ ), 63.35 (C-6), 50.58 (C-2), 51.26 ( $\text{OCH}_3$ ), 22.85 ( $\text{NHCOCH}_3$ ). **8**.  $^1\text{H}$  NMR (360 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  6.05 (dd, 1H, H-3), 5.48 (d, 1H,  $J = 2.9$  Hz, H-4), 5.17 (d, 1H,  $J = 8.3$  Hz, H-1), 3.65 (s, 3H,  $\text{OCH}_3$ ), 1.88 (s, 3H,  $\text{NHCOCH}_3$ ). **10**.  $^1\text{H}$  NMR (360 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  5.03 (d, 1H,  $J = 3.8$  Hz, H-1'), 4.47 (d, 1H,  $J = 8.4$  Hz, H-1), 4.46 (d, 1H,  $J = 8.4$  Hz, H-1'), 4.62 (q, 1H, H-5''), 3.64 (s, 3H,  $\text{OCH}_3$ ), 2.30 (t, 2H,  $\text{CH}_2\text{CO}$ ), 1.94 (s, 3H,  $\text{NHCOCH}_3$ ); FABMS:  $m/z$  738  $[\text{M}+\text{Na}]^+$ .