

## Studies on Covalently Linked Porphyrin–C<sub>60</sub> Dyads: Stabilization of Charge-Separated States by Axial Coordination

Francis D'Souza,<sup>\*,†</sup> Suresh Gadde,<sup>†</sup> Melvin E. Zandler,<sup>†</sup> Klykov Arkady,<sup>†</sup>  
Mohamed E. El-Khouly,<sup>‡</sup> Mamoru Fujitsuka,<sup>‡</sup> and Osamu Ito<sup>\*,‡</sup>

Department of Chemistry, Wichita State University, 1845 Fairmount, Wichita, Kansas 67260-0051, and  
Institute of Multidisciplinary Research for Advanced Materials, Tohoku University, Katahira,  
Sendai, 980-8577, Japan

Received: August 21, 2002

The effect of axial ligation on the photoinduced charge separation and charge recombination of a series of covalently linked porphyrin–C<sub>60</sub> dyads is investigated. Toward this, *meso*-tetraphenylporphyrin and its zinc(II) derivative are functionalized at the *ortho* or *para* positions of one of the aryl groups to bear a fulleropyrrolidine entity through a flexible ethylene dioxide bridge to probe the effect of intramolecular electron transfer phenomena. In *o*-dichlorobenzene, 0.1 M (TBA)ClO<sub>4</sub>, the synthesized dyads exhibit seven one electron reversible redox reactions within in the potential window of the solvent and the measured redox potentials and UV–visible absorption spectra reveal charge–transfer interactions between the electron donor, porphyrin, and the electron acceptor, fullerene entities. The geometric and electronic structures of the dyads probed by *ab initio* B3LYP/3-21G(\*) methods also revealed the existence of charge–transfer interactions. The excited state photochemical events are monitored by both steady state and time-resolved emission as well as transient absorption techniques. In *o*-dichlorobenzene or in benzonitrile, the main quenching pathway involves charge separation from the excited porphyrin to the moiety. The *k*<sub>cs</sub> and *k*<sub>cr</sub> are found to depend on the type of substitution (*ortho* or *para*) and the metal ion in the porphyrin cavity. Relatively long-lived charge-separated states are observed upon coordinating pyridine axial ligands to the central metal ion of the zinc porphyrin–C<sub>60</sub> dyads, and this has been attributed to the electronic and geometric effects caused by the axial coordinated ligand.

### Introduction

Studies on light-induced electron transfer in covalently linked donor–acceptor systems have witnessed enormous growth in recent years mainly to address the mechanistic details of electron transfer in chemistry and biology, to develop artificial photosynthetic systems for light energy harvesting, and also to develop molecular electronic devices.<sup>1,2</sup> Toward constructing such dyads, fullerenes<sup>3</sup> and porphyrins<sup>1</sup> have been utilized as important constituents owing to their rich redox,<sup>4</sup> optical,<sup>5</sup> and photochemical<sup>6</sup> properties. In contrast to the traditionally used two-dimensional aromatic electron acceptors, the fullerene in donor–acceptor dyads accelerates photoinduced charge separation and slows down charge recombination. Imahori and Sakata<sup>7</sup> attributed the slower charge recombination process to the small reorganization energy of C<sub>60</sub>. The small reorganization energy of C<sub>60</sub> predicts that *k*<sub>cs</sub> moves upward along the normal region into the top region of the Marcus curve, and *k*<sub>cr</sub> lies to be significantly downward into the inverted region.

More recently, elegantly designed porphyrin–C<sub>60</sub> dyads have been synthesized to probe the effect of molecular topology and distance and orientation effects of the donor and acceptor entities on photoinduced charge separation and charge recombination processes.<sup>8–11</sup> Supramolecular triads, tetrads, pentads, etc. have also been synthesized and studied. In these supramolecular systems, distinctly separated donor–acceptor radical ion pairs, in succession, are generated upon initial charge separation by

charge migration reactions along the well-tuned redox gradients. Photosynthetic reaction centers composed of self-assembled donor and acceptor entities produce long-lived, highly energetic, charge-separated states with quantum yields close to unity by using this mechanism of charge migration.<sup>12</sup>


In the present study, we report the effect of axial ligation on the photoinduced charge separation and charge recombination of covalently linked porphyrin–C<sub>60</sub> dyads. Toward this, first, a new synthetic methodology has been developed to covalently link porphyrin and fullerene entities with flexible bonds, which allows a direct special interaction between the porphyrin and the fullerene entities. Here, *meso*-tetraphenylporphyrin is functionalized at the *ortho* or *para* positions of one of the aryl groups to bear a fulleropyrrolidine entity through an ethylene dioxide bridge (Scheme 1). The *ortho* and *para* positions of the phenyl entity are primarily chosen to probe the orientation effects. The free-base porphyrin and zinc porphyrin are selected to visualize the effect of the free energy changes on the kinetics of photoinduced charge separation and charge recombination. Next, a series of substituted pyridine ligands are utilized to form penta-coordinated zinc porphyrin–C<sub>60</sub> dyads (Scheme 1). Here, the axial pyridine ligands enhance the electron donor ability of zinc(II) tetraphenylporphyrin and are expected to slow the charge recombination processes upon initial charge separation of the zinc porphyrin–C<sub>60</sub> dyad by the electronic and geometric effects caused by the axial pyridine ligand. Time-resolved emission and transient absorption spectral techniques have been employed to probe these reactions.

<sup>†</sup> Wichita State University.

<sup>‡</sup> Tohoku University.



## SCHEME 2




Similarly, for the ZnPp-C<sub>60</sub> derivative,  $R_{D-A}$  and  $R_{ee}$  are found to be, respectively, 6.4 and 3.9 Å, suggesting close proximity of the two entities irrespective of the nature of the substitution (*ortho* or *para*). For the H<sub>2</sub>Po-C<sub>60</sub> dyad, the  $R_{D-A}$  and  $R_{ee}$  are found to be slightly longer than those observed for the corresponding ZnPo-C<sub>60</sub> and are about 7.08 and 3.67 Å, respectively. As shown in Figure 2, little or no distortion of the porphyrin ring is observed for the *para*-substituted derivatives, while substantial distortion of the porphyrin ring atoms near the substituted *meso*-phenyl ring is observed in the case of the *ortho*-substituted derivative.

The existence of intramolecular interactions between the porphyrin ring and the C<sub>60</sub> moiety is evidenced by the frontier highest occupied molecular orbital (HOMO) and lowest unoccupied molecular orbital (LUMO) molecular orbitals. Figure 3a shows the HOMO and LUMO for the ZnPp-C<sub>60</sub> dyad. The majority of the HOMO is located on the porphyrin ring while small amounts of the HOMO are also found on the C<sub>60</sub> carbons located within the interacting distances, suggesting through space interactions. Similarly, the majority of the LUMO is located on the C<sub>60</sub> spheroid, with small amounts located on the porphyrin

ring atoms. The delocalization of HOMO and LUMO over the zinc porphyrin and C<sub>60</sub> entities suggests through space charge-transfer interactions both in the ground state and in the excited state of the zinc porphyrin-C<sub>60</sub> dyads. Similar charge-transfer interactions for the ZnPo-C<sub>60</sub> are observed for only the HOMO but not for the LUMO. However, for the H<sub>2</sub>Po-C<sub>60</sub>, because of the larger separation between the porphyrin and the C<sub>60</sub> entities, such effects are not observed; that is, the HOMO is found entirely located on the porphyrin ring while the LUMO is on the C<sub>60</sub> spheroid (see Supporting Information). The gas phase HOMO-LUMO energy gap is found to be 1.66, 1.59, and 1.81 eV for the ZnPo-C<sub>60</sub>, ZnPp-C<sub>60</sub>, and H<sub>2</sub>Pp-C<sub>60</sub> dyads, respectively.

**Cyclic Voltammetry Studies.** Determination of the redox potentials of the newly formed molecular donor-acceptor type systems is important to probe the existence of charge-transfer interactions between the donor and the acceptor in the ground state and also to evaluate the energetics of electron-transfer reactions. With this in mind, we have performed a systematic study to evaluate the redox behavior of the dyads using the cyclic voltammetric technique. The cyclic voltammograms of



**Figure 1.** (a) Normalized absorption spectra in the Soret region of (TPP)Zn (dotted), ZnPo-C<sub>60</sub> (solid), and ZnPp-C<sub>60</sub> (dashed) in DCB. (b) Normalized absorption spectra in the Q-band region of equimolar (TPP)Zn and 2-propyl fulleropyrrolidine (solid), ZnPo-C<sub>60</sub> (dashed), and ZnPp-C<sub>60</sub> (dotted) in DCB.


ZnPo-C<sub>60</sub> and H<sub>2</sub>Po-C<sub>60</sub> as representative examples are shown in Figure 4, and the data for all of the derivatives are summarized in Table 2.

Both free-base porphyrin and zinc porphyrin dyads exhibited a total of seven reversible redox processes within the accessible potential window of *o*-dichlorobenzene (DCB) containing 0.1 M (TBA)ClO<sub>4</sub> (Figure 4). The first and second redox potentials corresponding to the oxidation of the zinc porphyrin in ZnPo-C<sub>60</sub> are located at  $E_{1/2} = 0.29$  and  $0.63$  V vs Fc/Fc<sup>+</sup>, respectively. The reduction potentials of the appended C<sub>60</sub> moiety are located at  $E_{1/2} = -1.17$ ,  $-1.55$ , and  $-2.07$  V vs Fc/Fc<sup>+</sup>, while the corresponding potentials for the zinc porphyrin ring reduction are located at  $E_{1/2} = -1.90$  and  $-2.29$  V vs Fc/Fc<sup>+</sup>, respectively. The first and second oxidation potentials of H<sub>2</sub>Po-C<sub>60</sub> shift to

**TABLE 1: Formation Constant for Axial Coordination of Pyridine Ligands to Zinc-Porphyrin C<sub>60</sub> Dyads in DCB at 298 K**


| compd                | axial ligand <sup>b</sup> | $K$ (M <sup>-1</sup> ) <sup>a</sup> |
|----------------------|---------------------------|-------------------------------------|
| ZnPo-C <sub>60</sub> | 4-acetyl pyridine         | $1.46 \times 10^4$                  |
|                      | pyridine                  | $1.71 \times 10^4$                  |
| ZnPp-C <sub>60</sub> | 4-dimethylamino pyridine  | $13.4 \times 10^4$                  |
|                      | 4-acetyl pyridine         | $1.18 \times 10^4$                  |
| (TPP)Zn              | pyridine                  | $1.56 \times 10^4$                  |
|                      | 4-dimethylamino pyridine  | $11.8 \times 10^4$                  |
|                      | 4-acetyl pyridine         | $1.06 \times 10^4$                  |
|                      | 4-dimethylamino pyridine  | $5.20 \times 10^4$                  |

<sup>a</sup> Error in  $K$  values =  $\pm 10\%$ . <sup>b</sup> Because of the limited solubility of 4-hydroxypyridine in DCB, the  $K$  values for this ligand binding to ZnP-C<sub>60</sub> could not be obtained. <sup>c</sup> From ref 11g.



**Figure 2.** Ab initio B3LYP/3-21G(\*)-calculated geometric structures of (a) ZnPo-C<sub>60</sub> and (b) ZnPp-C<sub>60</sub>.

more positive potential by ca. 0.2 V as compared with those of ZnPo-C<sub>60</sub>. The reduction potentials of the appended C<sub>60</sub> moiety of H<sub>2</sub>Po-C<sub>60</sub> are almost the same as those of ZnPo-C<sub>60</sub>, while the corresponding potentials for the free-base porphyrin ring reduction shift to less negative values. The overall electrochemical behavior of the *para*-substituted derivatives is found to be similar to that of the *ortho*-substituted derivatives and revealed slight changes in the redox potential values (Table 2). A comparison of the redox potential values between the differently substituted dyads and their starting monomeric species, porphyrins and fulleropyrrolidine, shows small changes, which are




**Figure 3.** Frontier HOMO and LUMO of (a) ZnPp-C<sub>60</sub> and (b) Py:ZnPp-C<sub>60</sub> calculated by ab initio B3LYP/3-21G(\*) methods.

in agreement with the existence of charge-transfer interactions as shown by the spectral and computational studies. The electrochemically measured HOMO-LUMO gap, that is, the difference between the first oxidation potential of porphyrin and fulleropyrrolidine (Table 2), follows a trend similar to that obtained by B3LYP/3-21G(\*) methods in the gas phase (vide supra). These results indicate a smaller HOMO-LUMO gap for the zinc porphyrin-C<sub>60</sub> dyads as compared to the free-base porphyrin-C<sub>60</sub> dyads.

It is of interest to probe the sequence of the site of electron transfer in the investigated dyads as shown for ZnP-C<sub>60</sub> in Scheme 3 by ab initio B3LYP/3-21G(\*) methods. In this regard, our recent studies on ferrocene-C<sub>60</sub>-nitroaromatic entities bearing molecular triads have demonstrated excellent tracking of the sequence of the site of electron transfer involving the three redox active entities by the computed HOMO and LUMO orbitals.<sup>17b,c</sup> Figure 5 illustrates the computed HOMO and LUMO orbitals for the ZnPp-C<sub>60</sub> triad. Similar results were obtained for the ZnPo-C<sub>60</sub> and the H<sub>2</sub>P-C<sub>60</sub> dyads. In agreement with the electrochemical results (Scheme 3), a majority of the HOMO and HOMO-1 orbitals are located on the zinc porphyrin entity while the majority of the LUMO and LUMO+1 are located on the C<sub>60</sub> entities. The subsequent LUMO orbitals follow the order C<sub>60</sub>, porphyrin-C<sub>60</sub>, C<sub>60</sub>, porphyrin-C<sub>60</sub>, and


this compares with the electrochemical sequence of redox reactions: porphyrin, C<sub>60</sub>, porphyrin, and C<sub>60</sub>.

The pyridines utilized to form the penta-coordinated species revealed an irreversible oxidation process located at  $E_{pa} = 0.25$  V vs Fc/Fc<sup>+</sup>, which is less positive than zinc porphyrin ( $E_{1/2} = 0.28$  V), suggesting an increased electron donor ability of zinc porphyrin when photoinduced charge separation takes place in ZnP-C<sub>60</sub>. During the titrations involving ZnP-C<sub>60</sub> and pyridines, the first and second oxidation peaks corresponding to the zinc porphyrin revealed a cathodic shift of about 15–20 mV (on addition of 3–4 equiv of pyridine) as a result of axial ligation. These results suggest that the zinc porphyrin in Py:ZnP-C<sub>60</sub> is slightly a better electron donor than that in ZnP-C<sub>60</sub>. The HOMO and LUMO orbitals generated for Py:ZnP-C<sub>60</sub> revealed delocalization of HOMO orbitals over the Py:ZnP entity (Figure 3b), similar to that reported earlier for zinc porphyrin-(2-(pyridyl))fulleropyrrolidine dyad.<sup>11g</sup> Additionally, the observed partial delocalization of HOMO on the C<sub>60</sub> entity and LUMO on the porphyrin entity of ZnPp-C<sub>60</sub> (Figure 3a) is not seen in Py:ZnPp-C<sub>60</sub> (Figure 3b). This may be due to the increased distance between the ZnP and the C<sub>60</sub> entities of the dyad upon axial ligation (~0.5 Å) and the pulling of the Zn from a 0.3 Å bias out of the porphyrin ring cavity toward the C<sub>60</sub> (in ZnPp-C<sub>60</sub>) to a 0.3 Å bias away from the



**Figure 4.** Cyclic voltammograms of (a) ZnPo-C<sub>60</sub> and (b) H<sub>2</sub>Po-C<sub>60</sub> in DCB, 0.1 M (TBA)ClO<sub>4</sub>. Scan rate = 100 mV/s.

### SCHEME 3



C<sub>60</sub> toward the pyridine as a result of Zn-N axial bond formation. The delocalization of the frontier HOMO over the axially bound pyridine ligand would dissipate the positive charge of ZnP<sup>+</sup> formed during initial charge separation and slow the charge recombination process. Additionally, the slightly increased distance between the ZnP and the C<sub>60</sub> entities of Py:ZnPp-C<sub>60</sub> as compared to that in ZnPp-C<sub>60</sub> might also contribute toward slowing down the charge recombination process.

**Fluorescence Emission Studies.** The photochemical behavior of the free-base porphyrin-C<sub>60</sub> and zinc porphyrin-C<sub>60</sub> dyads is investigated first by using steady state fluorescence measurements. Figure 6 shows fluorescence spectra of the zinc porphyrin-C<sub>60</sub> and free-base porphyrin-C<sub>60</sub> dyads along with ZnP and H<sub>2</sub>P used as a reference compounds. The peak maxima corresponding to porphyrin emission of all of the dyads revealed a red shift of 1–2 nm as compared to their unsubstituted derivatives. In the case of ZnPp-C<sub>60</sub>, the porphyrin emission bands are quenched by ca. 73% while this quenching for the ZnPo-C<sub>60</sub> is found to be over 95% as compared with the emission intensity of ZnP. Similar observations are made for the free-base porphyrin dyads. The corresponding quenching of emission of H<sub>2</sub>Pp-C<sub>60</sub> and H<sub>2</sub>Po-C<sub>60</sub> is found to be ca. 78 and 97%, respectively, as compared to H<sub>2</sub>P. These observations suggest efficient quenching of the singlet excited state of the porphyrins by the appended fullerene entity, and such quenching for the *ortho*-substituted derivatives is higher than the *para*-substituted ones. It may also be mentioned here that changing the solvent from DCB to a more polar benzonitrile (BN) increased the overall quenching efficiency by another 2–5%.

Interestingly, on extending the wavelength scan to a higher wavelength, a weak emission at ~715 nm was observed in the case of zinc porphyrin derivatives corresponding to the emission of fulleropyrrolidine entity<sup>17b</sup> (Figure 6). However, the 715 nm emission was usually hidden by the tail of the huge emission from zinc porphyrin itself. Hence, it was difficult to confirm the energy transfer from the singlet excited porphyrin to the fullerene entity in these dyads.


Addition of Py (up to 10 equiv) to a solution of ZnP-C<sub>60</sub> resulted in further red shift (~10 nm) and a decrease of the intensity of the emission bands (20–30%) suggesting the formation of a penta-coordinated complex and enhanced fluorescence quenching of zinc porphyrin. To further understand the intramolecular quenching mechanism and follow the kinetics of photoinduced processes occurring in these dyads and triads, picosecond time-resolved emission, as well as subpicosecond and nanosecond transient absorption studies, have been performed.

**Fluorescence Lifetimes.** The time-resolved fluorescence spectral features of the dyads track that of the steady state measurements. The fluorescence decay-time profiles of the investigated dyads of ZnP (monitored at 600 nm) and H<sub>2</sub>P (monitored at 650 nm) show an enhanced decay rate as compared with those of (TPP)Zn and (TPP)H<sub>2</sub> in both DCB and in BN (Figure 7). In DCB, all of the investigated compounds

**TABLE 2: Electrochemical Half-Wave Redox Potentials (vs Fc/Fc<sup>+</sup>) of the Porphyrin–Fulleropyrrolidine Dyads in DCB, 0.1 M (TBA)ClO<sub>4</sub>**

| compd                             | potential (V vs Fc/Fc <sup>+</sup> ) |                  |                                |                                 |                  |                                  |                   | HOMO–LUMO, V <sup>a</sup> | ΔG <sub>cs</sub> <sup>o</sup> , V <sup>b</sup> |
|-----------------------------------|--------------------------------------|------------------|--------------------------------|---------------------------------|------------------|----------------------------------|-------------------|---------------------------|------------------------------------------------|
|                                   | P <sup>+/2+</sup>                    | P <sup>0/+</sup> | C <sub>60</sub> <sup>0/-</sup> | C <sub>60</sub> <sup>-2/-</sup> | P <sup>0/-</sup> | C <sub>60</sub> <sup>2-/3-</sup> | P <sup>-2/-</sup> |                           |                                                |
| ZnP                               | 0.62                                 | 0.28             |                                |                                 | -1.92            |                                  | -2.23             |                           |                                                |
| ZnPo-C <sub>60</sub>              | 0.63                                 | 0.29             | -1.17                          | -1.55                           | -1.90            | -2.07                            | -2.29             | 1.46                      | -0.84                                          |
| ZnPp-C <sub>60</sub>              | 0.61                                 | 0.27             | -1.18                          | -1.55                           | -1.91            | -2.06                            | -2.25             | 1.45                      | -0.86                                          |
| Py:ZnPo-C <sub>60</sub> C         | 0.61                                 | 0.26             | -1.18                          | -1.56                           | -1.91            | -2.07                            | -2.29             | 1.44                      | -0.83                                          |
| Py:ZnPp-C <sub>60</sub> C         | 0.60                                 | 0.26             | -1.18                          | -1.56                           | -1.92            | -2.07                            | -2.26             | 1.44                      | -0.84                                          |
| H <sub>2</sub> P                  | 0.79                                 | 0.53             |                                |                                 | -1.75            |                                  | -2.05             |                           |                                                |
| H <sub>2</sub> Po-C <sub>60</sub> | 0.81                                 | 0.50             | -1.18                          | -1.56                           | -1.73            | -2.10 <sup>d</sup>               |                   | 1.68                      | -0.44                                          |
| H <sub>2</sub> Pp-C <sub>60</sub> | 0.80                                 | 0.51             | -1.19                          | -1.55                           | -1.74            | -2.08 <sup>d</sup>               |                   | 1.69                      | -0.44                                          |

<sup>a</sup> HOMO–LUMO = P<sup>0/+</sup> - C<sub>60</sub><sup>0/-</sup>. <sup>b</sup> ΔG<sub>cs</sub><sup>o</sup> = E<sub>1/2</sub>(D<sup>+/+</sup>/D) - E<sub>1/2</sub>(A/A<sup>+</sup>) - ΔE<sub>0-0</sub> + ΔG<sub>s</sub> where E<sub>1/2</sub>(D<sup>+/+</sup>/D) is the first one electron oxidation potential of the donor porphyrin, E<sub>1/2</sub>(A/A<sup>+</sup>) is the first one electron reduction potential of the C<sub>60</sub> electron acceptor, ΔE<sub>0-0</sub> is the energy of the 0–0 transition energy gap between the lowest excited state and the ground state, and ΔG<sub>s</sub> refers to the solvation energy, calculated by using the “dielectric continuum model” according to the following equation: ΔG<sub>s</sub> = e<sup>2</sup>/4πε<sub>0</sub>[(1/2R<sub>+</sub> + 1/2R - 1/R<sub>D-A</sub>)(1/ε<sub>s</sub>) - (1/2R<sub>+</sub> + 1/2R)(1/ε<sub>R</sub>)]. From Figure 2, values of R<sub>+</sub> = 4.8 Å, R<sub>-</sub> = 4.2 Å, R<sub>D-A</sub> = 6.7 Å for *ortho* and R<sub>D-A</sub> = 6.4 Å for *para* derivatives were employed. <sup>c</sup> Obtained by addition of 3 equiv of pyridine to the ZnP-C<sub>60</sub> dyads. <sup>d</sup> Overlap of C<sub>60</sub><sup>2-/3-</sup> and P<sup>-2/-</sup> processes.



**Figure 5.** Ab initio B3LYP/3-21G(\*)-calculated frontier (a) HOMO-1, (b) HOMO, (c) LUMO, (d) LUMO+1, (e) LUMO+2, (f) LUMO+3, (g) LUMO+4, and (h) LUMO+5 orbitals of ZnPp-C<sub>60</sub>.

**TABLE 3: Fluorescence Lifetimes ( $\tau_f$ )<sup>a</sup>, Charge Separation Rate Constants ( $k_{cs}^{singlet}$ ), Charge Separation Quantum Yields ( $\Phi_{cs}^{singlet}$ ), and Charge Recombination Rate Constants ( $k_{cr}$ ) for Porphyrin-C<sub>60</sub> Dyads in DCB and BN Solvents**

| compd                             | solvent         | $\tau_f$ (ns) (fraction %) | $k_{cs}^{singlet}$ (s <sup>-1</sup> ) <sup>b</sup> | $\Phi_{cs}^{singlet}$ <sup>b</sup> | $k_{cr}$ (s <sup>-1</sup> ) | $k_{cs}/k_{cr}$ |
|-----------------------------------|-----------------|----------------------------|----------------------------------------------------|------------------------------------|-----------------------------|-----------------|
| ZnPp-C <sub>60</sub>              | DCB             | 0.33 (100)                 | $2.6 \times 10^9$                                  | 0.84                               | $5.9 \times 10^7$           | 44              |
|                                   | BN              | 0.26 (100)                 | $3.4 \times 10^9$                                  | 0.88                               | $1.8 \times 10^6$           | 1888            |
| Py:ZnPp-C <sub>60</sub> C         | DCB             | 0.29 (100)                 | $3.0 \times 10^9$                                  | 0.86                               | $3.4 \times 10^6$           | 882             |
|                                   | Py <sup>d</sup> | 0.23 (100)                 | $3.9 \times 10^9$                                  | 0.89                               | $1.7 \times 10^6$           | 2294            |
| ZnPpo-C <sub>60</sub>             | DCB             | 0.70 (100)                 | $9.5 \times 10^8$                                  | 0.68                               | $>6.0 \times 10^7$          | 15              |
|                                   | BN              | 0.64 (100)                 | $1.1 \times 10^9$                                  | 0.69                               | $3.0 \times 10^6$           | 366             |
| Py:ZnPp-C <sub>60</sub> C         | DCB             | 0.61 (100)                 | $1.1 \times 10^9$                                  | 0.71                               | $5.5 \times 10^6$           | 212             |
|                                   | Py <sup>d</sup> | 0.54 (100)                 | $1.3 \times 10^9$                                  | 0.74                               | $2.5 \times 10^6$           | 660             |
| H <sub>2</sub> P-C <sub>60</sub>  | DCB             | 0.87 (100)                 | $1.1 \times 10^9$                                  | 0.94                               | $4.3 \times 10^7$           | 26              |
|                                   | BN              | 0.44 (60) <sup>e</sup>     | $2.2 \times 10^9$                                  | 0.97                               | $4.4 \times 10^7$           | 50              |
| H <sub>2</sub> Pp-C <sub>60</sub> | DCB             | 3.33 (100)                 | $2.3 \times 10^8$                                  | 0.75                               | $1.7 \times 10^7$           | 14              |
|                                   | BN              | 2.70 (100)                 | $3.0 \times 10^8$                                  | 0.81                               | $2.3 \times 10^7$           | 13              |


<sup>a</sup> The  $\tau_f$  for reference compounds: (TPP)Zn = 2.1 ns and (TPP)H<sub>2</sub> = 13.6 ns in DCB and BN. <sup>b</sup>  $k_{cs}^{singlet} = (1/\tau_f)_{sample} - (1/\tau_f)_{ref}$ . <sup>c</sup>  $\Phi_{cs}^{singlet} = [(1/\tau_f)_{sample} - (1/\tau_f)_{ref}] / (1/\tau_f)_{sample}$ . For biexponential fitting,  $\tau_f$  from the initial decay component was employed. <sup>d</sup> Obtained by the addition of 0.1 mM pyridine. <sup>e</sup> In neat pyridine. <sup>f</sup> 0.97 ns (40%).

revealed a monoexponential decay, while in BN the decay of H<sub>2</sub>Po-C<sub>60</sub> could be fitted satisfactory to a biexponential decay. Substantial quenching of the fluorescence lifetimes is observed for the investigated dyads; the *ortho* derivatives clearly show a higher efficiency of quenching than those of *para* derivatives.

By assuming that the quenching is due to electron transfer from the singlet excited porphyrin to the fullerene, the rates of charge separation ( $k_{cs}^{singlet}$ ) and quantum yields ( $\Phi_{cs}^{singlet}$ ) are evaluated in an usual manner employed for the intra-molecular electron-transfer process and the data are given in Table 3. An examination of Table 3 reveals the following: (i) The experimentally determined  $k_{cs}^{singlet}$  and  $\Phi_{cs}^{singlet}$  are higher for *ortho*-substituted derivatives as compared to the *para*-substituted derivatives. This observation is consistent with the


close proximity of the porphyrin and fullerene entities in the *ortho*-substituted derivatives obtained from the fluorescence spectral and computational methods. (ii) The  $k_{cs}^{singlet}$  is higher for ZnP-C<sub>60</sub> as compared with H<sub>2</sub>P-C<sub>60</sub>, which is in agreement with the electrochemical redox potentials and the calculated free-energy change,  $\Delta G_{cs}^0$ , for photoinduced electron-transfer reactions. This considers the reorganization energy ( $\lambda$ ) to be close to the  $\Delta G_{cs}^0$  values of ZnP-C<sub>60</sub> and the  $\lambda$  values between the  $\Delta G_{cs}^0$  values of ZnP-C<sub>60</sub> and H<sub>2</sub>Po-C<sub>60</sub>. (iii) In agreement with the steady state emission results, the  $k_{cs}^{singlet}$  and  $\Phi_{cs}^{singlet}$  are higher in more polar BN solvent as compared to the values obtained in less polar DCB solvent.

Interestingly, upon addition of Py (10 equiv) to a solution of either ZnPp-C<sub>60</sub> or ZnPp-C<sub>60</sub> in DCB to form the penta-




**Figure 6.** Fluorescence spectrum of (a) (i) (TPP)Zn, (ii) ZnPp-C<sub>60</sub>, and (iii) ZnPo-C<sub>60</sub> in DCB ( $\lambda_{\text{ex}}$  = 550 nm) and (b) (i) (TPP)H<sub>2</sub>, (ii) H<sub>2</sub>Pp-C<sub>60</sub>, and (iii) H<sub>2</sub>Po-C<sub>60</sub> in DCB ( $\lambda_{\text{ex}}$  = 515 nm).

coordinated species, Py:ZnPo-C<sub>60</sub> or Py:ZnPp-C<sub>60</sub>, the fluorescence decay of ZnP revealed additional quenching while maintaining the monoexponential decay. Accordingly, the calculated  $k_{\text{cs}}^{\text{singlet}}$  and  $\Phi_{\text{cs}}^{\text{singlet}}$  are found to be slightly higher for the pyridine-coordinated dyads as compared to the corresponding dyads in the absence of any axial ligand (Table 3). These higher values of  $k_{\text{cs}}^{\text{singlet}}$  and  $\Phi_{\text{cs}}^{\text{singlet}}$  have been attributed to the easier oxidation of the penta-coordinated zinc porphyrin in the dyads (vide supra). Further increases in fluorescence decay rates were observed in neat pyridine, suggesting that all of the ZnPo-C<sub>60</sub> or ZnPp-C<sub>60</sub> are coordinated by excess pyridine, resulting in further increases in  $k_{\text{cs}}^{\text{singlet}}$  and  $\Phi_{\text{cs}}^{\text{singlet}}$ . Further studies involving subpicosecond and nanosecond transient absorption technique have been performed to monitor the charge separation and charge recombination processes in the dyads and self-assembled triads. The forthcoming sections discuss the results of these studies.




**Figure 7.** Fluorescence decay profiles of (a) (TPP)Zn, ZnPp-C<sub>60</sub>, and ZnPo-C<sub>60</sub> and (b) (TPP)H<sub>2</sub>, H<sub>2</sub>Pp-C<sub>60</sub>, and H<sub>2</sub>Po-C<sub>60</sub> in BN;  $\lambda_{\text{ex}}$  = 410 nm. The concentrations of porphyrins were maintained at 0.05 mM.



**Figure 8.** Upper panel: picosecond transient absorption spectra obtained by the 455 nm laser light excitation of H<sub>2</sub>Po-C<sub>60</sub> (0.1 mM) in BN. Lower panel: time profile at 455 nm.

**Subpicosecond Time-Resolved Transient Absorption Spectral Studies.** To confirm the occurrence of a charge separation process, the transient absorption spectra were monitored during the first 1000 ps after pulsing the samples by a 150 fs laser light. Figure 8 illustrates the spectral profiles obtained for ZnPo-C<sub>60</sub> in BN as a representative example. Immediately after the laser pulse, an intense absorption band appeared at 455 nm (spectrum at 0 ps; immediately after 150 fs laser pulse), which is attributed to the  $S_1 \rightarrow S_n$  transition of the ZnP moiety. The decay rate of this  $S_1$  band shown in the lower panel is about  $5 \times 10^9 \text{ s}^{-1}$ , which is slightly higher than that evaluated from the fluorescence lifetime measurements ( $3.4 \times 10^9 \text{ s}^{-1}$ ) but are within the estimated experimental error. As shown in the figure at the 300 ps regime, the decay of the  $S_1$  band of the ZnP moiety is accompanied by an appearance of an absorption band at 600 nm corresponding to the formation of ZnP<sup>+</sup>. In the longer wavelength region, a new absorption band at 1000 nm is also observed in this time interval corresponding to the formation of radical anion of the C<sub>60</sub> moiety. In DCB, weaker absorption was observed at the 600 and 1000 nm regions. On addition of pyridines to form the penta-coordinated complex in DCB,



**Figure 9.** Nanosecond transient absorption spectra obtained using 532 nm laser excitation of H<sub>2</sub>Po-C<sub>60</sub> (0.1 mM) in Ar-saturated DCB. Inset: time profiles for the 700 and 1000 nm bands.


spectral features similar to that shown in Figure 8 were observed. For the H<sub>2</sub>P-C<sub>60</sub> dyads both in DCB and in BN, similar transient spectra were observed in the picosecond region thus confirming the occurrence of charge separation from the singlet excited porphyrin to the covalently linked fullerene moiety.

**Nanosecond Time-Resolved Absorption Spectra of the Dyads in DCB.** The nanosecond transient absorption spectrum of pristine ZnP exhibited absorption peaks at 630 and 840 nm corresponding to the triplet state.<sup>18,19</sup> Similarly, pristine H<sub>2</sub>P exhibited transient absorption bands at 620 and 800 nm corresponding to its triplet state.<sup>18,19</sup> The transient absorption spectrum of 2-(4'-pyridyl)fulleropyrrolidine revealed a band around 700 nm corresponding to its triplet state.<sup>18</sup> The anion radical of 2-(4'-pyridyl)fulleropyrrolidine is expected to reveal a band around 1000 nm, while the cation radicals of H<sub>2</sub>P and ZnP are expected to appear in the 620–630 nm range.

The nanosecond transient absorption spectrum of H<sub>2</sub>Po-C<sub>60</sub> in DCB showed a fast decaying peak of the radical anion of the C<sub>60</sub> moiety around 1000 nm in addition to a slow decaying band at 700 nm corresponding to the triplet C<sub>60</sub> and 620 and 800 nm bands corresponding to the triplet state of the H<sub>2</sub>P (Figure 9).<sup>19</sup> A part of the transient absorption band at 626 nm also showed quick decay, and this has been attributed to the formation of H<sub>2</sub>P<sup>•+</sup>. The decay of the 1000 nm transient band corresponds to the rate of the charge recombination ( $k_{cr}$ ), and the values of  $k_{cr}$  thus calculated are given in the last column in Table 3.

The nanosecond transient absorption spectrum of ZnPo-C<sub>60</sub> in DCB is quite similar to Figure 9 for H<sub>2</sub>Po-C<sub>60</sub>, showing a fast decaying peak of the radical anion of the C<sub>60</sub> around 1000 nm ( $k_{cr} = 5.8 \times 10^7 \text{ s}^{-1}$ ) in addition to the slow decaying 700 nm band of the triplet C<sub>60</sub> and 850 nm band corresponding to the triplet state of the zinc porphyrin. In the case of ZnPp-C<sub>60</sub>, the transient absorption spectra also showed the 1000 nm band ( $k_{cr} \geq 6 \times 10^7 \text{ s}^{-1}$ ) in addition to the triplet states of C<sub>60</sub> and ZnP. A comparison between the  $k_{cr}$  values in DCB suggests lower rates for H<sub>2</sub>P-C<sub>60</sub> dyads as compared to the ZnP-C<sub>60</sub> dyads, indicating longer lifetimes of the charge-separated state of H<sub>2</sub>P-C<sub>60</sub> dyads. This suggests that the  $\Delta G_{cr}^\circ$  values for both H<sub>2</sub>P-C<sub>60</sub> and ZnP-C<sub>60</sub> dyads are in the inverted region and that  $\Delta G_{cr}^\circ$  values of ZnP-C<sub>60</sub> dyads are near the top region of the Marcus parabola.

**Nanosecond Time-Resolved Absorption Spectra of the Py:ZnP-C<sub>60</sub> in Py and DCB.** Figure 10 shows the nanosecond transient absorption spectra of Py:ZnP-C<sub>60</sub> in neat pyridine. For the penta-coordinated complex, the lifetime of the ion pair states increased considerably. From the oxidation potential of




**Figure 10.** Nanosecond transient absorption spectra obtained using 532 nm laser excitation of ZnPp-C<sub>60</sub> (0.1 mM) in Ar-saturated neat pyridine. Inset: time profile for the 1000 nm band.

Py:ZnP-C<sub>60</sub>, the hole (cation radical of ZnP moiety) generated by the initial charge separation is likely to delocalize to over the pyridine entity, thus slowing down the charge recombination process due to an increased distance between the C<sub>60</sub> radical anion and the radical cation of Py:ZnP. The spectrum at 100 ns indicates that all of the transient species observed immediately after the nanosecond laser light pulse decayed simultaneously within 1.0  $\mu\text{s}$ . As expected, charge stabilization in neat pyridine is found to be much more pronounced as revealed by the higher  $k_{cs}/k_{cr}$  values as compared with addition of pyridine in DCB (Table 3). Utilization of substituted pyridines as axial ligands revealed similar observations. Smaller  $k_{cr}$  values were obtained when 4-hydroxypyridine or 4-acetylpyridine was used as an axial ligand ( $1.1 \times 10^6 \text{ s}^{-1}$  or  $1.4 \times 10^6 \text{ s}^{-1}$  for Py:ZnPp-C<sub>60</sub> in DCB, respectively) while for 4-dimethylamino pyridine with an electron-donating group the  $k_{cr}$  values ( $2.2 \times 10^6 \text{ s}^{-1}$  for Py:ZnPp-C<sub>60</sub> in DCB) were found to be almost the same as that obtained for pyridine in DCB.

**Nanosecond Time-Resolved Absorption Spectra of the Dyads in BN.** In BN, the transient absorption spectrum of ZnPp-C<sub>60</sub> (see Supporting Information for figure) showed a clear absorption band of the radical anion of the C<sub>60</sub> moiety at 1000 nm with a longer lifetime of the ion pair state (330 ns), which is 20 times longer than that in DCB. Similarly, the lifetime of the ion pair state of ZnPo-C<sub>60</sub> in BN is found to be 550 ns, which is 33 times longer than that in DCB. The transient spectrum recorded at 100 ns for the ZnP-C<sub>60</sub> dyads revealed the existence of a weak absorption at 620 nm, which has been ascribed to the radical cation of the ZnP moiety. It is also noticeable that the triplet state absorption of the C<sub>60</sub> moiety at 700 nm decays more quickly in BN than that in DCB; the decay rates of the triplet states are also increased and are comparable to the decay rates of the ion pairs. This could be due to a mixing of the radical ion pair states with the triplet states, which would mean the existence of the radical ion pair character of the triplet state.<sup>20</sup> From the similarity of the  $k_{cr}$  values in BN to those in pyridine, the observed longer lifetime of the radical ion pairs in BN could be ascribed to the coordination of BN to the ZnP moiety.

**Energy Level Diagrams.** The results of the present investigation are summarized with the help of an energy level diagram shown in Figure 11. By utilizing the oxidation and reduction potentials and the calculated distance between the donor and the acceptor, the free energies ( $\Delta G^\circ$ ) for light-induced electron transfer were estimated for constructing the energy level diagram. On the basis of these  $\Delta G^\circ$  values and the reported energies for the singlet and triplet states, the energy levels in



**Figure 11.** Energy level diagrams showing the different photochemical events of (a)  $\text{H}_2\text{P}-\text{C}_{60}$  dyads and (b)  $\text{ZnP}-\text{C}_{60}$  dyads in DCB. The energy of  $(\text{Py}:\text{ZnP})^{+\bullet}-\text{C}_{60}^{-\bullet}$  state in DCB (1.22 eV) and  $(\text{ZnP})^{+\bullet}-\text{C}_{60}^{-\bullet}$  state in BN (1.12 eV) is shown as open rectangles.

Figure 11 showing the different photochemical processes of the tetra- and penta-coordinated dyads reveal the following. The main difference in the  $k_{\text{CS}}$  between  $\text{H}_2\text{P}-\text{C}_{60}$  and  $\text{ZnP}-\text{C}_{60}$  is the energy differences between the singlet excited state of the porphyrin moiety and the radical ion pair states; the reorganization energy ( $\lambda$ ) is rather near  $\Delta G_{\text{CS}}^0$  values of  $\text{ZnP}-\text{C}_{60}$  when the top region of the Marcus parabola exists between  $\Delta G_{\text{CS}}^0$  values of  $\text{ZnP}-\text{C}_{60}$  and  $\text{H}_2\text{P}-\text{C}_{60}$ . The enhanced charge separation in coordinating solvents, pyridine and BN, as compared to that in less polar DCB for the dyads is noteworthy. This trend is not reasonable for  $\text{ZnP}-\text{C}_{60}$  dyads belonging to the Marcus inverted region. The prolonged charge recombination processes observed for the radical ion pair of  $\text{ZnP}-\text{C}_{60}$  dyads in BN and not for the  $\text{H}_2\text{P}-\text{C}_{60}$  dyads may be attributed to the coordination to Zn, which renders the relative position of the radical ion pair center apart from the radical anion center,  $\text{C}_{60}$ . In addition, in the case of  $\text{ZnP}-\text{C}_{60}$  dyads, the radical ion pair states are lower than the triplet states, while the radical ion pair state of  $\text{H}_2\text{P}-\text{C}_{60}$  is higher than the triplet states even in BN; thus, the charge recombination yielding the triplet states is efficient for  $\text{H}_2\text{P}-\text{C}_{60}$ .

Efficient charge separation and the slower charge recombination processes have been observed upon axial coordination of pyridine to the  $\text{ZnP}-\text{C}_{60}$  dyads. The higher rates of charge separation can be attributed to the easier oxidation of the penta-coordinated  $\text{ZnP}$ , while the slower charge recombination can be attributed to the longer distance between radical anion and radical cation caused by the coordinated pyridine ligand; delocalization of the  $\pi$ -cation radical of  $\text{ZnP}$  to the coordinated pyridine ligand and the longer space between them stabilizes the radical ion pair state. More importantly, an evaluation of the  $k_{\text{CS}}/k_{\text{CR}}$  ratio, a measure of the excellence in the photoinduced electron-transfer systems, is more than 1000 for  $\text{ZnP}-\text{C}_{60}$  in neat pyridine or in BN.

## Experimental Section

**Chemicals.** Buckminsterfullerene,  $\text{C}_{60}$  (+99.95%), was from SES Research, (Houston, TX). DCB and BN in sure seal bottles, sarcosine, pyrrole, benzaldehyde derivatives (benzaldehyde, 4-bromoethoxybenzaldehyde, and *o*- and *p*-hydroxybenzaldehyde), and pyridine derivatives (pyridine, 4-hydroxypyridine, 4-dimethylaminopyridine, and 4-acetylpyridine) were from Aldrich Chemicals (Milwaukee, WI). Tetra-*n*-butylammonium perchlorate,  $(\text{TBA})\text{ClO}_4$ , was from Fluka Chemicals. All chemicals were used as received. Syntheses and purification of  $(\text{TPP})\text{H}_2$  and  $(\text{TPP})\text{Zn}$  were carried out according to the literature procedure.<sup>14</sup>

**Synthesis of 5-(2' or 4'-Hydroxyphenyl)-10,15,20-triphenylporphyrin.** The title compounds were synthesized by reacting 2'- or 4'-hydroxybenzaldehyde (1.5 g, 12 mM), benzaldehyde (3.9 g, 37 mM), and pyrrole (3.3 g, 49 mM) in refluxing propionic acid.<sup>11f</sup> The crude product was purified on a basic alumina column with chloroform/methanol (95:5 v/v) as eluent. Yield ~5%.  $^1\text{H}$  NMR in  $\text{CDCl}_3$ :  $\delta$  *ortho*-hydroxy: 8.89 (m, 8H,  $\beta$ -pyrrole-H), 8.27 (m, 6H, *ortho*-phenyl-H), 7.79 (m, 9H, *meta*- and *para*-phenyl-H), 8.08–7.21 (m, 4H, substituted phenyl-H), 5.35 (s, br, 1H, hydroxy-H), –2.81 (s, 2H, imino-H). *para*-hydroxy: 8.89 (m, 8H,  $\beta$ -pyrrole-H), 8.27 (m, 6H, *ortho*-phenyl-H), 7.79 (m, 9H, *meta*- and *para*-phenyl-H), 8.08–7.21 (d,d, 4H, substituted phenyl-H), 5.35 (s (br), 1H, hydroxy-H), –2.81 (s, 2H, imino-H).

**Synthesis of  $\text{H}_2\text{Po}-\text{CHO}$  and  $\text{H}_2\text{Pp}-\text{CHO}$ .** These were synthesized by stirring a solution of the respective hydroxy-derivatized porphyrins (0.1 mmol in 100 mL of DMF) and 4-(2-bromoethoxy)benzaldehyde (0.5 mmol) in the presence of  $\text{K}_2\text{CO}_3$  for 12 h. At the end of the reaction, the DMF was evaporated and the compound was extracted in  $\text{CHCl}_3$ . Further purification of the compounds was carried out on a silica gel column using toluene and chloroform (95:5 v/v) as eluent. Yield 45%.  $^1\text{H}$  NMR in  $\text{CDCl}_3$ :  $\delta$   $\text{H}_2\text{Po}-\text{CHO}$ : 10.03 (s, 1H, aldehyde-H), 8.83 (m, 8H,  $\beta$ -pyrrole-H), 8.16 (m, 6H, *ortho*-phenyl-H), 7.74 (m, 9H, *meta*- and *para*-phenyl-H), 8.14–7.41 (m, 4H, substituted phenyl-H), 5.96 (d, 2H, phenyl-H), 5.53 (d, 2H, phenyl-H), 4.21, 3.51 (t,t, 2H,2H,  $\text{CH}_2-\text{CH}_2$ ), –2.84 (s, 2H, imino-H).  $\text{H}_2\text{Pp}-\text{CHO}$ : 9.90 (s, 1H, aldehyde-H), 8.84 (m, 8H,  $\beta$ -pyrrole-H), 8.21 (m, 6H, *ortho*-phenyl-H), 7.84 (m, 9H, *meta*- and *para*-phenyl-H), 7.76–7.12 (d,d, 4H, substituted phenyl-H), 4.61 (m, 4H, phenyl-H), 4.39, 3.77 (t,t, 2H,2H,  $\text{CH}_2-\text{CH}_2$ ), –2.75 (s, 2H, imino-H).

**Synthesis of  $\text{H}_2\text{Po}-\text{C}_{60}$  and  $\text{H}_2\text{Pp}-\text{C}_{60}$ .** These were synthesized by refluxing a mixture of the respective  $\text{H}_2\text{P}-\text{CHO}$  derivatives (0.045 mmol),  $\text{C}_{60}$  (0.24 mmol), and sarcosine (0.24 mmol) in 100 mL of toluene for 10 h. At the end, the solvent was evaporated and the compound was purified on a silica gel column using toluene and hexane (95:5 v/v) as eluent. Yield

56.4 mg (~80%). <sup>1</sup>H NMR in CDCl<sub>3</sub>: δ H<sub>2</sub>Po-C<sub>60</sub>: 8.76 (m, 8H, β-pyrrole-H), 8.16 (m, 6H, *ortho*-phenyl-H), 7.72 (m, 9H, *meta*- and *para*-phenyl-H), 7.97–7.14 (d,d,t, 4H, substituted phenyl-H), 5.94 (s, 4H, phenyl-H), 4.15, 3.48 (t, t, 2H, 2H, CH<sub>2</sub>-CH<sub>2</sub>), 4.63, 3.77 (d,d, 2H, pyrrolidine-H), 3.80 (s, 1H, pyrrolidine-H), 2.34 (s, 3H, pyrrolidine N-CH<sub>3</sub>), -2.77 (s.br, 2H, imino-H). ESI mass in CH<sub>2</sub>Cl<sub>2</sub>: calcd, 1526; found, 1526.1. UV-visible in CH<sub>2</sub>Cl<sub>2</sub>: λ<sub>max</sub> 308, 326 (sh), 418, 515, 549, 591, 648. H<sub>2</sub>Pp-C<sub>60</sub>: 8.83 (m, 8H, β-pyrrole-H), 8.19 (m, 6H, *ortho*-phenyl-H), 7.75 (m, 9H, *meta*- and *para*-phenyl-H), 8.04–7.15 (d,d, 4H, substituted phenyl-H), 7.49, 7.06 (m, 4H, phenyl-H), 4.57, 4.49 (t,t, 2H, 2H, CH<sub>2</sub>-CH<sub>2</sub>), 4.88, 4.14 (d,d, 2H, pyrrolidine-H), 4.80 (s, 1H, pyrrolidine-H), 2.32 (s, 3H, pyrrolidine N-CH<sub>3</sub>), -2.81 (s.br, 2H, imino-H). ESI mass in CH<sub>2</sub>-Cl<sub>2</sub>: calcd, 1526; found, 1527.1. UV-visible in CH<sub>2</sub>Cl<sub>2</sub>: λ<sub>max</sub> 306, 325 (sh), 418, 515, 551, 590, 647.

**Synthesis of ZnPo-C<sub>60</sub> and ZnPp-C<sub>60</sub>.** These were synthesized by metalation of the respective free-base porphyrin derivatives (0.037 mmol) in CHCl<sub>3</sub> by using excess of zinc acetate in methanol. The course of the reaction was followed spectroscopically. At the end (~1 h), the solvent was evaporated and the product was purified on silica gel column using toluene as eluent. Yield 93%. <sup>1</sup>H NMR in CDCl<sub>3</sub>: δ ZnPo-C<sub>60</sub>: 8.88 (m, 8H, β-pyrrole-H), 8.18 (m, 6H, *ortho*-phenyl-H), 7.71 (m, 9H, *meta*- and *para*-phenyl-H), 7.96–7.15 (m, 4H, substituted phenyl-H), 5.91 (s, 4H, phenyl-H), 4.15, 3.51 (t,t, 2H, 2H, CH<sub>2</sub>-CH<sub>2</sub>), 4.62, 3.78 (d,d, 2H, pyrrolidine-H), 3.85 (s, 1H, pyrrolidine-H), 2.33 (s, 3H, pyrrolidine N-CH<sub>3</sub>). ESI mass in CH<sub>2</sub>Cl<sub>2</sub>: calcd, 1589.4; found, 1608.5 (M + H<sub>2</sub>O). UV-visible in CH<sub>2</sub>Cl<sub>2</sub>: λ<sub>max</sub> 309, 328 (sh), 421, 548, 584 (weak). ZnPp-C<sub>60</sub>: 8.92 (m, 8H, β-pyrrole-H), 8.20 (m, 6H, *ortho*-phenyl), 7.67 (m, 9H, *meta*- and *para*-phenyl-H), 8.04–7.11 (d,d, 4H, substituted phenyl-H), 7.29 (m, 4H, phenyl-H), 4.62, 4.53 (t,t, 2H, 2H, CH<sub>2</sub>-CH<sub>2</sub>), 4.92, 4.05 (d,d, 2H, pyrrolidine-H), 4.85 (s, 1H, pyrrolidine-H), 2.79 (s, 3H, pyrrolidine N-CH<sub>3</sub>). ESI mass in CH<sub>2</sub>Cl<sub>2</sub>: calcd, 1589.4; found, 1608.7 (M + H<sub>2</sub>O). UV-visible in CH<sub>2</sub>Cl<sub>2</sub>: λ<sub>max</sub> 308, 328 (sh), 421, 548, 585 (weak).

**Instrumentation.** The UV-visible spectral measurements were carried out with a Shimadzu model 1600 UV-visible spectrophotometer. The fluorescence emission was monitored by using a Spex Fluorolog-τ spectrometer. A right angle detection method was used. The <sup>1</sup>H NMR studies were carried out on a Varian 400 MHz spectrometer. Tetramethylsilane (TMS) was used as an internal standard. Cyclic voltammograms were recorded on a EG&G model 263A potentiostat using a three electrode system. A platinum button or glassy carbon electrode was used as the working electrode. A platinum wire served as the counter electrode, and a Ag wire (Ag/Ag<sup>+</sup>) was used as the reference electrode. Ferrocene/ferrocenium redox couple was used as an internal standard. All of the solutions were purged prior to electrochemical and spectral measurements using argon gas. The computational calculations were performed by ab initio B3LYP/3-21G(\*) methods with GAUSSIAN 98<sup>21</sup> software package on various PCs and a SGI ORIGIN 2000 computer. The graphics of HOMO and LUMO coefficients were generated with the help of GaussView software. The ESI-mass spectral analyses of the newly synthesized compounds were performed by using a Fennigan LCQ-Deca mass spectrometer. For this, the compounds (about 1 mM concentration) were prepared in CH<sub>2</sub>Cl<sub>2</sub>, freshly distilled over calcium hydride.

**Time-Resolved Emission and Transient Absorption Measurements.** The picosecond time-resolved fluorescence spectra were measured using an argon-ion-pumped Ti:sapphire laser

(Tsunami) and a streak scope (Hamamatsu Photonics). The details of the experimental setup are described elsewhere.<sup>20</sup> The subpicosecond transient absorption spectra were recorded by the pump and probe method. The samples were excited with a second harmonic generation (SHG, 388 nm) output from a femtosecond Ti:sapphire regenerative amplifier seeded by SHG of a Er-dropped fiber (Clark-MXRCPA-2001 plus, 1 kHz, fwhm 150 fs). The excitation light was depolarized. The monitor white light was generated by focusing the fundamental of laser light on flowing D<sub>2</sub>O/H<sub>2</sub>O cell. The transmitted monitor light was detected with a dual MOS linear image sensor (Hamamatsu Photonics, C6140) or InGaAs photodiode array (Hamamatsu Photonics, C5890-128). Nanosecond transient absorption spectra in the NIR region were measured by means of laser flash photolysis; 532 nm light from a Nd:YAG laser was used as the exciting source and a Ge-avalanche-photodiode module was used for detecting the monitoring light from a pulsed Xe lamp as described in our previous paper.<sup>20</sup>

**Acknowledgment.** We are thankful to the donors of the Petroleum Research Fund administered by the American Chemical Society, National Institutes of Health (to F.D.), Japan Ministry of Education, Science, Technology, Culture and Sports, and Mitsubishi Foundation (to O.I.) for support of this work. We are also thankful to the High Performance Computing Center of the Wichita State University for lending SGI ORIGIN 2000 computer time.

**Supporting Information Available:** Ab initio B3LYP/3-21G(\*)-calculated frontier HOMO and LUMO of ZnPo-C<sub>60</sub> and H<sub>2</sub>Po-C<sub>60</sub>. Nanosecond transient absorption spectrum of ZnPp-C<sub>60</sub> in BN. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## References and Notes

- (1) (a) Maruyama, K.; Osuka, A. *Pure Appl. Chem.* **1990**, 62, 1511. (b) Gust, D.; Moore, T. A. *Science* **1989**, 244, 35. (c) Gust, D.; Moore, T. A. *Top. Curr. Chem.* **1991**, 159, 103. (d) Gust, D.; Moore, T. A.; Moore, A. L. *Acc. Chem. Res.* **1993**, 26, 198. (e) Wasielewski, M. R. *Chem. Rev.* **1992**, 92, 435. (f) Paddon-Row, M. N. *Acc. Chem. Res.* **1994**, 27, 18. (g) Sutin, N. *Acc. Chem. Res.* **1983**, 15, 275. (h) Bard, A. J.; Fox, M. A. *Acc. Chem. Res.* **1995**, 28, 141. (i) Meyer, T. J. *Acc. Chem. Res.* **1989**, 22, 163. (j) Piotrowiak, P. *Chem. Soc. Rev.* **1999**, 28, 143.
- (2) (a) Ward, M. W. *Chem. Soc. Rev.* **1997**, 26, 365. (b) Hayashi, T.; Ogoshi, H. *Chem. Soc. Rev.* **1997**, 26, 355. (c) Sessler, J. S.; Wang, B.; Springs, S. L.; Brown, C. T. In *Comprehensive Supramolecular Chemistry*; Atwood, J. L.; Davies, J. E. D.; MacNicol, D. D.; Vögtle, F., Eds.; Pergamon: Elmsford, NY, 1996; Chapter 9. (d) Feldheim, D. L.; Keating, C. D. *Chem. Soc. Rev.* **1998**, 27, 1. (e) *Introduction of Molecular Electronics*; Petty, M. C.; Bryce, M. R.; Bloor, D., Eds.; Oxford University Press: New York, 1995. (f) Emmelius, M.; Pawlowski, G.; Vollmann, H. W. *Angew. Chem., Int. Ed. Engl.* **1989**, 28, 1445.
- (3) (a) Kroto, H. W.; Heath, J. R.; O'Brien, S. C.; Curl, R. F.; Smalley, R. E. *Nature* **1985**, 318, 162. (b) Kratschmer, W.; Lamb, L. D.; Fostiropoulos, F.; Huffman, D. R. *Nature* **1990**, 347, 345. (c) *Fullerene and Related Structures*; Hirsch, A., Ed.; Springer: Berlin, 1999; Vol. 199.
- (4) (a) Allemand, P. M.; Koch, A.; Wudl, F.; Rubin, Y.; Diederich, F.; Alvarez, M. M.; Anz, S. J.; Whetten, R. L. *J. Am. Chem. Soc.* **1991**, 113, 1050. (b) Xie, Q.; Perez-Cordero, E.; Echegoyen, L. J. *Am. Chem. Soc.* **1992**, 114, 3978.
- (5) Ajie, H.; Alvarez, M. M.; Anz, S. J.; Beck, R. E.; Diederich, F.; Fostiropoulos, K.; Huffman, D. R.; Kratschmer, W.; Rubin, Y.; Schriver, K. E.; Sensharma, E.; Whetten, R. L. *J. Phys. Chem.* **1990**, 94, 8630.
- (6) (a) Guldi, D. M. *Chem. Commun.* **2000**, 321. (b) Guldi, D. M. *Chem. Soc. Rev.* **2002**, 31, 22.
- (7) Imahori, H.; El-Khouly, M. E.; Fujitsuka, M.; Ito, O.; Sakata, Y.; Fukuzumi, S. *J. Phys. Chem. A* **2001**, 105, 325.
- (8) (a) Khan, S. I.; Oliver, A. M.; Paddon-Row, M. N.; Rubin, Y. J. *Am. Chem. Soc.* **1993**, 115, 4919. (b) Sariciftci, N. S.; Wudl, F.; Heeger, A. J.; Maggini, M.; Scorrano, G.; Prato, M.; Bourassa, J.; Ford, P. C. *Chem. Phys. Lett.* **1995**, 247, 510. (c) Liddell, P. A.; Sumia, J. P.; Macpherson, A. N.; Noss, L.; Seely, G. R.; Clark, K. N.; Moore, A. L.; Moore, T. A.; Gust, D. *Photochem. Photobiol.* **1994**, 60, 537. (d) Williams, R. M.; Zwier, J.

- M.; Verhoever, J. W. *J. Am. Chem. Soc.* **1995**, *117*, 4093. (e) Imahori, H.; Hagiwara, K.; Aoki, M.; Akiyama, T.; Taniguchi, S.; Okada, T.; Shirakawa, M.; Sakata, Y. *J. Am. Chem. Soc.* **1996**, *118*, 11771. (f) Kuciauskas, D.; Lin, S.; Seely, G. R.; Moore, A. L.; Moore, T. A.; Gust, D.; Drovetskaya, T.; Reed, C. A.; Boyd, P. D. W. *J. Phys. Chem.* **1996**, *100*, 15926. (g) Guldi, D. M.; Maggini, M.; Scorrano, G.; Prato, M. *J. Am. Chem. Soc.* **1997**, *119*, 974. (h) Imahori, H.; Yamada, K.; Hasegawa, M.; Taniguchi, S.; Okada, T.; Sakata, Y. *Angew. Chem., Int. Ed. Engl.* **1997**, *36*, 2626. (h) Liddell, P. A.; Kuciauska, D.; Sumida, J. P.; Nash, B.; Nguyen, D.; Moore, A. L.; Moore, T. A.; Gust, D. *J. Am. Chem. Soc.* **1997**, *119*, 1400.
- (9) (a) Dietel, E.; Hirsh, A.; Eichhorn, E.; Rieker, A.; Hackbarth, S.; Roder, B. *Chem. Commun.* **1998**, 1981. (b) Gareis, T.; Kothe, O.; Daub, J. *Eur. J. Org. Chem.* **1998**, 1549. (c) Da Ros, T.; Prato, M.; Guldi, D.; Alessio, E.; Ruzzi, M.; Pasimeni, L.; Carano, M.; Paolucci, F.; Ceroni, P.; Roffia, S. In *Recent Advances in the Chemistry and Physics of Fullerenes and Related Materials*; Kadish, K. M., Ruoff, R. S., Eds.; The Electrochemical Proceedings Series: Pennington, NJ, 1998; p 1074. (d) Martin, N.; Sanchez, L.; Illescas, B.; Perez, I. *Chem. Rev.* **1998**, *98*, 2527. (e) Gareis, T.; Kothe, O.; Daub, J. *Eur. J. Org. Chem.* **1998**, 1549.
- (10) (a) Imahori, H.; Tamaki, K.; Guldi, D. M.; Luo, C.; Fujitsuka, M.; Ito, O.; Sakata, Y.; Fukuzumi, S. *J. Am. Chem. Soc.* **2001**, *123*, 2607. (b) Armaroli, N.; Marconi, G.; Echegoyen, L.; Bourgeois, J.-P.; Diederich, F. *Chem. Eur. J.* **2000**, *6*, 1629. (c) Guldi, D. M.; Luo, C.; Da Ros, T.; Prato, M.; Dietel, E.; Hirsch, A. *Chem. Commun.* **2000**, 375. (d) Luo, C.; Guldi, D. M.; Imahori, H.; Tamaki, K.; Sakata, Y. *J. Am. Chem. Soc.* **2000**, *122*, 6535. (e) Kuciauskas, D.; Liddell, P. A.; Lin, S.; Stone, S. G.; Moore, A. L.; Moore, T. A.; Gust, D. *J. Phys. Chem. B* **2000**, *104*, 4307. (f) Schuster, D. I.; Cheng, P.; Wilson, S. R.; Prokhorenko, V.; Katterie, M.; Holzwarth, A. R.; Braslavsky, S. E.; Klihm, G.; Williams, R. M.; Luo, C. *J. Am. Chem. Soc.* **2000**, *121*, 11599. (g) Tashiro, K.; Aida, T.; Zheng, J.-Y.; Kinbara, K.; Saigo, K.; Sakamoto, S.; Yamaguchi, K. *J. Am. Chem. Soc.* **1999**, *121*, 9477. (h) Tkachenko, N. V.; Rantala, L.; Tauber, A. Y.; Helaja, J.; Hynninen, P. H.; Lemmetyinen, H. *J. Am. Chem. Soc.* **1999**, *121*, 9378. (i) Imahori, H.; Norieda, H.; Yamada, H.; Nishimura, Y.; Yamazaki, I.; Sakata, Y.; Fukuzumi, S. *J. Am. Chem. Soc.* **2001**, *123*, 100. (j) Imahori, H.; Tkachenko, N. V.; Vehmanen, V.; Tamaki, K.; Lemmetyinen, H.; Sakata, T.; Fukuzumi, S. *J. Phys. Chem. A* **2001**, *105*, 1750. (k) Fukuzumi, S.; Imahori, H.; Yamada, H.; El-Khouly, M. E.; Fujitsuka, M.; Ito, O.; Guldi, D. M. *J. Am. Chem. Soc.* **2001**, *123*, 2571. (l) Imahori, H.; Tamaki, K.; Guldi, D. M.; Luo, C.; Fujitsuka, M.; Ito, O.; Sakata, Y.; Fukuzumi, S. *J. Am. Chem. Soc.* **2001**, *123*, 2607. (m) Imahori, H.; Guldi, D. M.; Tamaki, K.; Yoshida, Y.; Luo, C.; Sakata, Y.; Fukuzumi, S. *J. Am. Chem. Soc.* **2001**, *123*, 6617.
- (11) (a) D'Souza, F.; Deviprasad, G. R.; Rahman, M. S.; Choi, J.-P. *Inorg. Chem.* **1999**, *38*, 2157. (b) Armaroli, N.; Diederich, F.; Echegoyen, L.; Habicher, T.; Flamigni, L.; Marconi, G.; Nierengarten, J.-F. *New J. Chem.* **1999**, 77. (c) Da Ros, T.; Prato, M.; Guldi, D. M.; Alessio, E.; Ruzzi, M.; Pasimeni, L. *Chem. Commun.* **1999**, 635. (d) D'Souza, F.; Rath, N. P.; Deviprasad, G. R.; Zandler, M. E. *Chem. Commun.* **2001**, 267. (e) Da Ros, T.; Prato, M.; Guldi, D. M.; Ruzzi, M.; Pasimeni, L. *Chem. Eur. J.* **2001**, *7*, 816. (f) D'Souza, F.; Deviprasad, G. R.; El-Khouly, M. E.; Fujitsuka, M.; Ito, O. *J. Am. Chem. Soc.* **2001**, *123*, 5277. (g) D'Souza, F.; Deviprasad, G. R.; Zandler, M. E.; Hoang, V. T.; Arkady, K.; VanStipdonk, M.; Perera, A.; El-Khouly, M. E.; Fujitsuka, M.; Ito, O. *J. Phys. Chem. A* **2002**, *106*, 3243. (h) D'Souza, F.; Deviprasad, G. R.; Zandler, M. E.; El-Khouly, M. E.; Fujitsuka, M.; Ito, O. *J. Phys. Chem. B* **2002**, *106*, 4952.
- (12) Deisenhofer, J.; Michel, H. In *Photochemical Energy Conversion*; Norris, J. R., Jr., Meisel, D., Eds.; Elsevier: New York, 1989.
- (13) Maggini, M.; Scorrano, G.; Prato, M. *J. Am. Chem. Soc.* **1993**, *115*, 9798.
- (14) Smith, K. M. *Porphyrins and Metalloporphyrins*; Elsevier: New York, 1977.
- (15) (a) D'Souza, F.; Hsieh, Y.-Y.; Deviprasad, G. R. *Inorg. Chem.* **1996**, *35*, 5747. (b) Nappa, M.; Valentine, J. S. *J. Am. Chem. Soc.* **1978**, *100*, 5075.
- (16) Scatchard, G. *Ann. N. Y. Acad. Sci.* **1949**, *51*, 661.
- (17) D'Souza, F.; Zandler, M. E.; Deviprasad, G. R.; Kutner, W. *J. Phys. Chem. A* **2000**, *104*, 6887. (b) D'Souza, F.; Zandler, M. E.; Smith, P. M.; Deviprasad, G. R.; Arkady, K.; Fujitsuka, M.; Ito, O. *J. Phys. Chem. A* **2002**, *106*, 649. (c) Zandler, M. E.; Smith, P. M.; Fujitsuka, M.; Ito, O.; D'Souza, F. *J. Org. Chem.* **2002**, in press.
- (18) Nojiri, T.; Watanabe, A.; Ito, O. *J. Phys. Chem. A* **1998**, *102*, 5215.
- (19) (a) Ghosh, H. N.; Pal, H.; Sapre, A. V.; Mittal, J. P. *J. Am. Chem. Soc.* **1993**, *115*, 11722. (b) Fujitsuka, M.; Ito, O.; Yamashiro, T.; Aso, Y.; Otsubo, T. *J. Phys. Chem. A* **2000**, *104*, 4876.
- (20) (a) Matsumoto, K.; Fujitsuka, M.; Sato, T.; Onodera, S.; Ito, O. *J. Phys. Chem. B* **2000**, *104*, 11632. (b) Komamine, S.; Fujitsuka, M.; Ito, O.; Morikawa, K.; Miyata, T.; Ohno, T. *J. Phys. Chem. A* **2000**, *104*, 11497. (c) Yamazaki, M.; Araki, Y.; Fujitsuka, M.; Ito, O. *J. Phys. Chem. A* **2001**, *105*, 8615.
- (21) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Zakrzewski, V. G.; Montgomery, J. A., Jr.; Stratmann, R. E.; Burant, J. C.; Dapprich, S.; Millam, J. M.; Daniels, A. D.; Kudin, K. N.; Strain, M. C.; Farkas, O.; Tomasi, J.; Barone, V.; Cossi, M.; Cammi, R.; Mennucci, B.; Pomelli, C.; Adamo, C.; Clifford, S.; Ochterski, J.; Petersson, G. A.; Ayala, P. Y.; Cui, Q.; Morokuma, K.; Malick, D. K.; Rabuck, A. D.; Raghavachari, K.; Foresman, J. B.; Cioslowski, J.; Ortiz, J. V.; Stefanov, B. B.; Liu, G.; Liashenko, A.; Piskorz, P.; Komaromi, I.; Gomperts, R.; Martin, R. L.; Fox, D. J.; Keith, T.; Al-Laham, M. A.; Peng, C. Y.; Nanayakkara, A.; Gonzalez, C.; Challacombe, M.; Gill, P. M. W.; Johnson, B. G.; Chen, W.; Wong, M. W.; Andres, J. L.; Head-Gordon, M.; Replogle, E. S.; Pople, J. A. *Gaussian 98*, revision A.7; Gaussian, Inc.: Pittsburgh, PA, 1998.