

Article

A Chlorin-based Nanoscale Metal-Organic Framework Systemically Rejects Colorectal Cancers via Synergistic Photodynamic Therapy and Checkpoint Blockade Immunotherapy

Kuangda Lu, Chunbai He, Nining Guo, Christina Chan, Kaiyuan Ni, Ralph R Weichselbaum, and Wenbin Lin

J. Am. Chem. Soc., Just Accepted Manuscript • DOI: 10.1021/jacs.6b06663 • Publication Date (Web): 30 Aug 2016 Downloaded from http://pubs.acs.org on August 30, 2016

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



Journal of the American Chemical Society is published by the American Chemical Society. 1155 Sixteenth Street N.W., Washington, DC 20036 Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

A Chlorin-based Nanoscale Metal-Organic Framework Systemically Rejects Colorectal Cancers via Synergistic Photodynamic Therapy and Checkpoint Blockade Immunotherapy

Kuangda Lu,^{a,†} Chunbai He,^{a,†} Nining Guo,^{a,b,†} Christina Chan,^a Kaiyuan Ni,^a Ralph R. Weichselbaum,^b Wenbin Lin^{a,*}

^aDepartment of Chemistry, The University of Chicago, Chicago, IL 60637, USA and ^bDepartment of Radiation and Cellular Oncology and The Ludwig Center for Metastasis Research, The University of Chicago, Chicago, IL 60637, USA

E-mail: wenbinlin@uchicago.edu

KEYWORDS (Word Style "BG_Keywords"). If you are submitting your paper to a journal that requires keywords, provide significant keywords to aid the reader in literature retrieval.

ABSTRACT: Photodynamic therapy can destroy local tumors and minimize normal tissue damage, but is ineffective at eliminating metastases. Checkpoint blockade immunotherapy has enjoyed recent success in the clinic, but only elicits limited rates of systemic antitumor response for most cancers due to insufficient activation of the host immune system. Here we describe a treatment strategy that combines photodynamic therapy (PDT) by a new chlorin-based nanoscale metal-organic framework (nMOF), TBC-Hf, and a small molecule immunotherapy agent, that inhibits Indoleamine 2,3-dioxygenase (IDO), encapsulated in the nMOF channels to induce systemic antitumor immunity. The synergistic combination therapy achieved effective local and distant tumor rejection in colorectal cancer models. We detected increased T cell infiltration in the tumor microenvironment after activation of the immuno system with the combination of the IDO inhibition by the small molecule immunotherapy agent and the immunogenic cell death induced by PDT. We also elucidated the underlying immunological mechanisms and revealed compensatory roles of neutrophils and B cells in presenting tumor-associated antigens to T cells in this combination therapy. We believe that nMOF-enabled PDT has the potential to significantly enhance checkpoint blockade cancer immunotherapy, affording clinical benefits for the treatment of many difficult-to-treat cancers.

INTRODUCTION

Photodynamic therapy (PDT) combines non-toxic photosensitizers (PSs), light, and tissue oxygen to generate reactive oxygen species (ROS) such as O_2 to achieve local tumor control/rejection.¹ PDT is highly effective for the treatment of esophageal cancer and skin lesions but cannot completely eradicate deep-seated tumors due to shallow penetration of light through tissues.² The localized light irradiation also renders PDT ineffective for treating systemically disseminated disease. Therefore, more efficient PSs and new treatment strategies are needed to enhance the efficacy of PDT in both eliminating local tumors and controlling distant metastases.

Significant efforts have been devoted to developing nanoparticle-based PSs to enhance PDT in the past decade.³ We first reported the use of porphyrin- and chlorin-based nanoscale metal-organic frameworks (nMOFs) as efficient PSs for PDT of cancer. The stable framework and crystalline structure of nMOFs allow for high PS loadings whereas the heavy metal cluster secondary building units (SBUs) facilitate intersystem crossing to enhance ¹O₂ generation.⁴ We further demonstrated in a recent study that both apoptosis and immunogenic cell death (ICD) contribute to the anticancer efficacy of nMOF-enabled PDT.^{4b} The acute inflammatory response of PDT can alter the immunosuppressive microenvironment of the local tumor to prime the host immune system.⁵ We hypothesized that the porous nMOF structure could be leveraged to accentuate immune response by delivering small molecule immunostimulatory agents in the open channels, combining PDT and immunotherapy to eradicate both local, treated tumors and reject distant, untreated tumors.

Immunotherapy has recently emerged as a highly effective cancer treatment strategy, with high overall response rates in multiple cancer types and durable tumor control enjoyed by a small subset of patients.⁶ Of particular interest is checkpoint blockade immunotherapy which uses small molecules or antibodies to stimulate the immunosuppressive microenvironment of tumors by modulating protein expressions and/or functions at dysregulated immune checkpoints.^{6b,7} Indoleamine 2,3-dioxygenase (IDO) is one such checkpoint, an immunoregulatory enzyme highly expressed in tumors that catalyzes the oxidative catabolism of tryptophan (Trp) to kynurenine (Kyn).⁸ The results of this catalytic process lead to depletion of Trp and production of Kyn, thus preventing the clonal expansion of T-cells and promoting T-cell anergy and apoptosis.⁹ Small molecule IDO inhibitors such as 1-methyl-tryptophan,^{9b, 10} INCB24360,¹¹ and NLG919,¹² can effectively block Trp catabolism, but show modest effect as monotherapies, due in part to ineffective antigen presentation and insufficient antitumor immunity.

We herein report the design of a new nMOF, TBC-Hf, constructed from a chlorin derivative, 5,10,15,20-tetra(p-benzoato)chlorin (H₄TBC) and Hf₆(μ_3 -O)₄(μ_3 -OH)₄ SBUs. We took advantage of the highly porous structure of the nMOF to load an IDO inhibitor (IDOi) into TBC-Hf to afford IDOi@TBC-Hf. The IDOi@TBC-Hf system enables a novel cancer treatment strategy by combining nMOF-enabled PDT and IDOi-based immunotherapy (Figure 1). We demonstrated the eradication of the primary, treated tumors and the regression of the distant, untreated tumors upon IDOi@TBC-Hf administration and light irradiation using two syngeneic colorectal cancer models, CT26 and MC38, and investigated the underlying immunological mechanisms.



Figure 1. Schematic presentation of combined PDT and immunotherapy by IDOi@TBC-Hf. Local injection of IDOi@TBC-Hf and light irradiation generate reactive oxygen species, causing immunogenic cell death (ICD), and releasing tumor-associated antigens which are presented to T-cells. Meanwhile, the IDO inhibitor released from IDOi@TBC-Hf modulates tryptophan/kynurenine catabolism to activate the immunosuppressive tumor microenvironment. The combination of antigen presentation from PDT and checkpoint blockade by IDO inhibition causes T-cell proliferation and infiltration, leading to not only eradication of local, treated tumors but also rejection of distant, untreated tumors.

RESULTS

Synthesis and Characterization of nMOFs

The new chlorin derivative 5,10,15,20-tetra(p-benzoato) chlorin (H_4 TBC) was synthesized as depicted in scheme 1. Briefly, 5,10,15,20-tetra(p-methylbenzoato)chlorin (Me_4 TBC) was synthesized by a partial reduction of 5,10,15,20-tetra(p-methylbenzoato)porphyrin (Me_4 TBP) with toluenesulfonhydrazide in 16% yield. Saponification of Me_4 TBC afforded H_4 TBC in 78% yield. Me_4 TBC and H_4 TBC were characterized by NMR and mass spectrometry (Figures S1-S5, supporting information [SI]).

Scheme 1. Synthesis of 5,10,15,20-tetra(p-benzoato) chlorin (H4TBC).



A solvothermal reaction between $HfCl_4$ and H_4TBC in N,N-dimethylformamide (DMF) at 80 °C afforded the purple powder of TBC-Hf in 87% yield. TBC-Hf was washed with copious amounts of DMF, 1% triethylamine (NEt₃) in ethanol (v/v), and ethanol successively and stored in ethanol as a stock suspension. As a control, TBP-Hf with the corresponding porphyrin ligand 5,10,15,20-tetra(p-benzoato)porphyrin (H₄TBP) was synthesized by a solvothermal reaction between HfCl₄ and H₄TBP in N,N-diethylformamide (DEF) at 120 °C and washed in the same fashion as for TBC-Hf.

TBC-Hf and TBP-Hf adopt the same structure as the previously reported TBP-Zr analog (MOF-545), based on powder X-ray diffraction (PXRD) (Figure 2a).¹³ The carboxylate moieties from the TBC ligands bridge 8 out of 12 edges of octahedral $Hf_6(\mu_3-O)_4(\mu_3-OH)_4$ SBUs. The remaining coordination sites on the SBUs are occupied by 4 water and 4 terminal hydroxy groups on the equatorial plane to afford a hexagonal framework of $Hf_6(\mu_3-O)_4($ $OH_4(OH_4(H_2O)_4(TBC)_2$ (Figure S6, SI). The nMOFs therefore feature large one-dimensional channels with diameters of 3.5 nm along the c axis, along with smaller channels with diameters of 1.0 nm along the c axis and 1.1 $nm \times 0.9$ nm windows perpendicular to the c axis. TBC-Hf and TBP-Hf exhibit BET surface area of 1077 and 1462 m²/g, respectively (Figure S₇, SI). During the course of this study, a few reports of TBP-based nMOFs with a cubic structure for PDT have appeared.¹⁴ However, we expect that our highly porous TBP-Hf control will outperform the cubic TBP-based nMOFs due to the enhanced ¹O₂ diffusion through the nMOF channels. In TBC-Hf and TBP-Hf, the porphyrin and chlorin rings are highly rigid in the nMOF structures to minimize energy transfer and self-quenching.



Figure 2. Morphology and structure of TBP-Hf and TBC-Hf. (a) PXRD patterns of TBP-Hf, TBC-Hf, and IDOi@TBC-Hf in comparison to MOF-545. TEM images of TBP-Hf (b-c), TBC-Hf (d-e) and TBC-Hf after incubation in Hank's balanced salt solution for 24 h (f).

Transmission electron microscopy (TEM) revealed nanorod/nanorice morphologies for TBP-Hf and TBC-Hf (Figure 2b,d). The rods are typically 50~100 nm long and 30-60 nm wide for TBC-Hf, and 50~100 nm long and 20-30 nm wide for TBP-Hf. High resolution images show the lattice fringes with the distances matching d_{200} and d_{001} spacings of the crystal structure (Figure 2c,e and S8-S9, SI). The average diameter of TBC-Hf and TBP-Hf are 83.2 nm and 72.7 nm, respectively, by dynamic light scattering (DLS) measurements (Figure S10, SI). The nMOFs are stable in biological environments, as evidenced by the lack of significant changes in PXRD pattern or morphology after incubation in Hank's balanced salt solution (HBSS) for 24 h (Figure 2a,f).

The Hf content was determined by inductively coupled plasma-mass spectrometry (ICP-MS) to be 30.4% (36.6% calculated). The TBC weight loss of 61.7% (56.8% calculated) determined by thermogravimetric analysis further confirms the framework formula (Figure S11a, SI). We encapsulated an INCB24360 analogue,¹¹ 4-amino-N-(3-chloro-4-fluoro-phenyl)-N'-hydroxy-1,2,5-oxadiazole-3-carboximidamide (Figure S11b, SI), into the TBC-Hf channels to afford IDOi@TBC-Hf for the co-delivery of IDOi with PSs. The IDOi weight percentage after loading is determined to be 4.7% by TGA (Figure S11a and S12, SI). The release of IDOi from IDOi@TBC-Hf was studied in HBSS (Figure S13, SI). IDOi slowly leached from TBC-Hf, reaching 83.3% release after incubation in HBSS for 24 h.

UV-visible absorption spectroscopy indicates that TBC-Hf more effectively absorbs red light compared to TBP-Hf (Figure 3a). H₄TBC has a major Soret peak at λ_{max} =420 nm and four Q-bands at 518, 546, 600, and 652 nm. The Soret absorption of TBC-Hf almost overlaps with that of the free ligand (λ_{max} =421 nm) while the Q-bands slightly redshift to 520, 548, 600, and 653 nm. In comparison, the Soret peaks of H₄TBP and TBP-Hf are at λ_{max} =420 nm and 418 nm, respectively, and the Q-bands are at 516, 549, 592 and 646 nm for H_4 TBP and 517, 550, 593 and 647 nm for TBP-Hf. The lowest energy Q-band of the chlorin therefore red-shifts by 6 nm after reduction from the porphyrin counterpart (652 nm compared to 646 nm). The extinction coefficient of H₄TBC at the lowest energy absorption $(\varepsilon = 44700 \text{ M}^{-1} \cdot \text{cm}^{-1})$ is ~9 fold greater than that of H₄TBP $(\epsilon = 4800 \text{ M}^{-1} \cdot \text{cm}^{-1})$ while the lowest energy absorption of TBC-Hf (ϵ =38500 M⁻¹·cm⁻¹) is 6-fold greater than that of TBP-Hf (ϵ =6400 M⁻¹·cm⁻¹). Notably, the lowest energy absorption of TBC-Hf (653 nm) is at a wavelength of 7-nm longer than that of our previously reported DBC-UiO and the ε of TBC-Hf is ~1.6 fold higher than DBC-UiO.

H₄TBC exhibits two fluorescence emissions at 655 nm and 715 nm with an intensity ratio of ~6.8 when excited at 420 nm (Figure 3c). H₄TBP fluorescence emissions are close to those of H_4 TBC, though the 651 nm/713 nm peak intensity ratio is ~2 (Figure 3b). After forming nMOFs, the fluorescence intensities drop significantly, consistent with other nMOF systems.⁴ For TBC-Hf, the fluorescence peaks slightly blue-shift to 650 nm and 712 nm and the intensities decrease by 5.3-fold and 4.7-fold, respectively; for TBP-Hf, the fluorescence peaks blue-shift to 648 nm and 704 nm and the fluorescence intensities drop by 9.0fold and 6.5-fold, respectively. The decrease in fluorescence intensity likely results from the enhanced intersystem crossing after coordination of the chlorin/porphyrin ligands to Hf on the SBUs. Time-Correlated Single Photon Counting measurement results also supported this interpretation: the fluorescence lifetime at 650 nm dropped from 9.56 ns for H₄TBC to 8.25 ns for TBC-Hf and from 9.25 ns for H₄TBP to 8.38 ns for TBP-Hf (Figure 3d-e and Table S1, SI).

Singlet Oxygen Generation

The ${}^{1}O_{2}$ generation efficiencies of H₄TBC, TBC-Hf, H₄TBP, and TBP-Hf were determined by Singlet Oxygen Sensor Green (SOSG) assay. The nMOF samples and ligand controls were irradiated with a LED (λ_{max} =650 nm, 20 mW/cm²) and the fluorescence enhancement upon reaction of SOSG with ${}^{1}O_{2}$ was measured by a fluorimeter (excitation/emission at 504/525 nm). The fluorescence intensity plotted against irradiation time fit exponential decay equations, as we had previously observed in other nMOF systems,⁴ indicating a pseudo first-order ${}^{1}O_{2}$ generation process (Figure 3f). TBC-Hf outperforms both TBP-Hf and H₄TBC in terms of ${}^{1}O_{2}$ generation efficiency.

TBC-Hf

Н₄ТВС

b) 6

Figure 3. Photophysics and photochemistry of the ligands and nMOFs. (a) UV-visible absorption spectra of H_4 TBP, H_4 TBC, TBP-Hf, and TBC-Hf. Inset shows expanded views of the Q-band regions. The fluorescence spectra of H_4 TBP and TBP-Hf (b), H_4 TBC and TBC-Hf (c) with excitation at 420 nm. Time-resolved fluorescence decay traces of H_4 TBP and TBP-Hf (d), H_4 TBC and TBC-Hf (e) along with instrument response function (IRF). (f) Singlet oxygen generation by H_4 TBP, H_4 TBC, TBP-Hf, and TBC-Hf with a 650 nm LED irradiation at 20 mW/cm², detected by Singlet Oxygen Sensor Green.

In vitro PDT efficacy and mechanistic study

We first confirmed efficient cellular uptake of TBC-Hf and TBP-Hf by cancer cells. TBC-Hf and TBP-Hf were incubated with colorectal cancer CT26 cells at a ligand concentration of 25 μ M for 24 h. The Hf contents inside cells were determined to be (2.07±0.26) and (1.79±0.17) nmol/10⁴ cells, respectively, by ICP-MS analysis (Figure S14, SI). The *in vitro* PDT efficacy of TBP-Hf, TBC-Hf, and

both ligands against CT₂₆ cells was next investigated. TBC-Hf outperformed TBP-Hf under 90 J/cm² light irradiation. Both nMOFs and ligands outperformed protoporphyrin IX (PpIX) upon irradiation, a commonly used PS, while no cytotoxicity was observed in dark control groups (Figure S15, SI). The IC₅₀ values of TBC-Hf, TBP-Hf, H₄TBC, H₄TBP and PpIX for CT26 cells were 4.82±0.50, 7.09±0.39, 10.04±1.22, 12.83±1.59 and 23.72±0.60 μ M, respectively, at an irradiation dose of 90 J/cm² (Figure 4a). These results were confirmed by a second colorectal cancer MC38 cell with IC50 values of 2.57±0.85 and 17.9±5.0 µM for TBC-Hf and H₄TBC, respectively. The IC₅₀ values exceeded 50 µM for TBP-Hf, H₄TBP and PpIX for MC₃8 cells (Figure 4b). To demonstrate the versatility of our nMOF system, unrelated B16F10 melanoma cells were employed to confirm the PDT efficacy, yielding IC_{50} values of 5.48±0.70, 9.72±0.78, 11.03±1.57, 20.10±4.28 and 16.48±0.77 µM for TBC-Hf, TBP-Hf, H₄TBC, H₄TBP and PpIX, respectively (Figure S16, SI).

Unlike most conventional cytotoxic agents, PDT is known to induce ICD via apoptosis and necrosis, thereby mediating antitumor immune response.⁵ We confirmed that both apoptosis and necrosis occur upon treatment with PDT by flow cytometry of Alexa Fluor 488 Annexin V-labeled dead cells. MC38 cells were incubated with TBC-Hf, TBP-Hf, H₄TBC and H₄TBP at 1 μ M (ligand concentration of 2 µM) followed by light irradiation at 90 J/cm^2 (650 nm, 100 mW/cm²). Significant amounts of cells underwent apoptosis/necrosis when treated with PDT of TBC-Hf, TBP-Hf, H₄TBC and H₄TBP with only 30.0%, 32.7%, 55.5%, and 57.6% healthy cells, respectively, compared to 80-90% healthy cells in dark control or PBStreated cells with irradiation (Figure 4c and Figure S17a, SI). These results were further confirmed in CT₂₆ cells under the same conditions (Figure S17b, SI). Taken together, the chlorin-based TBC-Hf is a more efficient PS than the porphyrin-based TBP-Hf at equivalent nMOF and light doses.

We also investigated the ability of PDT treatment to induce ICD by determining cell-surface expression of calreticulin (CRT). MC38 cells were incubated with TBC-Hf, TBP-Hf, H_4 TBC and H_4 TBP at 1 μ M (ligand concentration of 2 μ M) followed by light irradiation at 90 J/cm². For flow cytometry analysis, cells were collected and stained with Alexa Fluor 488-CRT antibody and propidium iodide (PI, Figure 4d and Figure S18, SI). The fluorescence intensity of stained cells was gated on PI-negative cells. For immunostaining analysis, the cells were stained with AlexaFluor 488-CRT and DAPI, and observed under confocal laser scanning microscopy (CLSM, Figure 4e and Figure S19, SI). In both instances, cells treated with nMOF or ligand showed significant CRT expression on the cellsurface upon irradiation but none with PBS control or without irradiation. Due to the higher 'O2 generation efficiency and in vitro PDT efficacy of TBC-Hf, we focus our subsequent in vivo studies on TBC-Hf and use TBP-Hf as a control wherever appropriate.

1

2

3

4

5

6

a)

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59 60

Figure 4. In vitro PDT efficacy and mechanistic study. PDT cytotoxicity of H₄TBP, TBP-Hf, H₄TBC, TBC-Hf, and PpIX at different PS concentrations in CT26 (a) and MC38 cells (b). (c) Annexin V/PI analysis of MC38 cells incubated with TBC-Hf or PBS with or without irradiation (90 J/cm²). The quadrants from lower left to upper left (counter clockwise) represent healthy, early apoptotic, late apoptotic, and necrotic cells, respectively. The percentage of cells in each quadrant is shown on each graph. (+) and (-) refer to with and without irradiation, respectively. (d) CRT exposure on the cell surface of MC38 was assessed after incubation with PBS and TBC-Hf with or without light irradiation (90 J/cm²) by flow cytometry analysis. (e) The fluorescence intensity was gated on PI-negative cells. Immunofluorescence microscopy of CRT expression on the cell surface of MC38 treated with PBS and TBC-Hf upon irradiation (90 J/cm²).

The Abscopal Effect of IDOi@TBC-Hf

The immunoregulatory enzyme IDO is often overexpressed in the tumor microenvironment, where it facilitates the survival and growth of tumor cells by suppressing antitumor immune responses.15 The small-molecule IDO inhibitor (IDOi) INCB24360, developed by Incyte, can reverse immunosuppression and inhibits tumor growth upon oral administration.¹⁶ We propose that by encapsulation of IDOi into the nMOF channels, IDOi@TBC-Hf can release IDOi both into the local tumor environment and enter blood circulation for systemic IDO inhibition. The effects of IDOi will alter the suppressive tumor microenvironment of both the treated and untreated tumors. We hypothesize that synergizing local PDT of IDOi@TBC-Hf with checkpoint blockade therapy can promote an efficient abscopal effect, regression of an untreated tumor at a distant site following the local treatment of another tumor in the same organism.

We evaluated the abscopal effect of IDOi@TBC-Hf and light irradiation in two immunocompetent mouse models

using bilateral tumor models of colorectal cancers CT₂₆ and MC38 in the flank regions of BALB/c mice and C57BL/6 mice, respectively. When the primary tumors reached ~100 mm³, mice received a single injection of H₄TBC, H₄TBC plus IDOi, TBC-Hf, or IDOi@TBC-Hf into the primary tumors at a ligand dose of 20 µmol/kg or a corresponding IDOi dose of 1.5 mg/kg. Twelve hours after injection, the primary tumors were irradiated at a light dose of 90 J/cm² (650 nm, 100 mW/cm²). Mice treated with IDOi@TBC-Hf without light irradiation served as dark controls. The primary tumor receiving intratumoral injection with or without irradiation was designated as the "treated tumor", while the contralateral tumor which received neither direct injection nor irradiation was designated as the "untreated tumor". As depicted in Figure 5 and Figure S20-S22 (SI), local nMOF injection with light irradiation led to near elimination of the treated primary tumors. At the endpoint, CT26 tumor-bearing mice treated with IDOi@TBC-Hf or TBC-Hf and PDT therapy had tumors only (1.1±0.2)% and (0.7±0.4)% the size of PBS treated tumors, respectively. Similarly, MC38 tumorbearing mice had tumors only (0.8±0.3)% and (0.9±0.4)% of the size of PBS treated tumors, respectively. H₄TBC with light irradiation and IDOi@TBC-Hf dark group failed to inhibit the tumor growth while H₄TBC plus IDOi with light irradiation slightly inhibits the tumor growth (Table S2). Moreover, only PDT treatment of IDOi@TBC-Hf successfully reduced the sizes of the untreated distant tumors. Tumors began shrinking on Day 6 and 5 after treatment in the CT₂₆ and MC₃₈ models, respectively, suggesting the treatment evoked systemic antitumor immunity in mice (Table S2-S3). TBC-Hf with light irradiation and IDOi@TBC-Hf dark control slightly inhibited the distant tumor growth, showing ineffectiveness of monotherapies.

Figure 5. *In vivo* anticancer efficacy showing abscopal effect. Growth curves for treated (a and c) and untreated (b and d) tumors of CT26 (a and b) or MC38 (c and d) tumor-bearing mice after PDT treatment. Black and red arrows refer to the time of injection and irradiation, respectively.

Antitumor Immunity

60

1

Journal of the American Chemical Society

We have shown that PDT of IDOi@TBC-Hf caused effective tumor regression of both primary, treated tumors and distant, untreated tumors in two syngeneic mouse models of colorectal cancer. We further investigated the underlying immunological mechanisms in the MC₃8 model by Enzyme-Linked ImmunoSpot (ELISPOT) and flow cytometry. We first performed an ELISPOT assay to detect the presence of tumor antigen-specific T cells 14 days after treatment. As shown in Figure 6a, the number of antigen-specific IFN- γ producing T cells increased to 799 per million splenocytes in mice treated with IDOi@TBC-Hf (vs. 128 per million splenocytes in PBS group, P=0.0879), indicating that PDT of IDOi@TBC-Hf induced *in situ* tumor vaccination to effectively generate tumor-specific T cell response.

Further analysis of the immune response was carried out by investigating the tumor-infiltrating leukocyte profiles in each tumor by flow cytometry. Previous research has shown that PDT can evoke acute inflammation,⁵ so we first evaluated the population of leukocytes responsible for innate immune response (neutrophils, B cells, dendritic cells, and macrophages) in both primary and distant tumors 12 hours after PDT treatment. As shown in Figure 6 b-c and Figure S23 (SI), IDOi@TBC-Hf administration and light irradiation led to significant increase in the percentage of tumor-infiltrating neutrophils and B cells with respect to the total number of cells in the tumor compared to PBS (neutrophils: P=0.0369 vs. PBS; B cells: P=0.0215 vs. PBS). To our surprise, the percentage of infiltrating dendritic cells appeared to decrease compared to PBS control in the primary tumor (Figure S23, SI), although the difference is not of statistical significance.

To better understand the roles these types of cells played in the antigen presentation after PDT, we further detected the major histocompatibility complex class II (MHC-II) expression on these cells 12 hours after PDT treatment. MHC-II mediates establishment of specific immunity by interacting with CD4 molecules on the surface of helper T cells.¹⁷ We found a significant decrease in MHC-II expression levels on dendritic cells, macrophages, and neutrophils in the primary treated tumors (DC: P<0.0001 vs. PBS; macrophages: P=0.0004 vs. PBS; neutrophils: P=0.006 vs. PBS, Figure S24, SI). These results imply that TBC-Hf mediated PDT may have an impact on the method of antigen presentation, which is typically the role of dendritic cells.

We profiled the tumor-infiltrating leukocytes on 7 days post PDT treatment. The mice treated with IDOi@TBC-Hf plus PDT did not show significant difference in all cell types except B cells from the control mice (Figure S25, SI). The percentage of B cells significantly increased with respect to the total number of cells in the primary tumors of IDOi@TBC-Hf and PDT treated mice (P=0.0018, Figure S25 c, SI), while the percentage of dendritic cells significantly decreased in the distant tumors of treated mice (P=0.0083, Figure S25 f, SI). We also found that the percentage of NK cells was extremely low on day 7 post PDT treatment both in IDOi@TBC-Hf treated mice and PBS treated mice (both percentage <0.05%, Figure S25 e, SI).

We further determined the tumor-infiltrating leukocyte profiles 12 days post PDT treatment. As shown in Figure 6 d-f, IDOi@TBC-Hf significantly increased the proportion of infiltrating CD8⁺ T cells relative to the total number of cells in the distant tumor (P=0.0012 vs. PBS), an essential step to mounting an antitumor immune response to induce the abscopal effect. In both primary and distant tumors, the percentages of infiltrating CD₄₅⁺ leukocytes and CD4⁺ T cells with respect to the total number of cells in the tumors were significantly increased in mice treated with PDT of IDOi@TBC-Hf (Treated tumor, CD45+: P=0.0061 vs. PBS; CD4+ T cells: P=0.0206 vs. PBS. Untreated tumor, CD45+: P=0.0001 vs. PBS; CD4+ T cells: P=0.0388 vs. PBS). Furthermore, the percentage of tumorinfiltrating NK cells significantly increased in the distant tumors of mice treated with IDOi@TBP-Hf and PDT (P=0.0034 vs. PBS, Figure S26, SI). Interestingly, IDOi@TBP-Hf administration and PDT also led to a significant increase in the percentages of tumor-infiltrating B cells and a decrease in the percentages of tumor infiltrating dendritic cells with respect to the total number of cells in the primary tumors 12 days post the treatment (B cells, P=0.0017 vs. PBS; DCs: P=0.0041 vs. PBS, Figure S27, SI).

We also performed T cell blocking experiments to confirm the involvement of T cells in the efficient abscopal response on bilateral subcutaneous MC38 mouse model. MC38 tumor-bearing mice were treated with IDOi@TBC-Hf and light irradiation as described before and received i.p. injection of Mouse IgG, anti-CD4, or anti-CD8 antibody at a dose of 200 µg/mouse/injection on Day 0 and Day 5 post the PDT treatment. While mice treated with Mouse IgG showed regression of both primary and distant tumors, no abscopal effect was observed on mice with blocking of CD4⁺ or CD8⁺ T cells (Figure 6 g-h). Furthermore, blocking of CD4⁺ and CD8⁺ T cells also diminished the effect of IDOi@TBC-Hf in regressing the primary tumors. These results indicated that CD4⁺ and CD8⁺ T cells were essential not only to abscopal effect but also to primary tumor rejection.

1

Figure 6. Antitumor immunity of PDT by IDOi@TBC-Hf. (a) Antigen-specific IFN-γ spot forming cells (SFC) 14 days after IDOi@TBC-Hf PDT treatment. The percentage of tumor-infiltrating neutrophils (b) and B cells (c) with respect to the total number of cells in the tumor compared to PBS, 12h after IDOi@TBC-Hf PDT treatment. The percentage of tumor-infiltrating CD8⁺ T cells (d), CD_4^+ T cells (e) and CD_{45}^+ cells (f) with respect to the total number of cells in the tumors 12 days after IDOi@TBC-Hf PDT treatment. Growth curves for primary, treated (g) and distant, untreated (h) tumors after PDT treatment (+) with IDOi@TBC-Hf and injections of Mouse IgG, anti-CD4, or anti-CD8 antibody, compared to PBS control without PDT treatment (-). Black, red and blue arrows refer to the time of IDOi@TBC-Hf injection, light irradiation and antibody injection, respectively.

Discussion

Approximately 140,000 patients are diagnosed with colorectal cancer in the US annually, with one-third dying from metastasis.¹⁸ Stimulation of the host immune system has been shown to generate an antitumor immunity capable of controlling metastatic tumor growth, and thus represents a promising treatment strategy for metastatic colon cancer.⁸ Clinical results suggest that immunotherapies have the potential to achieve systemic and adaptable cancer control. However, combination with other treatment modalities is necessary to maximize the benefits of immunotherapy and increase the tumor-specific immune response rate.^{6b} Here we report a new treatment strategy that combines local PDT of a new nMOF, TBC-Hf, in combination with IDO inhibition with nMOF-delivered small molecules to achieve effective and consistent abscopal responses in mouse models of colorectal cancers. We have shown that IDOi released from locally injected IDOi@TBC-Hf reversed the suppressive tumor microenvironment in both treated and untreated tumors, which further synergized with TBC-Hf mediated PDT to stimulate the immune system for activating both acute innate and prolonged adaptive immune response to achieve efficient local tumor regression and a consistent abscopal response. Our treatment method maximizes the benefits of local treatment with systemic immune response for the rejection of both primary and distant tumors while minimizing side effects, potentially affording an effective systemic therapy for metastatic colorectal cancers.

Owing to the outstanding photophysical properties of the chlorin-based TBC ligand, TBC-Hf shows potent PDT efficacy and outperforms its porphyrin counterpart TBP-Hf. The tetracarboxylate ligand constructs a robust framework with very large channels for small molecular inhibitor loading and ROS diffusion, and the coordination to Hf₆ SBUs enhances intersystem crossing of TBC to enhance ¹O₂ generation. As we¹⁹ and others²⁰ have demonstrated the synthetic tunability and potential biomedical applications of nMOFs in the past decade, we believe that the nMOF compositions and structures can be further optimized in order to enhance light absorption and 'O2 generation/diffusion. We thus believe that nMOF-based synergistic PDT and immunotherapy have the potential for future clinical translation for the treatment of metastatic cancers.

We propose that IDOi@TBC-Hf administration and light irradiation causes highly efficient tumor regression of both primary treated tumors and distant untreated tumors owing to two factors. First, TBC-Hf based PDT causes ICD of cancer cells in the primary tumors, which activates innate immune system and promotes antigen presentation (Figure 1). The massive stressed and dying necrotic tumor cells in the PDT-treated primary tumor sites are engulfed by the innate immune effector cells followed by presenting tumor-derived antigenic peptides to T cells, thus stimulating a tumor-specific T cell response. Second, the IDOi is released from intratumorally injected IDOi@TBC-Hf to systemically inhibit IDO activity to reverse the immunosuppressive tumor environments. Alternatively, the recruited immune cells likely undergo IDO inhibition in the treated tumors and migrate to the distant tumors to cause an abscopal effect. The two treatment modalities, PDT and IDOi checkpoint blockade therapy, synergize with each other to kill cancer cells locally and create an immunogenic tumor microenvironment systemically, leading to strong and consistent abscopal effects.

Dendritic cells are one of the most important antigenpresenting cells and prevalently believed to play a key role in antitumor immune response.²¹ However, we observed a decrease of dendritic cell population percentage at differ-

ent time points along with a significant decrease of MHC-II expression level on dendritic cells after PDT treatment with IDOi@TBC-Hf, indicating an altered role for dendritic cells after PDT treatment. Meanwhile, the increase of both neutrophils and B cell population percentages implies a compensating effect of other antigen presenting cells to present tumor-associated antigens and initiate the antitumor immune response. Previous studies also suggested that neutrophils, rather than dendritic cells, directly affect T cell proliferation upon PDT treatment.²² We also observed significant increase in the percentage of tumor infiltrating NK cells from 0.02% on day 7 to 3.83% on day 12 post PDT treatment in distant tumors, suggesting the important role of tumor-infiltrating NK cells in tumor rejection in addition to those played by CD8⁺ T cells, CD4⁺ T cells, and B cells after IDOi@TBC-Hf and PDT treatment. Our future efforts will be directed toward elucidating the roles of neutrophils and B cells in initiating antitumor immune response after the PDT of IDOi@TBC-Hf.

Conclusion

In this work, we have rationally designed a chlorinbased nMOF with large channels for highly efficient PDT, while simultaneously loaded an IDO inhibitor into its channels to achieve combination therapies of PDT and checkpoint blockade immunotherapy. We consistently observed an abscopal effect in mice receiving treatment with PDT of IDOi@TBC-Hf. The in situ vaccination induced by PDT treatment and IDOi immunotherapy synergize with each other and effectively generate systemic antitumor immunity. We believe the present strategy has the potential to significantly increase the systemic tumorspecific immune response rates of checkpoint blockade cancer immunotherapy and lead to clinical benefits for the treatment of metastatic colorectal cancers and other difficult-to-treat cancers.

Materials and Methods

Materials, cell lines, and animals

All of the starting materials were purchased from Sigma-Aldrich and Fisher (USA), unless otherwise noted, and used without further purification. INCB24360 analogue, 4-amino-N-(3-chloro-4-fluoro-phenyl)-N'hydroxy-1,2,5-oxadiazole-3-carboximidamide was purchased from Medkoo Biosciences, USA.

Murine colon adenocarcinoma cells CT26 and MC38 and murine melanoma cells Bi6Fio were purchased from the American Type Culture Collection (Rockville, MD, USA) and cultured in Dulbecco's Modified Eagle's Medium (DMEM) medium (Gibco, Grand Island, NY, USA) supplemented with 10% FBS.

C57BL/6 female mice (6 weeks, 20-22 g) and BALB/c female mice (6 weeks, 20-22 g) were provided by Harlan Laboratories, Inc (USA). The study protocol was reviewed and approved by the Institutional Animal Care and Use Committee (IACUC) at the University of Chicago.

Synthesis of TBC-Hf

To a 2-dram glass vial was added 1 mL of $HfCl_4$ solution [2 mg/mL in N,N-dimethylformamide (DMF), 6.2 µmol], 1 mL of the H₄TBC solution (1.9 mg/mL in DMF, 2.4 µmol), and 60 µL of 88% formic acid (1.4 mmol). The reaction mixture was kept in an 80 °C oven for 2 days. The purple powder was collected by centrifugation and washed with DMF, 1% triethylamine in ethanol (v/v) and ethanol.

Singlet Oxygen Generation

A light-emitting diode (LED) array with peak emission at 650 nm was used as the light source of singlet oxygen generation test. The irradiance of LED is 20 mW/cm². Singlet oxygen sensor green (SOSG) reagent (Life Technologies) was employed for the detection of singlet oxygen. H₄TBP, H₄TBC, TBP-Hf and TBC-Hf samples were prepared in 1 µM solutions/suspensions in water (for the nMOFs, the concentration was calculated as ligand equivalents). To 2 mL each of these solutions/suspensions, SOSG stock solution (5 µL at 5 mM) was added (final concentration=12.5 µM) before fluorescence measurement. For a typical measurement, fluorescence intensity was acquired on a spectrofluorophotometer (RF-5301PC, Shimadzu, Japan) with excitation at 504 nm and emission at 525 nm (slit width 1.5 nm/3 nm for ex/em). Fluorescence was measured after irradiation by LED for o (as background), 0.5, 1, 1.5, 2, 3, 4, 5, 7 and 10 min.

In vitro PDT efficacy

The cytotoxicity of TBC-Hf and TBC was evaluated in murine colorectal cancer cell CT26 and murine melanoma cell Bi6Fio, respectively. CT26 cells or Bi6Fio cells were seeded on 96-well plates at 1000 cells/well. The cells were treated with TBC-Hf and H₄TBC at various ligand concentrations (2, 4, 10, 15, 20, and 50 μ M base on ligand concentrations). A further incubation of 4 h was allowed followed by replacing the culture medium with 100 μ L of fresh medium. The cells were irradiated with LED light (650 nm) at 100 mW/cm² for 15 min (total light dose 90 J/cm²) or kept in dark, respectively. The cells were further incubated to achieve a total incubation time of 72 h. The cell viability was detected by (3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium) (MTS) assay (Promega, USA).

Abscopal Effect

The PDT efficacy of IDOi@TBC-Hf was investigated using a bilateral CT26 mouse colorectal cancer model and a bilateral MC38 mouse colorectal cancer model. Tumor bearing mice were established by subcutaneous inoculation of CT26 or MC38 cell suspension (1×10^6 cells per mouse on the right flank and 2×10^5 cells per mouse on the left flank) into 6-week female BALB/c or 6-week female C57BL/6 mice, respectively. When the right tumors reached 100 mm³, the CT26 tumor bearing mice received intratumoral injection only into the right tumors of PBS, H₄TBC, H₄TBC+IDOi, TBC-Hf, or IDOi@TBC-Hf at a ligand dose of 20 µmol/kg and IDOi dose of 1.5 mg/kg. Twelve hours post injection, CT26 tumor bearing mice were anesthetized with 2% (v/v) isoflurane and right flank tumors were irradiated with a 650 nm LED at 0.1 W/cm² for 15 min (90 J/cm²). IDOi@TBC-Hf without light irradiation served as a dark control. The nMOF injection and light irradiations were performed every three days for a total two injections and irradiations. For MC38 tumor model, mice received intratumoral injection only into the right tumors of PBS, TBC-Hf, or IDOi@TBC-Hf at a ligand dose of 20 µmol/kg and IDOi dose of 1.5 mg/kg. Twelve hours post injection, MC38 tumor bearing mice were anesthetized with 2% (v/v) isoflurane and right flank tumors were irradiated with a 650 nm LED at 0.1 W/cm² for 15 min (90 J/cm²). IDOi@TBC-Hf without light irradiation served as a dark control. The nMOF injection and light irradiations were performed only once on MC38 tumor bearing mice. For both CT26 and MC38 models, no nMOF injection or light irradiation was performed on the left tumors.

To evaluate the therapeutic efficacy, the tumor size was measured with a digital caliper every day. Tumor volumes were calculated as follows: (width² × length)/2. Finally, all mice were sacrificed when the tumor size of control group exceeded 2 cm³, and the excised tumors were photographed and weighed.

T cell blocking

The abscopal effect of IDOi@TBC-Hf was evaluated on bilateral subcutaneous MC38 model on C57BL/6 mice with CD4⁺ T cell or CD8⁺ T cell depletion. When the right tumors reached ~100 mm3, IDOi@TBC-Hf was intratumorally injected to the mice at a dose of 20 µmol/kg. Anti-CD4 (for CD4⁺ T cell depletion), anti-CD8 (for CD8⁺ T cell depletion), or mouse IgG (control) were intraperitoneally injected to the mice (200 µg/mouse/injection) on Day o and Day 5 post the first treatment. Twelve hours post injection, mice were anesthetized with 2% (v/v) isoflurane and the right tumors were irradiated with LED light irradiation (100 mW/cm², 90 J/cm², 650 nm). Single IDOi@TBC-Hf injections followed by single light irradiations were carried out. To evaluate the therapeutic efficacy, the tumor size was measured with a digital caliper every other day. Tumor volumes were calculated as follows: $(width^2 \times length)/2$.

ELISPOT

Tumor-specific immune responses to IFN- γ was measured *in vitro* by ELISPOT assay (Mouse IFN gamma ELISPOT Ready-SET-Go![®]; Cat. No. 88-7384-88; eBioscience). A Millipore Multiscreen HTS-IP plate was coated overnight at 4°C with anti-Mouse IFN- γ capture antibody. Single-cell suspensions of splenocytes were obtained from MC38 tumor-carrying mice and seeded onto the antibody-coated plate at a concentration of 2×10⁵ cells/well. Cells were incubated with or without KSPWFTTL (KSP) stimulation (10 µg/mL; in purity > 95%; PEPTIDE 2.0) for 48 h at 37 °C and then discarded. The plate was then incubated with biotin-conjugated anti-IFN- γ detection antibody at r.t. for 2 h, followed by incubation with Avidin-HRP for 2 h at r.t. AEC substrate solution (Sigma, Cat. AEC101) was added for cytokine spot detection.

Flow Cytometry

Tumors were harvested, treated with 1 mg/mL collagenase I (Gibco[™], USA) for 1 h, and ground by the rubber end of a syringe. Cells were filtered through nylon mesh filters and washed with PBS. The single-cell suspension was incubated with anti-CD16/32 (clone 93; eBiosciences) to reduce nonspecific binding to FcRs. Cells were further stained with the following fluorochrome-conjugated antibodies: CD45 (30-F11), CD3e (145-2C11), CD4 (GK1.5), CD8 (53-6.7), Foxp3 (FJK-16s), CD11b (M1/70), Ly6C (HK1.4), Ly6G (RB6-8C5), F4/80 (BM8), B220 (RA3-6B2), NKp46 (29A1.4) and PI staining solution (all from eBioscience). LSR FORTESSA (BD Biosciences) was used for cell acquisition, and data analysis was carried out using FlowJo software (TreeStar, Ashland, OR).

MHC-II expression

Tumors were harvested 12 hours after LED irradiation, treated with 1 mg/mL collagenase I, and ground by the rubber end of a syringe. Cells were filtered through nylon mesh filters and washed with PBS. The single cell suspension was incubated with anti-CD16/32 (clone 93; eBiosciences) for 10 minutes and then stained with CD45, CD3e, CD11b, Ly6G, B220, F4/80, CD11c and MHC-II. The expression levels of MHC-II on CD11b+Ly-6G+, CD3e-B220+, CD3e-CD11c+ and CD3e-CD11c-F4/80 populations were determined respectively.

ASSOCIATED CONTENT

Supporting Information. Experimental details and supporting results for the synthesis and characterization of H4TBC, TBC-Hf, IDOi@TBC-Hf, cellular uptake, in vitro PDT efficacy, abscopal effect, and antitumor immunity. This material is available free of charge via the Internet at http://pubs.acs.org."

AUTHOR INFORMATION

Corresponding Author

*wenbinlin@uchicago.edu

Author Contributions

[†]These authors contributed equally.

ACKNOWLEDGMENT

We thank Mr. Christopher Poon and Mr. Zekai Lin for experimental help. We acknowledge the National Cancer Institute (Uo1–CA198989), the University of Chicago Medicine Comprehensive Cancer Center (NIH CCSG: P30 CA014599), the Cancer Research Foundation, and the Ludwig Institute for Metastasis Research for funding support.

REFERENCES

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38 39 40

2. Stolik, S.; Delgado, J.; Perez, A.; Anasagasti, L. Journal of Photochemistry and Photobiology B: Biology 2000, 57 (2), 90-93.

3. (a) Ng, K. K.; Zheng, G. Chem. Rev. 2015, 115 (19), 11012-11042; (b) Lovell, J. F.; Jin, C. S.; Huynh, E.; Jin, H.; Kim, C.; Rubinstein, J. L.; Chan, W. C.; Cao, W.; Wang, L. V.; Zheng, G. Nature materials 2011, 10 (4), 324-332; (c) Huynh, E.; Leung, B. Y.; Helfield, B. L.; Shakiba, M.; Gandier, J.-A.; Jin, C. S.; Master, E. R.; Wilson, B. C.; Goertz, D. E.; Zheng, G. Nat Nanotechnol 2015, 10 (4), 325-332; (d) Carter, K. A.; Shao, S.; Hoopes, M. I.; Luo, D.; Ahsan, B.; Grigoryants, V. M.; Song, W.; Huang, H.; Zhang, G.; Pandey, R. K.; Geng, J.; Pfeifer, B. A.; Scholes, C. P.; Ortega, J.; Karttunen, M.; Lovell, J. F. Nature communications 2014, 5, 3546; (e) Kim, J.; Piao, Y.; Hyeon, T. Chem Soc Rev 2009, 38 (2), 372-390; (f) Wang, W.; Wang, L.; Li, Z.; Xie, Z. Chem Commun 2016, 52 (31), 5402-5405; (g) Padilla, R.; Maza, W. A.; Dominijanni, A. J.; Winkel, B. S.; Morris, A. J.; Brewer, K. J. Journal of Photochemistry and Photobiology A: Chemistry 2016, 322, 67-75; (h) Han, K.; Wang, S.-B.; Lei, Q.; Zhu, J.-Y.; Zhang, X.-Z. ACS nano 2015, 9 (10), 10268-10277; (i) Chen, J.-X.; Wang, H.-Y.; Li, C.; Han, K.; Zhang, X.-Z.; Zhuo, R.-X. Biomaterials 2011, 32 (6), 1678-1684; (j) Wang, C.; Tao, H.; Cheng, L.; Liu, Z. Biomaterials 2011, 32 (26), 6145-6154; (k) Idris, N. M.; Gnanasammandhan, M. K.; Zhang, J.; Ho, P. C.; Mahendran, R.; Zhang, Y. Nat. Med. 2012, 18 (10), 1580-1585; (1) Huang, P.; Lin, J.; Wang, S.; Zhou, Z.; Li, Z.; Wang, Z.; Zhang, C.; Yue, X.; Niu, G.; Yang, M. Biomaterials 2013, 34 (19), 4643-4654; (m) Park, Y. I.; Kim, H. M.; Kim, J. H.; Moon, K. C.; Yoo, B.; Lee, K. T.; Lee, N.; Choi, Y.; Park, W.; Ling, D. Adv Mater 2012, 24 (42), 5755-5761; (n) Levine, D. H.; Ghoroghchian, P. P.; Freudenberg, J.; Zhang, G.; Therien, M. J.; Greene, M. I.; Hammer, D. A.; Murali, R. Methods 2008, 46 (1), 25-32; (o) Cheng, Y.; C. Samia, A.; Meyers, J. D.; Panagopoulos, I.; Fei, B.; Burda, C. J. Am. Chem. Soc. 2008, 130 (32), 10643-10647; (p) Rieffel, J.; Chen, F.; Kim, J.; Chen, G.; Shao, W.; Shao, S.; Chitgupi, U.; Hernandez, R.; Graves, S. A.; Nickles, R. J. Adv Mater 2015, 27 (10), 1785-1790; (q) Luo, D.; Carter, K. A.; Miranda, D.; Lovell, J. F. Adv. Sci. 2016, DOI: 10.1002/advs.201600106.

4. (a) Lu, K.; He, C.; Lin, W. J. Am. Chem. Soc. **2014**, 136 (48), 16712-16715; (b) Lu, K.; He, C.; Lin, W. J. Am. Chem. Soc. **2015**, 137 (24), 7600-7603.

5. Castano, A. P.; Mroz, P.; Hamblin, M. R. *Nature Reviews Cancer* **2006**, *6* (7), 535-545.

Pardoll, D. M. *Nature Reviews Cancer* **2012**, *12* (4), 252-264.

7. Sharma, P.; Allison, J. P. *Science* **2015**, *348* (6230), 56-61.

8. Zou, W. *Nature Reviews Cancer* **2005**, *5* (4), 263-274.

9. (a) Gilboa, E. *Nature Reviews Cancer* **2004**, *4* (5), 401-411; (b) Munn, D. H.; Shafizadeh, E.; Attwood, J. T.; Bondarev, I.; Pashine, A.; Mellor,

A. L. The Journal of experimental medicine **1999**, 189 (9), 1363-1372.

10. Munn, D. H.; Zhou, M.; Attwood, J. T.; Bondarev, I.; Conway, S. J.; Marshall, B.; Brown, C.; Mellor, A. L. *Science* **1998**, *281* (5380), 1191-1193.

11. Yue, E. W.; Douty, B.; Wayland, B.; Bower, M.; Liu, X.; Leffet, L.; Wang, Q.; Bowman, K. J.; Hansbury, M. J.; Liu, C. *J. Med. Chem.* **2009**, *52* (23), 7364-7367.

12. Khleif, S.; Munn, D.; Nyak-Kapoor, A.; Mautino, M. R.; Kennedy, E.; Vahanian, N. N.; Link, C. J ASCO Annual Meeting Proceedings, 2014; p TPS3121.

13. Morris, W.; Volosskiy, B.; Demir, S.; Gándara, F.; McGrier, P. L.; Furukawa, H.; Cascio, D.; Stoddart, J. F.; Yaghi, O. M. *Inorg Chem* **2012**, *51* (12), 6443-6445.

14. (a) Park, J.; Jiang, Q.; Feng, D.; Mao, L.; Zhou, H.-C., *J. Am. Chem. Soc.* **2016**, *138* (10), 3518-3525; (b) Liu, J.; Yang, Y.; Zhu, W.; Yi, X.; Dong, Z.; Xu, X.; Chen, M.; Yang, K.; Lu, G.; Jiang, L., *Biomaterials* **2016**, *97*, 1-9.

15. Uyttenhove, C.; Pilotte, L.; Théate, I.; Stroobant, V.; Colau, D.; Parmentier, N.; Boon, T.; Van den Eynde, B. J. *Nat. Med.* **2003**, *9* (10), 1269-1274.

16. Newton, R. C.; Scherle, P. A.; Bowman, K.; Liu, X.; Beatty, G. L.; O'Dwyer, P. J.; Gajewski, T.; Bowman, J.; Schaub, R.; Leopold, L ASCO Annual Meeting Proceedings, 2012; p 2500.

17. Janeway, C. A.; Travers, P.; Walport, M.; Capra, J. D., Immunobiology: the immune system in health and disease. **2005**.

18. Siegel, R.; DeSantis, C.; Jemal, A. CA Cancer J. Clin. 2014, 64 (2), 104-117.

19. (a) Huxford, R. C.; Della Rocca, J.; Lin, W. *Curr. Opin. Chem. Biol.* **2010**, *14* (2), 262-268; (b) Lin, W.; Rieter, W. J.; Taylor, K. M. *Angewandte Chemie International Edition* **2009**, *48* (4), 650-658; (c) Della Rocca, J.; Liu, D.; Lin, W. Accounts Chem Res **2011**, *44* (10), 957-968; (d) He, C.; Liu, D.; Lin, W. Chem. Rev. **2015**, *115* (19), 11079-11108.

20. (a) Cui, Y.; Yue, Y.; Qian, G.; Chen, B. *Chem. Rev.* **2011**, *112* (2), 1126-1162; (b) Furukawa, H.; Cordova, K. E.; O'Keeffe, M.; Yaghi, O. M. *Science* **2013**, *341* (6149), 1230444; (c) Horcajada, P.; Gref, R.; Baati, T.; Allan, P. K.; Maurin, G.; Couvreur, P.; Ferey, G.; Morris, R. E.; Serre, C. *Chem. Rev.* **2011**, *112* (2), 1232-1268.

21. Palucka, K.; Banchereau, J. Nature Reviews Cancer 2012, 12 (4), 265-277.

22. Kousis, P. C.; Henderson, B. W.; Maier, P. G.; Gollnick, S. O. *Cancer Res.* 2007, 67 (21), 10501-10510.

