

TABLE I
 ANTIBACTERIAL ACTIVITY

Compd	Solvent	Zone of inhib, mm					
		<i>M.</i> <i>smegmatis</i>	<i>S.</i> <i>aureus</i>	γ <i>Strep.</i>	<i>E. coli</i>	<i>P.</i> <i>vulgaris</i>	<i>Ps.</i> <i>aeruginosa</i>
III	EtOH	25 ^b (35) ^c					
III	H ₂ O	17 (25)					
VIII	EtOH	25 (33)	20	24			
IX	H ₂ O						
X	EtOH	42 (54)	19	20	18 (22)	19	19
XI	H ₂ O	15 (21)	20				
XII	EtOH ^a						
XIII	H ₂ O	36 (69)					15
XIV	EtOH	18	17				15
XV	H ₂ O	19	18				
XVI-XVIII	EtOH ^c						

^a Suspension. ^b Complete inhibition. ^c Complete and partial inhibition.

greater odor than the other derivatives and gave qualitative P and S tests.

Synthesis.—The products were prepared using previously described methods^{2,7} whereby II,⁶ VI,¹⁵ VI,¹⁶ or VII¹⁶ and the appropriate amine, alcohol, hydrazine, or hydrazide were treated to yield VIII (C₁₁H₁₈N₃O₃PS, 125°, 66%), IX (C₈H₁₈N₃O₃PS, 81°, 49%), X (C₁₁H₁₇N₄O₄PS, 147°, 38%), XI (C₁₀H₂₁N₆O₆PS₂, >80°, 53%, deliquescent), XII (C₈H₁₅N₄O₂PS₂, >115°, 76%), XIII (C₁₆H₁₉N₈O₄PS₂, >105°, 94%), XV (C₁₂H₂₄N₆O₃PS₃, >53°, 14%, deliquescent), XVI (C₁₃H₂₇N₆O₄PS₃, >95°, 91%), XVII (C₁₃H₂₇N₆O₃PS₄, >130°, 84%), and XVIII (C₉H₁₃N₆PS₄, >175°, 28%). Ether was the solvent, except in the case of XIII (CH₃CN), XVI and XVII (CH₃CN-ether, 1:1), and X (no solvent). Addition of II to hydrazine or 4-methyl-3-thiosemicarbazide (CH₃CN) gave III (C₅H₁₂N₃O₃PS, 129°, 61%). Consecutive additions of ethanethiol and aziridine to IV (ether) produced V (C₆H₁₂N₃O₂PS₂, >147°, 57%). Similar addition of phenylhydrazine and ethyl carbazate to IV (ether) yielded XIV (C₁₃H₂₁N₆O₄PS₂, >65°, 25%) which separated as an oil before solidifying. One mixture (XI) was heated (35°, 1 hr) to ensure complete reaction.

Antibacterial Screening.—The antibacterial spectra were determined by saturating filter paper disks (12.7 mm) with 2 drops of an aqueous or alcoholic solution or suspension of the compound (20 mg/ml) and placing these on agar (Bacto Nutrient Agar, Difco Lab.) seeded with 48-hr culture broths (Nutrient Broth, Baltimore Biological Lab.) of the test organisms (0.5 ml). The microbial spectrum consisted of *Mycobacterium smegmatis*, *Escherichia coli*, *Proteus vulgaris*, *Staphylococcus aureus*, a γ *Streptococcus*, and *Pseudomonas aeruginosa* from the collections maintained at the Biology Department, University of Houston. V did not dissolve or suspend well in either solvent and was not tested. Alcohol controls were also run. The zones of inhibition around the disks were measured after 4 days of incubation (37°).

The MIC of VIII, X, XIII, INH, and alcohol against *M. smegmatis* and XIII and INH against *M. tuberculosis*¹⁷ were determined by a serial broth dilution method similar to that employed by Glasser and Doughty.¹⁸ The tubes, containing concentrations of 1000 to 1 μ g/ml, were examined for bacterial growth after incubation periods (37°) of 7 (*M. smegmatis*) and 10 days (*M. tuberculosis*).

Acknowledgments.—The author wishes to thank the following faculty and staff members of the University of Houston: Dr. M. R. Willcott for recording the nmr spectra, Dr. J. E. Evans and Mr. Harlan Brown for suggestions concerning the microbiological work, and Mr. N. O. Walker for preparing two starting materials and conducting some of the elemental analyses.

Some Aza Analogs of Amino Acids¹

TOMMY J. McCORD, LLEWELLYN D. BOOTH,
AND ALVIE L. DAVIS

Department of Chemistry, Abilene Christian College,
Abilene, Texas

Received April 5, 1968

In previous work, we described the preparation of 4-azaleucine (2-amino-3-dimethylaminopropionic acid) and its potent microbiological activities as a specific and competitive antagonist of leucine.² More recently, a compound identified as 2-amino-3-dimethylaminopropionic acid by structural studies was reported to have been isolated from culture media of a certain *Streptomyces* strain.³ In view of the uniqueness of 4-azaleucine as a natural product and our particular interest in aza analogs of amino acids,^{4,5} we undertook the synthesis of 4-azanorleucine, 5-azanorleucine, and 4-azanorvaline in order to determine their microbiological properties.

5-Azanorleucine was prepared in relatively poor over-all yield via an acetamidomalonic ester synthesis. Ethyl acetamidomalonic ester was condensed with N-benzyl-N-methyl-2-chloroethylamine in the presence of sodium ethoxide to form ethyl 2-acetamido-2-(N-benzyl-N-methyl-2-ethylamino)malonate. Acid hydrolysis of the condensation product gave the corresponding intermediate amino acid, 2-amino-4-(N-benzylmethylamino)butyric acid. Subsequent hydrolysis of the latter compound resulted in the formation of 5-azanorleucine.

The synthesis of 4-azanorleucine and 4-azanorvaline was accomplished by using the same general procedure in which the appropriately substituted amine underwent addition with 2-acetamidoacrylic acid to yield the

(15) B. C. Saunders, J. J. Stacey, F. Wild, and I. G. E. Wilding, *J. Chem. Soc.*, 699 (1948).

(16) B. S. Green, D. B. Sowerby, and K. J. Wihksne, *Chem. Ind. (London)*, 1306 (1960).

(17) Obtained from Jefferson Davis Hospital, Tuberculosis Laboratory, Houston, Texas, through the courtesy of Mr. Henry Gonzales.

(18) A. C. Glasser and R. M. Doughty, *J. Pharm. Sci.*, **51**, 1031 (1962).

(1) The support of this work in part by Grant R-085 from the Robert A. Welch Foundation of Houston, Texas, and by Public Health Service Research Grant No. AM-4060-08 from the National Institute of Arthritis and Metabolic Diseases is gratefully acknowledged.

(2) S. S. Smith, N. L. Bayliss, and T. J. McCord, *Arch. Biochem. Biophys.*, **102**, 313 (1963).

(3) A. D. Argoudelis, R. R. Herr, D. J. Mason, T. P. Pyke, and J. F. Zieserl, *Biochemistry*, **6**, 165 (1967).

(4) T. J. McCord, D. E. Cook, and L. G. Smith, *Arch. Biochem. Biophys.*, **105**, 349 (1964).

(5) T. J. McCord, D. C. Foyt, J. L. Kirkpatrick, and A. L. Davis, *J. Med. Chem.*, **10**, 353 (1967).

TABLE I
PHYSICAL CONSTANTS, YIELDS, AND ANALYTICAL DATA OF AMINO ACID ANALOGS

Compd	Yield, ^a %	Mp, ^a °C	<i>R_f</i> ^b	Formula ^c
5-Azanorleucine·2HCl	72	200–201	0.085 (A), 0.34 (B), 0.33 (C)	C ₅ H ₁₂ N ₂ O ₂ ·2HCl
4-Azanorleucine·2HCl	75	175–177	0.14 (A), 0.64 (B), 0.48 (C)	C ₅ H ₁₂ N ₂ O ₂ ·2HCl
4-Azanorvaline·2HCl	63	181–183	0.10 (A), 0.52 (B), 0.48 (C)	C ₆ H ₁₀ N ₂ O ₂ ·2HCl

^a Melting points and yields are for analytical samples. ^b The *R_f* values given for the appropriate solvents are indicated by letter: A, *n*-BuOH–AcOH–H₂O (4:1:1); B, 65% pyridine; C, 95% MeOH. ^c All compounds were analyzed for C, H, N.

corresponding 2-acetamido-3-alkylaminopropionic acid, followed by mild acid hydrolysis.

Biological Results.—Using *Escherichia coli* 9723 as the test organism, 5-azanorleucine, 4-azanorleucine, and 4-azanorvaline did not exert any observable inhibitory activity even at a concentration level of 2 mg/ml.

Experimental Section⁶

Ethyl 2-Acetamido-2-(N-benzyl-N-methyl-2-ethylamino)malonate.—To a solution of 21.7 g of ethyl acetamidomalonate in 100 ml of Mg-dried EtOH containing 2.3 g of Na was added 18.3 g of freshly distilled N-benzyl-N-methyl-2-chloroethylamine⁷ in small increments, and the reaction mixture was heated at reflux for 20 hr. The salt was removed by filtration, and the filtrate was taken to dryness *in vacuo*. Recrystallization of the residue (EtOH–H₂O) gave 5.0 g (14%) of product, mp 34–36°. *Anal.* (C₁₉H₂₈N₂O₅) C, H.

5-Azanorvaline (2-Amino-4-methylaminobutyric Acid) Dihydrochloride.—A sample of 4.0 g of ethyl 2-acetamido-2-(N-benzyl-N-methyl-2-ethylamino)malonate was hydrolyzed in the presence of 40 ml of 6 *N* HCl for 2 hr. The hydrolysis reaction mixture was taken to dryness under reduced pressure with warming. The resulting oil was freed of residual HCl by the repeated addition and evaporation of EtOH and the remaining residue was recrystallized (EtOH–Et₂O) to yield 2.1 g (65%) of product. The extreme hygroscopic nature of the sample was such that a melting point could not readily be determined and an acceptable analysis was not obtained. The material gave a positive test with ninhydrin, and the *R_f* values of this crude reaction product in *n*-BuOH–AcOH–H₂O (4:1:1), 65% pyridine, and 95% MeOH were 0.37, 0.80, and 0.63, respectively.

A sample of 0.40 g of the hygroscopic 2-amino-4-(N-benzyl-methylamino)butyric acid in 50% MeOH–H₂O was agitated in the presence of 50 mg of Pd black under about 3.52 kg/cm² pressure of H₂ for 3 hr. The catalyst was removed by filtration and the filtrate was acidified to a pH 1 by addition of concentrated HCl. After the resulting solution was reduced to dryness *in vacuo*, the residue was recrystallized from absolute EtOH–Et₂O to yield 0.20 g of product (see Table I).

2-Acetamido-3-ethylaminopropionic Acid.—The synthesis of this compound was patterned after a previously described procedure for the general preparation of 3-amino-substituted derivatives of acetamidopropionic acid.⁸ A mixture of 3.2 g of acetamidoacrylic acid and 50 ml of 33% aqueous EtNH₂ was allowed to stand at 40° for 72 hr. The reaction mixture was taken to dryness *in vacuo*, and the solid residue was recrystallized from absolute EtOH to yield 2.7 g (57%) of the product, mp 160–161°. *Anal.* (C₇H₁₄N₂O₃·H₂O) C, H.

4-Azanorleucine (2-Amino-3-ethylaminopropionic Acid) Dihydrochloride.—A solution of 1.0 g of 2-acetamido-3-ethylaminopropionic acid in 20 ml of 6 *N* HCl was heated under reflux for 4 hr. The reaction mixture was concentrated by removal of the acid under reduced pressure and absolute EtOH was then added. After chilling in the refrigerator, 0.80 g of product was obtained (see Table I).

2-Acetamido-3-methylaminopropionic Acid.—Using the general

(6) All melting points are corrected. The microanalyses were performed by International Chemical and Nuclear Corp., City of Industry, Calif. All *R_f* data were determined using the ascending technique of paper chromatography in the solvents indicated, and ninhydrin reagent was used for the development of the spots. Where analyses are indicated only by symbols of the elements, analytical results obtained by those elements were within ±0.4% of the theoretical values.

(7) J. B. Wright, E. H. Lincoln, R. V. Heinzelmann, and J. H. Hunter, *J. Am. Chem. Soc.*, **72**, 3536 (1950).

(8) L. Z. Eiger and J. P. Greenstein, *Arch. Biochem. Biophys.*, **19**, 467 (1948).

method⁸ from 3.6 g of 2-acetamidoacrylic acid there was recovered 2.7 g (60%) of product, mp 164–165°, lit.⁸ mp 164°.

4-Azanorvaline (2-Amino-3-methylaminopropionic Acid) Dihydrochloride.—2-Acetamido-3-methylaminopropionic acid (1 g) in 20 ml of 6 *N* HCl was heated at reflux for 4 hr. There was recovered 0.75 g of product from the hydrolysis reaction mixture after facilitating crystallization by the addition of absolute EtOH (see Table I).

Microbiological Assays.—A previously described inorganic salts–glucose medium⁹ was used for *E. coli* 9723, and experimental detail has been reported elsewhere.¹⁰

The amino acid analogs were dissolved in sterile H₂O and added aseptically to the previously autoclaved assay tubes. In all assays the amount of growth was determined photometrically at 625 mμ with a Bausch and Lomb Spectronic 20 spectrophotometer, in terms of absorbance readings of the turbid culture medium against a blank of uninoculated medium set at zero absorbance.

(9) E. H. Anderson, *Proc. Natl. Acad. Sci. U. S. A.*, **32**, 120 (1946).

(10) L. T. H. Dien, J. M. Ravel, and W. Shive, *Arch. Biochem. Biophys.*, **49**, 283 (1954).

Grignard Addition to 17α-Acetoxy-6α-methylprogesterone

ARVIN P. SHROFF

Division of Organic Chemistry, Ortho Research Foundation,
Raritan, New Jersey 08869

Received May 17, 1968

Grignard addition to a steroid α,β-unsaturated ketone has been reported by Musgrave¹ in 1951. He carried out 1,2 addition on cholestenone with methylmagnesium bromide to give 3-methylcholest-4-en-3-ol. A number of other investigators^{2–4} have used cuprous chloride and cuprous acetate along with the Grignard reagent to obtain 1,4 addition. We wish to report in this communication a one-step 1,2 addition of methylmagnesium bromide to the α,β-unsaturated ketone, 17α-acetoxy-6α-methylprogesterone (**1**).

An attempt to add a stoichiometric amount of methylmagnesium bromide to **1** at room temperature gave only precipitates owing to complex formation. Varying the proportions of reactants and the temperature of the reaction gave mixtures of products. However, the use of a large excess of Grignard reagent at room temperature gave 3ξ,17α-dihydroxy-3ξ,6α-dimethylpregn-4-en-20-one 17-acetate (**2**) if acid was avoided in isolating the desired product. If acetic acid or dilute HCl was added in the work-up of the reaction, then dehydration occurred and 17α-acetoxy-3,6-dimethylpregn-3,5-dien-20-one (**3**) was isolated. When

(1) O. C. Musgrave, *J. Chem. Soc.*, 3121 (1951).

(2) Merck and Co., Inc., British Patent 877,987 (1961).

(3) Hiromu Mori, *Chem. Pharm. Bull. (Tokyo)*, **12**, 1224 (1964).

(4) Bohumil Pele, Czechoslovakian Patent 111,047 (1964).