

Oxidative Dimerization of Quinolinic Nitroxides in the Presence of Trichloro- and Trifluoro- Acetic Acid. Crystal Structures of 6,6'-bis-(1-oxide-1,2,6,8a-tetrahydroquinoline)ylidene and of 2,3-Diphenylquinoline

Patricia Carloni, Elisabetta Damiani, Lucedio Greci,* and Pierluigi Stipa

Dipartimento di Scienze dei Materiali e della Terra, Università, Via Brecce Bianche,
I-60131 Ancona, Italy.

Corrado Rizzoli, Paolo Sgarabotto,* and Franco Ugozzoli

Istituto di Strutturistica Chimica, Università, Centro di Studio per la Strutturistica Diffraattometrica del C.N.R.,
Viale delle Scienze, I-43100 Parma, Italy.

(Received in UK 22 March 1993)

Abstract: Quinolinic nitroxides **1a-c** react with trichloro- (TCA) and trifluoro- (TFA) acetic acid to give dimers **3a-c** and quinolines **4a-c** as the main products. Products **3a-c** are explained as arising via the intermediate formation of the radical cation **5**, which forms the final dimer through an oxidative dimerization promoted by the TCA or TFA. The formation of products **4a-c** is explained by a Wagner-Meerwein type transposition of the phenyl group from C-2 to C-3. The disproportionation, which generally occurs in the presence of acid for nitroxides having the N=O function conjugated with a π -system, has been excluded even though quinoneimine N-oxides **2a-c** were isolated. The structure of compounds **3a** and **4a** were determined by crystal X-ray analysis.


The use of nitroxide radicals in the last decade has received a great development in several fields for many applications. They are studied as "contrast agents" in MR-imaging,¹ used in pharmacological investigations² and more recently they have been proposed as superoxide dismutation mimics.³ Nitroxides are used for studying membranes,⁴ enzymes⁵ and for oxygen determination (oxymetry).⁶ Some of them are used or generated *in situ* via radical trapping by suitable scavengers, mostly PBN (N-*tert*-butyl- α -phenylnitrone)⁷ and DMPO (5,5-dimethyl-1-pyrroline-N-oxide), in particular media such as acidic solutions or complex biological systems.⁸ Several derivatives of the tetramethyl-piperidine and -pyrrolidine are also used as probes for studying structural properties of polymers.⁹ Consequently, the study of their behaviour in different conditions can be useful.

Most of the nitroxides used or generated have the unpaired electron localized in the nitroxide function and are structurally similar to the tetramethyl-piperidine,¹⁰ -pyrrolidine¹⁰ and imidazolyl series.¹¹ Instead, the indolinic and the quinolinic nitroxides,¹² and their derivatives,¹² have the N=O group in a conjugated position with the π system of the benzenic ring of the indolic or the quinolinic nucleus and they possess a very high stability if compared to diaryl and alkyl-aryl-nitroxides.

Since we are developing the applications of the latter in biology,¹³ in polymers¹⁴ and in the study of reaction mechanisms on the basis of their reactivity,^{15,16} in the present paper we report the study of some quinolinic nitroxides under acid treatment by TCA and TFA.


RESULTS

Quinolinic nitroxides **1a-c** when treated at room temperature for 10 minutes with a triple quantity of TCA or TFA in a benzene solution carefully degassed with nitrogen or argon, gave the dimers **3a-c** and the quinolines **4a-c**, as the main products, together with traces of quinoneimine N-oxides **2a-c** (Scheme 1).



Scheme 1

C-4 for **4a**, at 1.16 and 3.69 ppm due to the hydrogens of the substituent at C-4 for **4b** and at 1.64 ppm due to the hydrogens of the methyl groups of the *iso*-propanol substituent at C-4 for **4c**. Compound **4c** also shows at 2.3 ppm the signal for the hydrogens of the methyl group at C-2. Quinoneimine N-oxides **2a-c**, which were also obtained by oxidizing the nitroxides **1a-c** in acetonitrile/water with cerium ammonium nitrate (see experimental) were identified by their analytical and spectroscopic data.



Scheme 2




Fig. 1. Perspective view of 6,6'-bis-(1-oxide-1,2,6,8a-tetra-hydro-quinoline)-ylidene 3a.

Their $^1\text{H-NMR}$ spectra show characteristic signals (see Table 3) in the region between 6.6 and 8.0 ppm, due to the hydrogens of the benzenic ring of the quinolinic moiety, and between 5.6 and 6.6 ppm due to the proton at C-3. Moreover compounds 2b and 2c also show the signals characteristic of the hydrogens of their substituents at C-4. The IR spectrum of compound 2c shows the OH band at 3330

cm^{-1} . Amines 8a-c (Scheme 4), which were prepared by refluxing nitroxides 1a-c with Fe/AcOH, were identified by their analytical and spectroscopic data (Table 3). The $^1\text{H-NMR}$ spectra show characteristic signals between 6.5 and 7.8 ppm due to the hydrogens of the benzenic ring, at ca. 5.8-5.9 ppm due to the hydrogen at C-3 for 8a and 8c, at 3.38 ppm due to the peak of the two hydrogens at C-3 for 8b, and at 1.6 ppm due to the hydrogens of the substituent at C-4 for 8c. The IR spectra show the bands due to the NH at ca. 3200-3400 cm^{-1} , and for compound 8c the OH band at 3560 cm^{-1} . All mass spectra gave the expected molecular ion peaks.

The reactions carried out with an excess of TFA gave the same products, in higher yields, whereas in the above conditions some starting nitroxide was recovered.

Molecular geometry

Figures 1 and 2 show a drawing of the molecule of compounds 3a and 4a with the arbitrary numbering scheme adopted in the crystallographic analysis; Table 1 lists bond distances and angles.*@

The conformational geometry of the molecule was deduced from the torsion angles of Table 1 and from the analysis of the planarity reported in Table 2.

For both compounds, bond distances and angles are mostly as expected from hybridization of the atoms. The main differences in the skeleton of the two-condensed six-membered rings concern bond distances and angles, according to the different hybridization of the carbon atom in position two, being sp^3 in compound 3a and sp^2 in 4a. Nevertheless the quinoline moiety adopts a conformation substantially planar in both compounds, being 1.3(1) and 0.4(1)* the values of the dihedral angles between the two mean planes of the six-membered rings in compounds 3a




Fig. 2. Perspective view of 2,3-diphenyl-quinoline 4a.

* The atomic coordinates are available from the Director of the Cambridge Crystallographic Data Centre, University Chemical Laboratories, Lensfield Road, Cambridge CB2 1EW.

Any request should be accompanied by the full literature citation for this communication.

@ Supplementary data available: Atomic fractional coordinates, list of thermal parameters and structure factors.

and 4a, respectively. In compound 4a the geometry of the quinoline system is in good agreement with that found in similar molecules reported in literature,¹⁷ slight differences in bond distances and angles being

Table 1. Bond Distances (Å), Angles (°) and Selected Torsion Angles(°) with e.s.d.'s in Parentheses.

3a			4a			3a			4a		
Bond Distances (Å)			C(33)-C(34)	1.368(6)		C(2)-C(11)-C(16)	119.4(4)				
O(1)-N(1)	1.282(4)		C(34)-C(35)	1.379(5)		C(12)-C(11)-C(16)	118.6(4)				
N(1)-C(2)	1.506(4)	1.326(3)	C(35)-C(36)	1.396(4)		C(11)-C(12)-C(13)	120.1(5)				
N(1)-C(10)	1.373(5)	1.371(3)	Angles (°)			C(12)-C(13)-C(14)	120.9(5)				
C(2)-C(3)	1.505(6)	1.440(3)	O(1)-N(1)-C(2)	115.2(3)		C(13)-C(14)-C(15)	119.1(6)				
C(2)-C(11)	1.528(6)		O(1)-N(1)-C(10)	120.0(3)		C(14)-C(15)-C(16)	120.7(5)				
C(2)-C(21)	1.536(6)	1.491(3)	C(2)-N(1)-C(10)	124.7(3)	119.2(3)	C(11)-C(16)-C(15)	120.6(4)				
C(3)-C(4)	1.327(6)	1.370(3)	N(1)-C(2)-C(3)	109.9(3)	123.0(3)	C(2)-C(21)-C(22)	120.6(4)	121.2(2)			
C(3)-C(31)		1.491(3)	N(1)-C(2)-C(11)	108.2(3)		C(2)-C(21)-C(26)	120.7(4)	120.3(2)			
C(4)-C(5)	1.432(5)	1.403(3)	N(1)-C(2)-C(21)	106.6(3)	114.3(3)	C(22)-C(21)-C(26)	118.7(4)	118.5(3)			
C(5)-C(6)	1.386(6)	1.414(3)	C(3)-C(2)-C(11)	106.9(3)		C(21)-C(22)-C(23)	121.3(5)	120.4(3)			
C(5)-C(10)	1.416(5)	1.419(3)	C(3)-C(2)-C(21)	111.7(3)	122.7(2)	C(22)-C(23)-C(24)	119.6(5)	120.2(3)			
C(6)-C(7)	1.401(5)	1.365(4)	C(11)-C(2)-C(21)	113.4(4)		C(23)-C(24)-C(25)	119.7(6)	119.8(4)			
C(7)-C(8)	1.412(7)	1.401(5)	C(2)-C(3)-C(4)	124.9(4)	117.0(2)	C(24)-C(25)-C(26)	120.9(5)	119.9(3)			
C(7)-C(7')	1.454(6)		C(2)-C(3)-C(31)		122.9(3)	C(21)-C(26)-C(25)	119.8(4)	121.1(3)			
C(8)-C(9)	1.369(6)	1.364(3)	C(4)-C(3)-C(31)		120.2(3)	C(3)-C(31)-C(32)		119.9(2)			
C(11)-C(12)	1.382(6)		C(3)-C(4)-C(5)	121.3(4)	121.8(3)	C(3)-C(31)-C(36)		122.1(2)			
C(11)-C(16)	1.390(5)		C(4)-C(5)-C(6)	122.8(3)	124.1(3)	C(32)-C(31)-C(36)		117.9(2)			
C(12)-C(13)	1.392(8)		C(4)-C(5)-C(10)	119.3(4)	117.5(2)	C(31)-C(32)-C(33)		120.2(3)			
C(13)-C(14)	1.371(8)		C(6)-C(5)-C(10)	117.9(4)	118.4(2)	C(32)-C(33)-C(34)		121.2(3)			
C(14)-C(15)	1.374(7)		C(5)-C(6)-C(7)	123.2(4)	120.5(3)	C(33)-C(34)-C(35)		119.4(4)			
C(15)-C(16)	1.379(7)		C(6)-C(7)-C(7')	122.8(4)		C(34)-C(35)-C(36)		119.9(3)			
C(21)-C(22)	1.378(6)	1.384(4)	C(6)-C(7)-C(8)	116.4(4)	120.8(3)	C(31)-C(36)-C(35)		121.5(3)			
C(21)-C(26)	1.385(7)	1.382(4)	C(7')-C(7)-C(8)	120.8(4)		Selected Torsion Angles(°)					
C(22)-C(23)	1.383(7)	1.393(4)	C(7)-C(8)-C(9)	122.7(4)	120.6(3)	O(1)-N(1)-C(2)-C(11)	62.4(4)				
C(23)-C(24)	1.376(10)	1.374(5)	C(8)-C(9)-C(10)	119.3(4)	120.0(3)	O(1)-N(1)-C(2)-C(21)	-60.0(4)				
C(24)-C(25)	1.368(8)	1.374(5)	N(1)-C(10)-C(5)	119.7(4)	121.6(2)	N(1)-C(2)-C(3)-C(4)	2.7(6)	-2.0(5)			
C(25)-C(26)	1.388(7)	1.385(4)	N(1)-C(10)-C(9)	119.8(4)	118.7(3)	C(21)-C(2)-C(3)-C(31)		-5.8(5)			
C(31)-C(32)	1.401(4)		C(5)-C(10)-C(9)	120.5(4)	119.7(3)	C(3)-C(2)-C(21)-C(22)		-61.7(5)			
C(31)-C(36)	1.379(3)		C(2)-C(11)-C(12)	121.8(4)		C(4)-C(3)-C(31)-C(32)	136.4(4)				
C(32)-C(33)	1.392(6)										

interpreted in terms of steric interactions of the two adjacent phenyl substituents.

The crystal structure of 3a consists of symmetric molecules for the presence of a two-fold crystallographic axis perpendicular to the C(7)-C(7') bond and inclined by 78.7° with respect to the mean planes of the two quinolinic rings. In this compound the geometry of the quinoline moiety is comparable to that found in the 1,2-dihydro-2,2-diphenyl-quinoline-1-oxyl previously studied;¹⁸ also the orientation of the N-oxide group with respect to the quinoline ring is similar and in agreement with its more marked planar arrangement. In fact the oxygen atom is out of the C-N-C plane by 0.037(3) Å corresponding to an angle of 1.6(4)° (Table 2). The N-O bond length of 1.282(4) Å, which is intermediate between the values characteristic for N=O double bond, 1.20 Å, and N-O-R single bond, 1.44 Å,¹⁹ is in agreement with that found for heterocycle N-oxides.²⁰

Table 2. Analysis of the Planarity

3a

4a

3a

4a

3a

4a

a) Distances ($\text{\AA} \times 10^3$) of relevant atoms from the mean plane with e.s.d.'s in parentheses: starred atoms were not used to define the plane.

Plane A: N(1)C(2)C(3)C(4)C(5)C(10)			Plane C: C(11)-C(16)			Plane E: C(31)-C(36)		
N(1)	9(3)	3(3)	C(11)	10(4)		C(31)		2(4)
C(2)	-22(4)	-10(3)	C(12)	-7(4)		C(32)		-3(5)
C(3)	10(4)	8(3)	C(13)	-1(6)		C(33)		1(5)
C(4)	9(4)	0(3)	C(14)	4(6)		C(34)		2(5)
C(5)	-7(3)	-7(3)	C(15)	5(6)		C(35)		-2(5)
C(10)	0(4)	5(3)	C(16)	-11(4)		C(36)		0(5)
O(1)*	8(3)		C(2)*	-58(4)		C(3)*		-66(4)
C(21)*		-88(3)	Plane D: C(21)-C(26)			b) Angles ($^\circ$) between planes		
C(31)*		78(3)	C(21)	-2(4)	1(4)	A - B	1.3(1)	0.4(1)
Plane B: C(5)-C(10)			C(22)	7(5)	9(4)	A - C	107.3(1)	
C(5)	9(3)	-1(3)	C(23)	-9(6)	-12(4)	A - D	105.0(1)	118.7(1)
C(6)	-2(4)	-2(4)	C(24)	-1(6)	3(4)	A - E		133.5(1)
C(7)	-9(3)	3(4)	C(25)	6(5)	9(4)	C - D	65.1(2)	
C(8)	11(3)	0(4)	C(26)	-3(5)	-9(4)	D - E		118.0(1)
C(9)	-5(4)	-2(3)	C(2)*	44(4)	-91(4)	AB - A'B'	157.4(1)	
C(10)	-17(4)	2(3)						


DISCUSSION

2,2,6,6-Tetramethylpiperidine-1-oxyl when treated with hydrochloric acid undergoes a disproportionation reaction forming the corresponding oxoammonium ion and hydroxylamine;²¹ no data are reported with car-

boxylic acid. Diaryl- and arylalkyl- nitroxides undergo a spontaneous disproportionation to the corresponding quinoneimine-N-oxides and amines,²² which is faster in the presence of carboxylic acids. Indolinonic nitroxides slowly afford to the corresponding amines and quinoneimine-N-oxides in the presence of TFA.²³

Nitroxides 1a-c, which are perfectly stable in solution, are transformed by TCA or TFA treatment, into the corresponding dimers 3a-c through a dimerization forming a carbon-carbon bond followed by an oxidative step. Dimerization involving only the nitroxide function was observed in the piperidino- and pyrrolidino-nitroxides series in particular conditions,²⁴ whereas spontaneous dimerization with formation of a carbon-carbon or carbon-oxygen bond was only observed with vinyl nitroxides.²⁵

In our case, the dimerization could be explained as proceeding through coupling of two molecules of the radical cation of the nitroxide at C-6 followed by deprotonation of the intermediate, forming the N,N'-dihydroxy dimer 6 (Scheme 2). Since the reactions were carried out in the absence of oxygen, the last



Scheme 3

oxidative step in which the N,N'-dihydroxy dimer is transformed into 3, is a complicated task. The possible candidate for oxidation could be the starting nitroxide which could act as hydrogen abstractor forming the hydroxylamine 7 in agreement with the equilibrium depicted in Scheme 3.


The fact that the hydroxylamine 7 corresponding to the nitroxide was never observed, could be due to its transformation to the quinoline 4, through protonation, transposition of a phenyl group and elimination of water, as shown in Scheme 3. The transformation of 7 into 4 may also contribute to shift the equilibrium depicted in Scheme 3 to the right. Since the nitroxide 1c is methyl and phenyl substituted at C-2, for compound 4c the migration of the phenyl group, which in general occurs better than alkyl group could be expected. Thus, on this basis, established in the Wagner-Meerwein type transpositions study,²⁶ we assigned to 4c the structure reported in Scheme 1. The spectroscopic data of 4c are insufficient for its structure determination.

The other unexpected candidate to oxidize 6 to 3 could be the acid present. Since it has been recently reported that TFA can promote oxidative processes,²⁷ we examined this possibility. The N,N'-dihydroxy dimer 6a, obtained from 3a by reduction with hydrazo-benzene, when treated with TFA was suddenly retransformed into the dimer 3a.

The presence of quinoneimine N-oxide 2a-c is difficult to explain. At first this product was thought to derive from the disproportionation of two molecules of nitroxide (Scheme 4) as occurs in the case of indolinic nitroxides. In order to verify this possibility, amines 8a-c were synthesized to obtain authentic samples for assaying their presence in the reaction mixture. The amines 8a-c were never found. The total absence of the amine from the reaction mixture and the fact that the yield of the quinoneimine 2a increases by 100%, when carrying out the reaction of nitroxide 1a in the presence of air, suggest that it is formed by an interaction of the nitroxide radical cation and oxygen of air. This kind of interaction was observed by us²⁸ and others²⁹ and proceeds as reported in the literature.

Generally, vinyl ethers in the presence of acid, undergo cleavage of the ethereal

group.³⁰ According to the literature reports the dimer 3b recovered in the reaction with acid, and the amine 8b, bear a ketonic group at C-4; we have no explanation at the moment for the fact that quinoneimine N-oxide 2b and quinoline 4b keep the ethereal function.



Scheme 4

EXPERIMENTAL

Melting points were measured on an Electrochemical Melting Point Apparatus and are uncorrected. IR spectra were recorded on a Perkin-Elmer model 298 spectrophotometer. ¹H-NMR spectra were recorded in CDCl₃ on a Varian Gemini 200 at 200 MHz. Mass spectra were recorded on a Carlo Erba QMD 1000 spectrometer. EPR spectra were recorded on a Varian E4 spectrometer.

Nitroxides 1a,b were prepared as reported in the literature.¹²

Synthesis of nitroxide 1c. A Grignard reagent solution of CH₃MgI (60 mmoles in 50 ml of Et₂O) was

added to a 2-phenyl-4-carbethoxyquinoline-N-oxide³¹ solution (10 mmol in 80 ml of THF) under stirring in N₂ atmosphere and at room temperature. After 2h, the mixture was treated with a 0.5 M solution of NH₄Cl and extracted with CHCl₃. The organic layer was dried on Na₂SO₄ and oxidized with PbO₂ under magnetic stirring for 2h. The solution was filtered, evaporated to dryness and the residue was chromatographed on SiO₂ column, eluting with benzene. The red nitroxide radical was isolated as a pure, red powder after concentration. Yield 60%. M.p.= 148°C; ν_{max} (Nujol) 3480 (OH), 1600 cm⁻¹; m/z 294 (M⁺), 236 (100%); h_{fcc}'s in CHCl₃: a_N=10.48, a_H(3)=1.07, a_H(5,7)=1.25, a_H(4,6)=3.35 Gauss; Analysis: Calcd. for C₁₉H₂₀NO₂: H, 6.85; C, 77.52; N, 4.76; Found: H, 6.72; C, 77.50; N, 4.80.

Preparation of Quinoneimine N-oxides 2a-c. 0.5 mmol of nitroxides 1a-c were dissolved in 10 ml CH₃CN. An aqueous solution of cerium ammonium nitrate (Ce^{IV}) (0.5 mmol in 1 ml of water) was added dropwise to the nitroxide solution. An additional mmol of solid Ce^{IV} salt was then added in small amounts to the reaction until complete disappearance of the nitroxide, by following the reaction through T.L.C.. The reaction was then poured into 20 ml water, extracted with 40 ml CHCl₃, washed twice with 20 ml water, dried over Na₂SO₄ and concentrated. The residue was chromatographed on preparative silica gel plates eluting with cyclohexane/ethyl acetate 8:2 for 2a-b, and cyclohexane/ethyl acetate 6:4 for 2c. From the yellow spots products 2a-c were obtained, crystallized from petroleum ether 80°-100°. Yields as follows: 2a (90%), 2b (65%), 2c (60%).

Preparation of Amines 8a-c. 1 mmol of nitroxides 1a-c, 5 ml glacial acetic acid and 500 mg of iron were refluxed altogether for 5 min. The iron was filtered off and the organic layer evaporated to dryness. The residue was taken up with 30 ml CHCl₃ and neutralized with 20 ml 10% Na₂CO₃. The organic layer was then washed twice with 20 ml water, dried over Na₂SO₄ and concentrated to a small volume. The residue was chromatographed on preparative silica gel plates eluting with cyclohexane/ethyl acetate 1:9 for 8a-b and cyclohexane/ethyl acetate 7:3 for 8c. The large fluorescent spot yielded 70% for 8a, 68% for 8b, and 64% for 8c (crystallized from hexane).

Reaction of 1a-c with Trichloroacetic Acid. Solutions of nitroxides 1a-c (1 mmol in 10 ml of benzene) and of Trichloroacetic acid (TCA) (3 mmol in 10 ml of benzene) were degassed separately under argon at room temperature, in a two-legged test-tube for 10 mins. The two solutions were then mixed together and degassed for a further 10 mins. The dark red colour of the nitroxide solution turned dark violet immediately on addition of the acid. The reaction was then neutralized with 10% Na₂CO₃ (50 ml), extracted with benzene (2 x 20 ml). The organic layer was washed with water (2 x 20 ml), dried over Na₂SO₄, evaporated to a small volume and then chromatographed on preparative silica gel plates eluting with cyclohexane/ethyl acetate 8:2 for nitroxides 1a-b, and cyclohexane/ethyl acetate 6:4 for nitroxide 1c. From the red spots, nitroxides 1a-c were recovered. Yields as follows: 1a (180 mg, 60%), 1b (193 mg, 56%), 1c (no nitroxide was recovered because of its instability). From the dark violet spots extracted with ethyl acetate and evaporated to dryness, fine bright fluorescent violet crystals of dimers 3a-c were extracted. Yields as follows: 3a (35 mg, 12%), 3b (50 mg, 16%), 3c (60 mg, 20%). The fluorescent spot was identified as 4-substituted 2,3-quinoline (4a-c). Yields as follows: 4a (26 mg, 8%), 4b (34 mg, 10%), 4c (25 mg, 7%). The yellow spot was identified as quinoneimine N-oxides 2a-c by comparison with authentic samples. Yields are as follows: 2a (10 mg, 3%), 2b (11 mg, 3%), 2c (15 mg, 5%). Other minor unidentified spots were extracted from the reactions.

Reaction of 1a-c with Trifluoroacetic Acid. Solutions of nitroxides 1a-c (1 mmol in 10 ml of benzene) and of Trifluoroacetic acid (TFA) (3 mmol in 10 ml of benzene) were reacted, worked up and chromatographed in the same way as for the reactions with TCA. From the red spots, nitroxides 1a-c were recovered. Yields as follows: 1a (135 mg, 45%), 1b (113 mg, 33%), 1c (60 mg, 20%). From the dark violet spots, dimers 3a-c resulted. Yields are as follows: 3a (48 mg, 16%), 3b (66 mg, 21%), 3c (78 mg, 27%). From the fluorescent spots quinolines 4a-c were recovered. Yields as follows: 4a (15 mg, 5%), 4b (23 mg, 7%), 4c (16 mg,

Table 3. Analytical and Spectroscopic Data of Compounds 2a-c, 3a-c, 4a-c and 8a-c.

Comp.	NMR (δ , ppm) in CDCl_3	IR (ν , cm^{-1})	Mass Spectra	Melting Point	Calc. % Found (%)
2a	6.23 (1H, d, $J=1.9$ Hz, H-C5); 6.60 and 6.67 (2H, AB, $J=10.1$ Hz, H-C3 and H-C4); 6.64 (1H, dd, $J=1.9$ and 10.2 Hz, H-C7); 7.34 (10H, m, arom.); 7.95 (1H, d, $J=10.2$ Hz, H-C8)	1625, 1580, 1550	313 (M^+), 313 (100%)	192°C	H 4.82 (4.77) C 80.49 (80.36) N 4.47 (4.50)
2b	1.43 (3H, t, $J=7.0$ Hz, OCH_2CH_3); 3.97 (2H, q, $J=7.0$ Hz, OCH_2CH_3); 5.59 (1H, s, H-C3); 6.63 (1H, dd, $J=2.1$ and 10.2 Hz, H-C7); 6.70 (1H, d, $J=1.8$ Hz, H-C5); 7.33 (10H, m, arom.); 7.92 (1H, d, $J=10.2$ Hz, H-C8)	1660, 1625, 1595, 1555	357 (M^+), 357 (100%)	183°C	H 5.36 (5.21) C 77.29 (77.17) N 3.92 (4.03)
2c	1.67 (6H, s, $\text{HO-C}(\text{CH}_3)_2$); 2.05 (3H, s, CH_3); 6.42 (1H, s, H-C3); 6.60 (1H, dd, $J=2.0$ and 10.2 Hz, H-C7); 7.18 (1H, d, $J=2.0$ Hz, H-C5); 7.36 (5H, m, arom.); 7.97 (1H, d, $J=10.2$ Hz, H-C8)	3330, 1615, 1580, 1545	309 (M^+), 309 (100%)	147°C	H 6.19 (6.05) C 73.77 (73.64) N 4.53 (4.70)
3a	not resolved	1600	594 (M^+), 501 (100%)	225°C	H 5.08 (4.99) C 84.82 (84.72) N 4.71 (4.88)
3b	3.40 (2H, s, $\text{H}_2\text{-C3}$); 6.71 (2H, m, arom.); 7.29 (11H, m, arom.); 7.74 (1H, bdd, $J=7.6$ and 1.3 Hz, arom.)	1730, 1610	626 (M^+), 497 (100%)	182°C	H 4.82 (4.72) C 80.49 (80.31) N 4.47 (4.55)
3c	1.67 (6H, s, $\text{HO-C}(\text{CH}_3)_2$); 2.05 (3H, s, CH_3); 6.43 (1H, s, H-C3); 6.61 (1H, dd, $J=1.9$ and 10.2 Hz, H-C7); 7.045 (1H, d, $J=1.8$ Hz, H-C5); 7.34 (5H, m, arom.); 7.98 (1H, d, $J=10.2$ Hz, H-C8)	3377, 1733, 1615, 1591	586 (M^+), 524 (100%)	151°C	H 6.53 (6.35) C 77.79 (77.60) N 4.77 (4.85)
4a	7.26 (9H, m, arom.); 7.46 (2H, m, arom.); 7.61 (1H, t, $J=6.1$ Hz, H-C6); 7.78 (1H, t, $J=6.1$ Hz, H-C7); 7.9 (1H, d, $J=6.1$ Hz, H-C5); 8.24 (1H, s, H-C4)	1615, 1590, 1020, 695	281 (M^+), 280 (100%)	77°C	H 5.37 (5.28) C 89.65 (89.52) N 4.98 (5.10)
4b	1.16 (3H, t, OCH_2CH_3); 3.69 (2H, q, OCH_2CH_3); 7.28 (11H, m, arom.); 7.60 (1H, bt, $J=7.6$ Hz, H-C6); 7.78 (1H, bt, $J=7.3$ Hz, H-C7); 8.23 (1H, bd, $J=8.2$ Hz, H-C5)	1580, 1570, 1105, 1055, 695	325 (M^+), 296 (100%)	122°C	H 5.89 (5.70) C 84.89 (84.65) N 4.30 (4.35)
4c	1.64 (6H, s, $\text{HO-C}(\text{CH}_3)_2$); 2.30 (3H, s, CH_3); 7.23 (3H, m, arom.); 7.46 (3H, m, arom.); 7.66 (1H, bt, $J=7.6$ Hz, H-C7); 8.06 (1H, bd, $J=7.6$ Hz, H-C5); 8.70 (1H, bd, $J=7.9$ Hz, H-C8)	3260, 1555, 1135, 760, 705	277 (M^+), 218 (100%)	150°C	H 6.90 (6.76) C 82.28 (82.22) N 5.05 (5.13)
8a	1.65 (1H, bs, NH); 5.91 (1H, d, $J=9.9$ Hz, H-C3); 6.51 (1H, d, $J=9.9$ Hz, H-C4); 6.52 (1H, t, $J=8.4$ Hz, H-C7); 6.62 (1H, d, $J=7.3$, arom.); 6.62 (1H, d, $J=7.4$, arom.); 7.01 (1H, t, $J=7.7$ Hz, H-C5); 7.3 (10H, m, arom.)	3400, 1605	283 (M^+), 206 (100%)	77°C	H 6.05 (6.00) C 89.01 (88.89) N 4.94 (5.03)
8b	3.38 (2H, s, $\text{H}_2\text{-C3}$); 5.16 (1H, bs, NH); 6.69 (1H, ddd, $J=8.0$, 7.0 and 1.0 Hz, arom.); 6.75 (1H, dd, $J=7.0$ and 1.0 Hz, arom.); 7.3 (11H, m, arom.); 7.73 (1H, dd, $J=8.0$ and 1.0 Hz, arom.)	3240, 1645, 1610, 1580	299 (M^+), 222 (100%)	205°C	H 5.72 (5.60) C 84.25 (84.17) N 4.68 (4.77)
8c	1.60 (6H, s, $\text{HO-C}(\text{CH}_3)_2$); 1.69 (3H, s, CH_3); 5.81 (1H, s, H-C3); 6.58 (1H, dd, $J=7.6$ and 1.5 Hz, H-C4); 6.67 (1H, dt, $J=7.6$ and 1.5 Hz, H-C5); 7.02 (1H, dt, $J=7.9$ and 1.5 Hz, H-C6); 7.29 (3H, m, arom.); 7.49 (2H, m, arom.); 7.81 (1H, dd, $J=7.9$ and 1.5 Hz, H-C7)	3560, 3370, 1710, 1600	279 (M^+), 264 (100%)	54°C	H 7.58 (7.42) C 81.68 (81.50) N 5.01 (5.20)

6%). From the yellow spots, products **2a-c** were obtained. Yields as follows: **2a** (9 mg, 3%), **2b** (7 mg, 4%), **2c** (3 mg, 1%). The reaction was repeated also with an excess of TFA (3 ml in 10 ml of benzene). Yields are as follows: **3a** (148 mg, 50%), **3b** (110 mg, 36%), **3c** (133 mg, 45%), **4a** (96 mg, 34%), **4b** (65 mg, 20%), **4c** (68 mg, 25%), **2a** (10 mg, 3%), **2b** (18 mg, 6%), **2c** (9 mg, 3%). Other minor unidentified spots were extracted from the reactions.

Crystal structure of 6,6'-bis-(1-oxide-1,2,6,8a-tetrahydroquinoline)ylidene (3a)

Crystals are dark violet prisms. Lattice constants were determined repeatedly rectifying on the diffractometer the values of $(\vartheta, \chi, \varphi)_{hkl}$ angles by thirty reflections to have the maximum of the peak when the angles were moving by 0.001° as a maximum.

Crystal data - $C_{42}H_{30}N_2O_2$, $M = 594.7$; monoclinic, $a = 20.547(4)$, $b = 14.381(3)$, $c = 11.097(2)$ Å; $\beta = 110.5(1)^\circ$, $U = 3071.4(23)$ Å³; $Z = 4$, $D_c = 1.29$ g cm⁻³; Cu-K α radiation, $\lambda = 1.5418$ Å, $\mu = 5.8$ cm⁻¹. Space group $C2/c$ (C_{2h}^6 , No. 15) from structure determination.

Intensity data were collected at 295 K on a Siemens AED single-crystal diffractometer in the range $3^\circ \leq 2\theta \leq 70^\circ$ using Ni-filtered Cu-K α radiation. The angles for every reflection were determined on the basis of the orientation matrix and the outline of the diffraction peak was collected in the ϑ - 2ϑ step scanning mode using a scan width from $(\vartheta - 0.60)^\circ$ to $(\vartheta + 0.60 + \Delta\lambda/\lambda \tan \vartheta)^\circ$. The intensities I_{hkl} were determined by analyzing the reflection profiles.³² 2917 Independent reflections were measured, 1072 of which were used in the crystal analysis having $I_{hkl} > 2\sigma(I_{hkl})$, $\sigma(I_{hkl})$ being based on statistic counting. One standard reflection, measured every 50 collected reflections to monitor crystal decomposition and instrumental linearity, showed no significant variation. Intensities I_{hkl} were corrected for Lorentz and polarization effects. The dimensions of the crystal were 0.05, 0.09, 0.24 mm. No absorption corrections were applied.

Structure analysis and refinement - The structure was solved by direct methods by use of the SHELXS86 program³² and refined by SHELX76³³ with cycles of full-matrix anisotropic least-squares (hydrogen atoms isotropically) up to $R = 0.034$, $R_w = 0.036$; $\sum w(F_o - F_c)^2$ minimized with $w = 0.1898/[\sigma^2(F_o) + 0.0033F_o^2]$. Atomic scattering factors from International Tables for X-ray Crystallography.³⁴

Crystal structure of 2,3-diphenylquinoline (4a)

Crystals are pale pink prisms. Cell parameters were derived as before.

Crystal data. $C_{21}H_{15}N$, $M = 281.4$. Monoclinic, $a = 8.734(2)$, $b = 16.674(3)$, $c = 11.112(3)$ Å; $\beta = 104.1(1)^\circ$, $U = 1569.5(9)$ Å³; $Z = 4$, $D_c = 1.19$ g cm⁻³; Cu-K α radiation, $\lambda = 1.5418$ Å, $\mu = 5.0$ cm⁻¹. Space group $P2_1/n$ (C_{2h}^6 , No. 15) from systematic absences.

Intensity data are collected as before. Of 3249 measured reflections ($-10 \leq h \leq 10$, $0 \leq k \leq 20$, $0 \leq l \leq 13$), 1759 symmetry independent were used in the crystal analysis. The dimensions of the crystal were 0.43, 0.10, 0.29 mm. No absorption corrections were applied.

Structure analysis and refinement. The structure was solved as before and refined by cycles of full-matrix anisotropic least-squares (hydrogen atoms isotropically) up to $R = 0.039$, $R_w = 0.044$. The weighting function was of the form $1/w = \sigma^2(F_o) + 0.0033F_o^2$.

All the hydrogen atoms were located in the difference-Fourier map. Atomic scattering factors from International Tables for X-ray Crystallography.³⁵

All the calculations were carried out on the GOULD 6040 POWERNODE computer of the Centro di Studio per la Strutturistica Diffraattometrica del C.N.R. di Parma and the Cambridge Crystallographic Data Files were used for bibliographic searches through the Servizio Italiano di Diffusione Dati Cristallografici di Parma.

REFERENCES

1. Swartz, H. M. in *Adv. Magn. Reson. Imaging*; Flig, E. Ed.; Ablex Publishing: Norwood, N.J., 1989.
2. Chignell, C. F. in *Spin Labelling II, Theory and Application*; Berliner, L. J. Ed.; Acad. Press: N. Y. London, 1979; p. 223.

3. Mitchell, J. B.; Samuni, A.; Krishna, N. C.; DeGraff, W. G.; Ahn, N. S.; Samuni, U.; Russo, A. *Biochem.* **1990**, 2802.
4. McConnell, H. N. in *Spin Labelling, Theory and Application*: Berliner, L. J. Ed.; Acad. Press: N. Y. London, 1976; p. 525.
5. Morrisett, J. D. *ibid.*, p. 274.
6. Swartz, H. M.; Pals, M. A. in *Handbook of Biomedicine of Free Radicals and Antioxidants*: Miguel, J.; Weber, H.; Quintanilha A. Eds.; CRC Press: Boca Raton, FL, 1989; Vol. 3, p.141.
7. Jansen, E. G.; Hair, D. L.; Coulter, G. A.; Stronks, H. J.; Krygsman, P. H.; Towner, R. A.; Hilborn, J. W. *J. Org. Chem.* **1989**, 54, 2915.
8. Ueno, I.; Kohno, M.; Mitsuta, K.; Mizuta, Y.; Kanegasaki, S. *J. Biochem.* **1989**, 105, 905; Mottley, C.; Mason, R. P. in *Biological Magnetic Resonance*: Berliner, L. J.; Reuben, J. Eds.; Plenum Press: N. Y. London, 1989; Vol. 8, p. 489.
9. Miller, W. G. in *Spin Labelling II, Theory and Application*: Berliner, L. J. Ed.; Acad. Press: N. Y. London, 1979; p. 173; Rånby, B.; Rabek, J. F. *ESR Spectroscopy in Polymer Research*: Springer Verlag: Berlin N. Y., 1977.
10. Rosantzev, E. G. *Free Nitroxyl Radicals*, Ulrich, H. Ed.; Plenum Press: N. Y. London, 1970.
11. Volodorsky, L. B. *Imidazoline Nitroxide*: Miguel, J.; Weber, H.; Quintanilha A. Eds.; CRC Press: Boca Raton, FL, 1988; Vol. I and III.
12. Berti, C.; Colonna, M.; Greci, L.; Marchetti, L. *Tetrahedron* **1976**, 32, 2147.
13. Antosiewicz, J.; Bertoli, E.; Damiani, E.; Greci, L.; Popinigis, J.; Przybylski, S.; Tanfani, F.; Wozniak, M. *Free Radical Biology and Medicine*, accepted for publication, 1993.
14. Carloni, P.; Greci, L.; Stipa, P.; Rizzoli, C.; Sgarabotto, P.; Ugozzoli, F. *Polymer Deg. and Stab.* **1993**, 39, 73; Alberti, A.; Carloni, P.; Greci, L.; Stipa, P.; Neri, C. *ibid.*, **1993**, 39, 215.
15. Cardellini, L.; Carloni, P.; Greci, L.; Stipa, P.; Faucitano, A. *Gazz. Chim. Ital.* **1989**, 119, 621, and references reported therein.
16. Carloni, P.; Greci, L.; Stipa, P.; Eberson, L. *J. Org. Chem.* **1991**, 56, 4733.
17. Lenner, M. and Lindgren, O. *Acta Cryst.* **1976**, B32, 1903; Folting, K. and L. Meritt, L. Jr. *Acta Cryst.* **1977**, B33, 3540; Gdaniec, M.; Jaskolski, M.; Kosturkiewicz, Z. *Acta Cryst.* **1977**, B33, 3558; Levina, O. I.; Rau, V. G.; Potekhin, K. A.; Kurkutova, E. N.; Pakhomov, I. G.; Struchkov, Yu. T. *Cryst. Struct. Comm.* **1982**, 11, 1899; Das, K.; Sinha, U. C.; Talwar, S. S.; Kamath, M. B.; Bohra, R. *Acta Cryst.* **1990**, C46, 2126.
18. Benassi, R.; Taddei, F.; Greci, L.; Marchetti, L.; Andreotti, G. D.; Bocelli, G.; Sgarabotto, P. *J. Chem. Soc. Perkin Trans. 2* **1980**, 786.
19. Pauling, L. *The Nature of the Chemical Bond*: Cornell University Press, 1960.
20. Rizzoli, C.; Sgarabotto, P.; Ugozzoli, F.; Carloni, P.; Damiani, E.; Greci, L.; Stipa, P. *J. Het. Chem.*, in press.
21. Rosantzev, E. G.; Sholle, V. D. *Synthesis* **1971**, 401.
22. Forrester, A. R.; Hay, J. M.; Thomson, R. H. *Organic Chemistry of Stable Free Radicals*: Academic Press: London, 1968; p. 225.
23. Greci, L. *Tetrahedron* **1983**, 39, 667.
24. Aurich, H. G. *Nitroxides in Amino, Nitroso and Nitro Compounds and their Derivatives, Part. 1*: Patai, S.; Wiley, J. and Sons, Eds.; N. Y., 1982; p. 598.
25. Aurich, H. G. *Pure and Appl. Chem. (2)* **1990**, 62, 183.
26. March, J.; *Advanced Organic Chemistry*: Wiley, J. and Sons, Eds.; N. Y., 1985; p. 950.
27. Eberson, L.; Radner, F. *J. Chem. Soc. Chem. Comm.* **1991**, 1233.
28. Carloni, P.; Eberson, L.; Greci, L.; Stipa, P.; Tosi, G. *J. Chem. Soc. Perkin Trans. 2* **1991**, 1779.
29. Effemberger, F.; Stohrer, W.; Mack, K.; Reisinger, F.; Seufert, W.; Kramer, H. E. A.; Föll, R.; Vogelmann, E. *J. Am. Chem. Soc.* **1990**, 112, 4849.
30. March, J.; *Advanced Organic Chemistry*: Wiley, J. and Sons, Eds.; N. Y., 1985; p. 331.
31. Monti, L.; Franchi, G. *Gazz. Chim. Ital.* **1951**, 81, 544.
32. Lehmann, M. S.; Larsen, F. K. *Acta Cryst.* **1974**, A30, 580.
33. Sheldrick, G. M. *SHELXS86 - Program for the Solution of Crystal Structures*: University of Göttingen, Germany, 1986.
34. Sheldrick, G. M. *SHELX76 - System of Computer programs for Crystal Structure Determination*: University of Cambridge, 1976.
35. *International Tables for X-ray Crystallography*: Kynoch Press: Birmingham, England, 1974; Vol. IV.