

Synthesis and Reactivity of Two *cis*-PtH₂-(Diphosphine) Complexes (Diphosphine = 1,3-Bis(diphenylphosphino)propane and 1,4-Bis(diphenylphosphino)butane)

ALBERTO SCRIVANTI*

Centro di Chimica Metallorganica del CNR, Via Marzolo 9, 35100 Padua, Italy


RENZO CAMPOSTRINI and GIOVANNI CARTURAN

Dipartimento di Ingegneria, Università di Trento, Mesiano, Trento, Italy


(Received June 19, 1987; revised August 27, 1987)

Square planar Pt(II) complexes holding two hydride ligands in the *cis* position are quite rare. Since the first report [1] a few other examples have been found for Pt(II) species stabilized either by bulky chelating diphosphines [2, 3] or basic monophosphines [4, 5].

The interest in these complexes stems from facets ranging from catalysis [6] and mechanisms of fundamental organometallic reactions [7], to intermediates in the synthesis of Pt–Pt dimeric complexes [2]. More recently, Clark [8] suggested a *cis*-dihydride $\leftrightarrow \eta^2$ -dihydrogen exchange to account for the dynamic behaviour of a series of *cis*-PtH₂-(diphosphine) complexes. Besides the intrinsic interest, such a type of exchange could be of paramount importance for the understanding of H₂ activation on metallic catalysts. Moreover, the Horiuti–Polanyi hydrogenation mechanism [9], where H₂ activation is claimed to be rate determining, has been recently reconsidered in the light of experimental data, which suggest the facile reaction [10, 11]:



This has a non-casual counterpart in the well-documented reaction [2–4, 8]:



Our interest in these topics [12, 13] spurred us to find new examples of *cis*-dihydride Pt(II) complexes and to study their reactivity and stability in the presence of H₂, ethylene and HCl.

Experimental

In the reactions oxygen was excluded by the standard techniques used for manipulation of air-sensitive compounds.

*Author to whom correspondence should be addressed.

¹H and ³¹P NMR spectra were recorded on a Varian FT 80A spectrometer operating at 79.542 and 32.203 MHz, respectively; IR spectra were registered on a Perkin-Elmer 983 spectrometer. Gas chromatographic/mass spectrometric analysis was carried out on a Finnigam-MAT 700 ion trap detector interfaced to a C. Erba Fractovap 4160 gas chromatograph.

Complexes PtCl₂(diphosphine) were prepared by reacting PtCl₂(COD) (COD = 1,5-cyclooctadiene) with the appropriate diphosphine ligand (Strem Chemicals).

Preparation of *cis*-PtH₂(Ph₂P–(CH₂)₃–PPh₂) (2a)

2.00 g of PtCl₂(Ph₂P–(CH₂)₃–PPh₂) are suspended in a 1:1 mixture of CH₂Cl₂/EtOH (10 ml) under H₂ atmosphere. The mixture is stirred at –10 °C for a few minutes, and solid NaBH₄ (0.5 g) is added in small amounts over 30 min. The resulting off-white suspension is left stirring for an additional 2 h. After this time a mixture of 5 ml MeOH and 20 ml EtOH is added and stirring is continued for 30 min. The suspension is quickly filtered in air and the white precipitate washed with H₂O, EtOH and n-hexane, and dried under vacuum. Yield 1.32 g (73%). *Anal.* Calc.: C, 53.2; H, 4.6. Found: C, 52.8; H, 4.7%.

The same procedure is used in the synthesis of complex 2b.

Preparation of Pt(C₂H₄)(Ph₂P–(CH₂)₃–PPh₂) (3a)

A 1:1 mixture of CH₂Cl₂/EtOH (5 ml) is saturated with ethylene at –78 °C. 0.5 g of complex 2a are added to yield a white suspension. The mixture is allowed to reach room temperature under ethylene atmosphere and stirred for 1 h. The white suspension is filtered, washed with n-hexane and dried *in vacuo*. Yield 0.3 g (60%). *Anal.* Calc. C, 54.8; H, 4.7. Found: C, 54.5; H, 4.7%.

Complex 3b is obtained in the same way.

Reaction of Complex 3a with HCl

0.32 g (0.5 mmol) of complex 3a are dissolved in 10 ml of CH₂Cl₂ under ethylene at 0 °C, and treated with the stoichiometric amount of dry HCl in diethyl-ether. Dilution of the resulting solution with diethyl-ether affords complex 4a as a white powder. Yield 0.25 g (75%).

The reaction of complex 3b with HCl is carried out in the same way.

Results and Discussion


The dichloro complexes *cis*-PtCl₂(L–L) (1a: L–L = 1,3-bis(diphenylphosphino)propane = dppp;

TABLE I. IR and NMR Data for Complexes *cis*-PtH₂(L-L) and Pt(C₂H₄)(L-L)^a

Complex	δH	<i>J</i> (P-H) _{cis}	<i>J</i> (P-H) _{trans}	<i>J</i> (Pt-H)	δP	<i>J</i> (Pt-P)	ν(Pt-H) (cm ⁻¹)
2a PtH ₂ (dppp) ^b	-2.21		152 ^c	1080	8.6	1945	2014
2b PtH ₂ (dppb) ^d	-2.91	± 19	± 170	1076	22.7	2054	2006
3a Pt(C ₂ H ₄)(dppp)	2.02 ^e			61	13.6	2711	
3b Pt(C ₂ H ₄)(dppb)	1.89 ^e			62	25.3	2528	

^aIR spectra are in CH₂Cl₂; NMR spectra are in CD₂Cl₂; chemical shifts are in ppm; coupling constants in Hz; ³¹P shifts are quoted relative to 85% H₃PO₄. ^bAt -70 °C. ^c[*J*(P-H)_{cis} + *J*(P-H)_{trans}] (see text), *J*_{cis} and *J*_{trans} are opposite in sign. ^dAt -80 °C. ^eEthylene protons.

b: L-L = 1,4-bis(diphenylphosphino)butane = dppb) upon reaction with NaBH₄ under H₂ atmosphere afford the *cis*-dihydride complexes **2a** and **2b**, respectively:



The preparative yields are generally good (>70%). Complexes **2a** and **2b** are recovered from the reaction mixture as white powders which can be safely handled in air for a short time; solutions of these species are air-sensitive but stable, when stored under H₂ atmosphere. The solutions are either colourless or pale-yellow, and turn dark-orange under vacuum or under the action of a nitrogen stream, which irreversibly leads to untractable products.


Complexes **2a** and **2b** have been characterized by analytical and spectroscopic data (IR, ¹H and ³¹P NMR).

The infrared spectra, registered in CH₂Cl₂ solution, show a broad absorption at ~2010 cm⁻¹ (see Table I) assigned to ν(Pt-H): this result agrees with the literature data which point out the difficulty in observing and assigning symmetrical and antisymmetrical Pt-H stretching vibrations in the case of *cis*-PtH₂(L₂) species [1, 2].

The ³¹P{¹H} NMR spectra of complexes **2a** and **2b** consist of a singlet flanked by ¹⁹⁵Pt satellites as expected from the magnetic equivalence of the two P atoms, under ¹H broad band decoupling conditions (the relevant data are reported in Table I); ¹*J*(Pt-P) values are in keeping with literature data.


The AA'XX' spin system of a *cis*-PtH₂(L-L) species should give rise to a pattern of ten separate lines in the hydride region of the ¹H NMR spectrum (disregarding ¹⁹⁵Pt satellites). In this spectral region complex **2a** shows a deceptively simple AA'XX' pattern which consists of only six separate lines flanked by ¹⁹⁵Pt satellites. A six line pattern for a AA'XX' spin system should be obtained if either *J*_{AA}, or *J*_{XX}, are ~0 Hz; however in our case, computer simulation with these parameters leads to poor agreement between the intensity of the calculated

lines and that of the observed ones (attempts to include possible couplings with the diphosphine protons were not made). It is likely that lower intensity lines are lost in the background and/or masked by some of the more intense ones. The spectral data obtained enable us to assign only |*J*(P-H)_{cis} + *J*(P-H)_{trans}| and these results are reported in Table I. The ¹H NMR spectrum of complex **2a** is not affected by temperature, while that of complex **2b** shows a dynamic behaviour. The low temperature limiting spectrum is obtained at -80 °C and shows only four equal-intensity lines for the central part of the hydride region. This pattern is consistent with an AA'XX' spin system in which both *J*_{AA}, and *J*_{XX}, are ~0 Hz, and allows straightforward assignment of *J*(P-H)_{cis} and *J*(P-H)_{trans} (see Table I). Increasing the sample temperature to ca. 20 °C leads to the broadening of the external lines of the quartet, whereas the internal ones remain unchanged. In the meanwhile no variation is observed in the ³¹P NMR spectrum, where the signals remain sharp and the coupling between ³¹P and ¹⁹⁵Pt is retained. This overall behaviour is very similar to that observed by Clark [8] and is consistent with an intramolecular process [14] leading to mutual exchange of the two hydrido ligands:



By comparing our ¹H NMR data with those in the literature it may be concluded that a deceptively simple AA'XX' spin system is a commonly encountered feature for this type of complex which mostly shows only four lines in the hydrido region. Other common features are the following: the hydrides resonate at relatively low fields (they cover the range 0.65 to -4.00 ppm); ²*J*(H-P)_{trans} are ~170 Hz, whereas ²*J*(H-P)_{cis} are ca. one order of magnitude smaller; ¹*J*(H-Pt) are >1000 Hz significantly larger than those observed for platinum dihydrides having *trans* geometry.

Both complexes **2a** and **2b** react with ethylene according to reaction (3):



This reaction does not lead to ethylene hydrogenation, as confirmed by GC-MS analysis. Complexes **3a** and **3b** are isolated as white powders, they are air-stable and the relevant spectroscopic data are reported in Table I.

Owing to our interest in the olefin hydroformylation with mono-hydrido Pt(II) species [15], we thought it interesting to study the reaction of complexes **2a** and **2b** with a stoichiometric amount of dry HCl, in order to obtain new *cis*-PtHCl(L-L) compounds [1]. However, all our attempts failed, since by reacting complexes **2a** and **2b** with a 1:1 stoichiometric amount of dry HCl we obtained a 50:50 mixture of the complexes **1a,b** and **2a,b**. The dichloro complexes **1a** and **1b** are the only products obtained by reacting complexes **2a** and **2b** with a two-fold stoichiometric amount of HCl.

Likewise, mono-hydrido-chloro complexes cannot be obtained by reacting complexes **3a** and **3b** with HCl: this reaction affords in good yield (>70%) the alkyl complexes *cis*-PtCl(C₂H₅)(L–L) (**4a,b**) which have been identified by comparing their spectroscopic properties with those of authentic samples [16].

References

- 1 C. J. Moulton and B. L. Shaw, *J. Chem. Soc., Chem. Commun.*, 365 (1976).
- 2 T. Yoshida, T. Yamagata, T. H. Tulip, J. A. Ibers and S. Otsuka, *J. Am. Chem. Soc.*, 100, 2063 (1978).
- 3 L. M. Venanzi, private communication; H. Rimml, *Ph.D. Thesis*, Diss. ETH No. 7562, Zurich, Switzerland, 1984.
- 4 R. S. Paonessa and W. C. Trogler, *J. Am. Chem. Soc.*, 104, 1138 (1982).
- 5 D. L. Packett, C. M. Jensen, R. L. Cowan, C. E. Strouse and W. C. Trogler, *Inorg. Chem.*, 24, 3578 (1985).
- 6 B. R. James, *Adv. Organomet. Chem.*, 17, 319 (1979); P. J. Brothers, *Prog. Inorg. Chem.*, 28, 1 (1981).
- 7 J. J. Low and W. A. Goddard, III, *Organometallics*, 5, 609 (1986); *J. Am. Chem. Soc.*, 106, 6928 (1984).
- 8 H. C. Clark and M. J. Hampden Smith, *J. Am. Chem. Soc.*, 108, 3829 (1986).
- 9 I. Koriuti and M. Polanyi, *Trans. Faraday Soc.*, 30, 1164 (1934).
- 10 S. J. Thomsom and G. Webb, *J. Chem. Soc., Chem. Commun.*, 526 (1976).
- 11 F. Zaera and G. A. Somorjai, *J. Am. Chem. Soc.*, 106, 2288 (1984).
- 12 R. Bertani, G. Carturan and A. Scrivanti, *Angew. Chem., Int. Ed. Eng.*, 22, 246 (1983).
- 13 G. Carturan and R. Campostrini, in Y. U. Yermakov and V. Likholobov (eds.), 'Homogeneous and Heterogeneous Catalysis', VNU Science Press, Utrecht, 1986, p. 923; G. Carturan, A. Scrivanti and M. Graziani, *J. Mol. Catal.*, 5, 139 (1979).
- 14 H. Azizian, K. R. Dixon, C. Eaborn, A. Pidcock, N. M. Shuhaib and J. Vinaixa, *J. Chem. Soc., Chem. Commun.*, 1020 (1982).
- 15 A. Scrivanti, A. Berton, L. Toniolo and C. Botteghi, *J. Organomet. Chem.*, 314, 369 (1986).
- 16 D. A. Slack and M. C. Baird, *Inorg. Chim. Acta*, 24, 277 (1977).