Radical Substitution of 2-Deoxy-2-lodo Pyranosides using 2,2,6,6-Tetramethylpiperidinoxy (TEMPO) Free Radical

Anthony G. M. Barrett* and David J. Rys

Department of Chemistry, Imperial College of Science, Technology and Medicine, London, UK SW7 2AY

Methyl 2-hydroxy- α -D-pyranosides were formed *via* photolytic radical substitution of methyl 2-deoxy-2-iodo- α -D-pyranosides with 2,2,6,6-tetramethylpiperidinoxy (TEMPO) free radical followed by zinc reduction.

Recently, during a synthesis of sucrose, we encountered difficulties effecting the S_N^2 substitution of iodide 1 by oxygen-centred nucleophiles.¹ However, radical-mediated substitution (Scheme 1)² under photolytic conditions with TEMPO free radical 4 provided hydroxylamine 2. Global deprotection *via* dissolving metal reduction gave sucrose which was isolated as the octacetate 3. We speculated that this method would also be applicable to iodoethers derived from glycosides, *via* epoxidation or iodoetherification *etc.*,³ and this methodology has recently been applied by Friesen and Danishefsky³ in oligosaccharide assembly. Herein, we report a method for the radical substitution of 2-deoxy-2-iodo- α -D-pyranosides with TEMPO to provide glycosides.

Methyl 2-deoxy-2-iodo- α -D-pyranosides (5,4 7 and 9) were synthesised from 3,4,6-tri-O-benzyl-D-glucal,⁵ -allal,⁶ and -galactal7 respectively via oxidative coupling with methanol {MeOH, [Ag(sym-collidine)₂]ClO₄, I₂, 4 Å molecules sieves, $CH_2Cl_2, 0 \circ C$.³ Subsequent photolysis [>200 nm, (Bu₃Sn)₂, 40 °C] in the presence of TEMPO (2.5 equiv.) afforded the hydroxylamine derivatives 6, 8 and 10 (Scheme 2). † Despite proceeding through a radical manifold, moderate diastereoselectivity was observed for each substrate. For example, the mannopyranoside 5 yielded a mixture of diastereoisomers‡ favouring the mannopyranoside 6a over the glucopyranoside **6b** (2.0:1). This modest selectivity is consistent with approach of the bulky TEMPO reagent from the less sterically congested face8 of the radical intermediate. Radical substitution of the altropyranoside 7 favoured the corresponding altropyranoside 8a over the allopyranoside 8b (4.5:1). In contrast, photolysis of iodide 9 in the presence of TEMPO (4) and hexabutyldistannane gave rise predominately to the cisglycoside (10a: 10b = 2.2: 1). Again both these selectivities are consistent with steric approach control. Unfortunately, efforts to increase the diastereoselectivities of these radical substitution reactions by lowering the temperature served only to slow the process. Dissolving metal reduction (activated Zn dust, 10% HOAc-EtOH-EtOAc 3:1:1, sonication) of 6a, 8a and 10a resulted in cleavage of the hydroxylamine nitrogen-oxygen bond affording the hydroxy pyranosides 119 (81%), 12 (78%) and 1310 (93%) respectively.†§

In conclusion a mild method for stereoselective radical substitution of iodine with oxygen functionality has been demonstrated on several 2-deoxy-2-iodo pyranosides. The diastereoselectivity of reaction is consistent with steric approach control of the TEMPO free radical. Recently, Nakamura *et al.*¹¹ have described an alternative method for the radical substitution of alkyl iodides using tributylstannane and oxygen.

We thank Glaxo Group Reserch Ltd. for the most generous endowment (to A. G. M. B.), the Wolfson Foundation for establishing the Wolfson Centre for Organic Chemistry in Medical Science at Imperial College, Merck Sharpe & Dohme Research Laboratories, Wyeth-Ayerst and the US Department of Education under the Graduate Assistance in Areas of National Need Program for Fellowship support (to D. J. R.), the National Institutes of Health (GM-40949) for support when this work started in the USA and the James Black Foundation, Parke Davis, The Proctor and Gamble Com-



Scheme 2 Reagents: i, TEMPO, (Bu₃Sn)₂, PhH, hv





pany, Quest International, Rhône-Poulenc Rorer Ltd., Roche Products Ltd., Rohm and Haas Company, G. D. Searle & Company and ZENECA Corporate Research and Technology for generous unrestricted support of our program.

Received, 24th January 1994; Com. 4/00425F

Footnotes

[†] Pyranosides **6a**, **7**, **8a**, **9**, **10a** and **12** were fully characterised by spectroscopic data and microanalyses. The minor diastereoisomers **6b** and **10b** were characterised by ¹H NMR, ¹³C NMR and MS only. The minor diastereoisomer **8b** could not be completely purified and its structural assignment is tentative.

[‡] The preparation of hydroxylamines (**6a/6b**) is representative: iodide **5** (51.6 mg, 90.0 μ mol) was dissolved in benzene (1.5 ml) in a quartz reaction vessel. TEMPO 4 (37.4 mg, 0.239 mmol) and (Bu₃Sn)₂ (67 μ l, 0.175 mmol) were added and the mixture was irradiated (>200 nm) for 2 h at 40 °C with a 450 W mercury immersion lamp equipped with a quartz cooling jacket. EtOAc (5 ml) was added and the solution washed with 10% aqueous KF (2 × 10 ml). The aqueous layers were combined and extracted with EtOAc (3 × 10 ml), and the extracts dried (Na₂SO₄), evaporated, and chromatographed on silica (eluant hexanes–Et₂O 3: 1) to yield **6a** (25 mg, 46%) and **6b** (12.6 mg, 23%) both as oils.

§ The preparation of pyranoside 11 is representative: to a solution of **6a** (42.6 mg, 71.0 μ mol) in 10% aqueous AcOH-EtOH-EtOAc (7 ml, 3:1:1) was added activated zinc dust (220 mg, 3.36 mmol). The mixture was sonicated for 14 h, extracted with EtOAc (3 × 5 ml),

J. CHEM. SOC., CHEM. COMMUN., 1994

dried Na_2SO_4) and evaporated. Chromatography on silica (eluant hexanes-EtOAc 7:3) yielded 11 (27 mg, 81%) as a clear oil.

References

- 1 A. G. M. Barrett, B. C. B. Bezuidenhoudt and L. M. Melcher, J. Org. Chem., 1990, 55, 5196.
- R. J. Kenney, W. D. Jones and R. G. Bergman, J. Am. Chem. Soc., 1978, 100, 7902; A. R. Howell and G. Pattenden, J. Chem. Soc., Chem. Commun., 1990, 2, 103.
 R. U. Lemieux and B. Fraser-Reid, Can. J. Chem., 1964, 42, 539,
- R. U. Lemieux and B. Fraser-Reid, Can. J. Chem., 1964, 42, 539, 547; R. U. Lemieux and B. Fraser-Reid, Can. J. Chem., 1965, 43, 1460; R. U. Lemieux and A. R. Morgan, Can. J. Chem., 1965, 43, 2190; R. W. Friesen and S. J. Danishefsky, J. Am. Chem. Soc., 1989, 111, 6656.
- 4 P. J. Garegg and B. J. Samuelssen, J. Chem. Soc., Perkin Trans. 1, 1980, 12, 2866.
- 5 W. Roth and W. Pigman, Methods in Carbohydrate Chemistry, vol. VI, p. 405.
- 6 D. C. Baker, D. Hartman and C. G. Tindall, Jr., Carbohydr. Res., 1972, 24, 192.
- 7 F. Shafizadeh, Methods in Carbohydrate Chemsitry, vol. VI, p. 409.
- 8 B. Giese, Radicals in Organic Synthesis: Formation of Carbon-Carbon Bonds, Pergamon, Oxford, 1986, pp. 62-63.
- 9 N. E. Franks and R. Montgomery, Carbohydr. Res., 1968, 6, 286.
- 10 P. A. J. Gorin, Carbohydr. Res., 1982, 101, 13.
- 11 E. Nakamura, T. Inubushi, S. Aoki and D. Machii, J. Am. Chem. Soc., 1991, 113, 8980.