

A simple route to chiral phosphinous acid–boranes†

Delphine Moraleda, David Gatineau, David Martin, Laurent Giordano and Gérard Buono*

Received (in Cambridge, UK) 19th February 2008, Accepted 26th March 2008

First published as an Advance Article on the web 22nd April 2008

DOI: 10.1039/b802817f

We report a simple one-pot synthesis of enantiomerically enriched alkyl- and arylphenylphosphinous acid–borane starting from readily available (*R_P*)-(–)-menthylhydrogenophenylphosphinate and organolithium reagents.

The availability of chiral organophosphorous compounds has allowed tremendous achievements in chemistry, in particular in the field of asymmetric metal catalysis.¹ However most of the popular chiral P-ligands, such as the ubiquitous BINAP, are based on chiral C-backbones. In comparison, the occurrence of P-stereogenic ligands in literature is lower because their synthesis often presents a challenge. As P-chiral compounds could not be found in the natural pool of chirality, they were initially obtained by the resolution of racemic mixtures by means of a chiral auxiliary. Nowadays, stereoselective syntheses are supplanting this methodology.² For instance, one of the most convenient and versatile approach to chlorophosphine–boranes or phosphinite–boranes, which are precursors of P-chiral tertiary phosphine–boranes,³ is the nucleophilic substitution of chiral 1,3,2-oxazaphospholidine–boranes with organolithium reagents. However, this method, which was initially described by Jugé *et al.*,⁴ suffers from limitations. For instance, the subsequent acidolysis (or methanolysis) step is limited to not sterically hindered substituents. As recently stated,^{2d} an overall methodology is still lacking.


Recently, Pietrusiewicz and Stankevici demonstrated the potentiality of chiral phosphinous acid–boranes as key intermediates for the preparation of various P-stereogenic compounds, such as bulky tertiary and secondary phosphine–boranes, phosphinite–boranes or boranatophosphinous–sulfinic anhydrides.⁵ Some enantiopure phosphinous acid–boranes are available, but only from racemic resolution.^{5e,f} Despite promising results, this methodology is at a very early stage of development. In particular, an efficient and versatile stereoselective route to chiral phosphinous acid–boranes remains to be designed. To our knowledge there are only two reports of stereoselective syntheses (Scheme 1), and none exceeds 73% of enantiomeric excess (ee).⁶ Herein we report a simple one-pot and stereoselective synthesis of phosphinous acid–boranes, with up to 99% ee.

The phosphinous acid–secondary phosphine oxide (SPO) tautomerism is well known.⁷ Usually, only the latter, which is the more stable, is observed.⁸ Curiously, despite the tauto-

meric equilibrium, SPOs do not react with boranes, under mild conditions. We knew from the report of Pietrusiewicz and co-workers^{5b} that the reaction could occur at room temperature when deprotonating the SPO. Recently, our interest in SPOs⁹ prompted us to design an enantioselective synthesis of such compounds from (*R_P*)-(–)-menthyl hydrogenophenylphosphinate **1**.^{9c} We felt that this approach could be adapted to the synthesis of phosphinous acid–boranes.

Initially, we examined the model reaction of enantiopure (*R_P*)-(–)-**1**¹⁰ with MeLi (2.5–3 equiv.) at –78 °C in THF. The first equivalent of organometallic reagents was required to abstract the proton carried by (*R_P*)-(–)-**1**, whereas the second equivalent was used for the substitution of menthyloxy group. At this stage, lithium phosphinate species (*S*)-**2a** was quenched with BH₃·SMe₂. The desired methylphenyl phosphinous acid–borane (*S*)-(–)-**3a** was obtained in 78% overall yield and 95% ee (Table 1, entry 1). Of note, when the reaction was carried out with MeMgBr instead of MeLi, (*S*)-(–)-**3a** was obtained with poorer yield (64%) and ee (86%).


Then, we briefly explored the scope of this first approach (method A) using various organolithium reagents in similar conditions. The phosphinous acid–boranes **3a–g** were obtained in good yields (Table 1). Good enantioselectivities (91–98% ee) were observed with aryllithium reagents (entries 4–6) and *n*-butyllithium (89% ee, entry 2). Conversely, **3g** and **3c** were obtained with modest 72 and 60% ee, respectively (entries 7 and 3). The latter case was particularly enlightening. Indeed, we knew that the first step of the procedure afforded **2c** with at least 84% ee.^{9c} In addition, ee's remained unchanged when stirring **2c** at room temperature for 3 days, suggesting that this compound did not racemize under our conditions. Thus, a significant loss of enantiopurity occurred upon the addition of BH₃·SMe₂. This statement is in agreement with the recent report of Stankevici *et al.*, who noticed that the deprotonation of resolved enantiopure *tert*-butylphenylphosphine oxide, followed by the addition of BH₃, afforded **3c** with only 74% ee.^{5a}



Scheme 1 Previously reported stereoselective routes to chiral phosphinous acid–boranes.

Université Aix-Marseille, Institut des Sciences Moléculaires de Marseille, ECM, CNRS, UMR 6263, Av. Escadrille Normandie Niemen, Marseille Cedex 20, 13397. E-mail: gerard.buono@univ-cezanne.fr

† Electronic supplementary information (ESI) available: Experimental procedures and spectroscopic analyses. See DOI: 10.1039/b802817f

Table 1 One-pot synthesis of phosphinous acid-boranes **3a–g** (method A)


Entry	R	Product	Yield ^a (%)	ee ^b (%)
1	Me	<i>S</i> -(–)- 3a	78	95
2	<i>n</i> -Bu	(–)- 3b	84	89
3	<i>t</i> -Bu	<i>S</i> -(–)- 3c	82	60
4	2-MeC ₆ H ₄	(+)- 3d	93	98
5	2-PhC ₆ H ₄	(–)- 3e	84	93
6	1-Naphthyl	<i>R</i> -(–)- 3f	88	91 ^c
7	1-Furyl	(+)- 3g	92	72


^a Yields after purification. ^b Enantiomeric excesses were determined by HPLC analysis on Chiralpak AS-H ($\lambda = 254$ nm; 1 mL min^{–1}; eluent: hexane-*i*-PrOH mixtures). ^c Enantiomeric excess was determined by HPLC analysis on Chiralpak OD-H ($\lambda = 254$ nm; 1 mL min^{–1}; eluent: hexane-EtOH) (8 : 2).

Unsatisfied with these results we decided to explore a new approach (method B). After the addition of (*R_P*)-(–)-**1** to organolithium reagents,¹¹ lithium phosphinate species **2a–g** were trapped by addition of Me₃SiCl. The resulting O-silylated product **4** was then protected using BH₃.SMe₂ and desilylated by treatment with aqueous HCl. All steps were conveniently monitored by ³¹P NMR. This one-pot procedure afforded the desired phosphinous acid-borane **3a–g** in 70–85% yields after purification. All results are summarised in Table 2.

Using this method almost all enantioselectivities have been improved (Table 2, entries 3, 5, 6 and 7). Of note, when the reaction was performed with *t*-BuLi, *S*-(–)-**3c** was obtained with 84% ee. This value, which corresponds to the enantioselectivity of the first step, seemed to indicate that the three last steps occurred without significant racemization. To verify this point, we synthesized *tert*-butylphenylphosphine oxide **5** by adding (*R_P*)-(–)-**1** to *tert*-butyllithium.^{9c} A single crystallisation afforded (*S*)-(–)-**5** with 99.6% ee.^{9b} Addition of Me₃SiCl and the subsequent addition of BH₃.SMe₂ afforded (*S*)-(–)-**3c** with >99% ee, as indicated by HPLC analysis (Scheme 2).


In conclusion, we reported simple diastereoselective one-pot syntheses of enantioenriched chiral phosphinous acid-boranes starting from cheap and readily available (*R_P*)-(–)-menthylhydrogenophenylphosphinate and organolithium reagents. Although direct trapping of the resulting lithium phosphinate species with BH₃.SMe₂ afforded the desired products with up to 98% ee, it generally resulted in partial racemization. Such loss could be avoided by using Me₃SiCl as trapping agent. This general approach affords an unprecedented versatile diastereoselective entry to chiral phosphinous acid-boranes, and therefore, to a broad range of P-chiral compounds.

We are grateful to CNRS and ANR project “SPOs Preli-gands” (Contract No. BLAN07-1_190839) for funding. D. G. thanks MENRT for the doctoral fellowship. We also thank Dr N. Vanthuyne (CNRS, UMR 6263) for HPLC analyses.

Table 2 One-pot synthesis of phosphinous acid-boranes **3a–g** (method B)


Entry	R	4a–g ($\delta^{31}\text{P}$ / ppm)	Product	Yield ^a (%)	ee ^b (%)
1	Me	100	<i>S</i> -(–)- 3a	78	95
2	<i>n</i> -Bu	106	(–)- 3b	70	89
3	<i>t</i> -Bu	115	<i>S</i> -(–)- 3c	70	84
4	2-Me(C ₆ H ₄)	97	(+)- 3d	75	97
5	2-Ph(C ₆ H ₄)	99	(–)- 3e	85	95
6	1-Naphthyl	92	<i>R</i> -(–)- 3f	75	99 ^c
7	1-Furyl	75	(+)- 3g	72	80

^a Yields after purification. ^b Enantiomeric excesses were determined by HPLC analysis on Chiralpak AS-H ($\lambda = 254$ nm; 1 mL min^{–1}; eluent: hexane-*i*-PrOH mixtures). ^c Enantiomeric excess was determined by HPLC analysis on Chiralpak OD-H ($\lambda = 254$ nm; 1 mL min^{–1}; eluent: hexane-EtOH) (8 : 2).

**Scheme 2** Two-step stereoselective synthesis of enantiopure **3c**.

Notes and references

- (a) *Comprehensive Asymmetric Catalysis*, ed. E. N. Jacobsen, A. Pfaltz and H. Yamamoto, Springer, Heidelberg, 1999; (b) *Catalytic Asymmetric Synthesis*, ed. I. Ojima, Wiley-VCH, New York, 2nd edn, 2000.
- (a) K. M. Pietrusiewicz and M. Zablocka, *Chem. Rev.*, 1994, **94**, 1375–1411; (b) T. Imamoto, K. V. L. Cr  py and K. Katagiri, *Tetrahedron: Asymmetry*, 2004, **15**, 2213–2218; (c) T. Imamoto, H. Tsuruta, Y. Wada, H. Masuda and K. Yamaguchi, *Tetrahedron Lett.*, 1995, **36**, 8271–8274; (d) A. Grabulosa, J. Granell and G. Muller, *Coord. Chem. Rev.*, 2007, **251**, 25–90.
- M. L. Clarke and J. M. J. Williams, in *A Practical Approach in Chemistry: Organophosphorus Reagents*, ed. P. J. Murphy, Oxford University Press, New York, 2004, pp. 35–41.
- (a) S. Jug  , M. Stephan, J. A. Laffitte and J. P. Genet, *Tetrahedron Lett.*, 1990, **31**, 6357–6360; (b) S. Jug  , M. Stephan, R. Merd  s, J. P. Genet and S. Halut-Desportes, *J. Chem. Soc., Chem. Commun.*, 1993, 531–533; (c) D. Moulin, C. Darcel and S. Jug  , *Tetrahedron: Asymmetry*, 1999, **10**, 4729–4743; (d) D. Moulin, S. Bago, C. Bauduin, C. Darcel and S. Jug  , *Tetrahedron: Asymmetry*, 2000, **11**, 3939–3956.
- (a) M. Stankevic and K. M. Pietrusiewicz, *Synlett*, 2003, 1012–1016; (b) M. Stankevic, G. Andrijewski and K. M. Pietrusiewicz, *Synlett*, 2004, 311–315; (c) M. Stankevic and K. M. Pietrusiewicz, *Synthesis*, 2005, 1279–1290; (d) M. Stankevic and K. M. Pietrusiewicz, *Curr. Org. Chem.*, 2005, **9**, 1883–1897; (e) M. Stankevic and K. M. Pietrusiewicz, *J. Org. Chem.*, 2007, **72**, 816–822; (f) M. Stankevic and K. M. Pietrusiewicz, *Tetrahedron: Asymmetry*, 2007, **18**, 552–556.

6. (a) J. Uziel, M. Stephan, E. B. Kaloun, J. P. Genêt and S. Jugé, *Bull. Soc. Chim. Fr.*, 1997, **134**, 379–389; (b) K. Nagata, S. Matsukawa and T. Imamoto, *J. Org. Chem.*, 2000, **65**, 4185–4188.
7. D. E. C. Corbridge, *Phosphorus, An Outline of its Chemistry, Biochemistry and Uses*, Elsevier Science BV, Amsterdam, 5th edn, 1995, p. 336.
8. A recently isolated stable phosphinous acid stands as an exception to this rule: (a) B. Hoge, S. Neufeind, S. Hettel, W. Wiebe and C. Thösen, *J. Organomet. Chem.*, 2005, **690**, 2382–2387; (b) B. Hoge, G. Placido, H. Willner and H. Oberhammer, *Chem.–Eur. J.*, 2006, **12**, 3567–3574, and references cited therein.
9. (a) J. Bigeault, L. Giordano and G. Buono, *Angew. Chem., Int. Ed.*, 2005, **44**, 4753–4757; (b) A. Leyris, D. Nuel, L. Giordano, M. Achard and G. Buono, *Tetrahedron Lett.*, 2005, **46**, 8677–8680; (c) A. Leyris, J. Bigeault, D. Nuel, L. Giordano and G. Buono, *Tetrahedron Lett.*, 2007, **48**, 5247–5250; (d) J. Bigeault, L. Giordano, I. de Riggi, Y. Gimbert and G. Buono, *Org. Lett.*, 2007, **9**, 3567–3570.
10. W. B. Farnham, R. K. Murray, Jr and K. Mislow, *J. Am. Chem. Soc.*, 1970, **92**, 5809–5810.
11. ‘Inverse’ addition of methyllithium or butyllithium on (*R_p*)-(–)-**1** was also performed: similar enantioselectivities were observed.