






$k_8$  have the values 0.44 and 0.56  $M^{-2} \text{ sec}^{-1}$  at 0°. <sup>20</sup> Again the ratio of  $k_7/k_8 = 0.79$  is significantly smaller than the statistically expected value of 4.0. Moreover, a comparison of the value  $k_7 + k_8 = 1.0 M^{-2} \text{ sec}^{-1}$  (0°) with the value  $k_9 = 9 M^{-1} \text{ sec}^{-1}$  (0°) for reaction 9<sup>21</sup> il-



lustrates once more the small effect of added chloride ion on the rate of inner-sphere reactions.

It is instructive to recast the chloride ion effects on the electron-transfer reactions considered above in the form of equilibrium constants involving transition states. Calculations have been previously carried out for acid dissociation constants<sup>22,23</sup> and can be generalized to many other formal reactions of transition states.<sup>24</sup> As an example, it can be seen that the proper combination of eq 10, 11, and 12 leads to eq 13 and a calculated value of  $Q_{13} = 5.8 \times 10^6 M^{-1}$ .



Calculated values of  $Q$  for the formal reactions of the transition states of interest in the present note are summarized in Table II. Although no new information is obtained by recasting the chloride ion effects in terms of the equilibrium constants listed in Table II, the approach provides a useful summary of these effects. For example, comparisons of eq A with B and C and of eq D with E and F clearly illustrate the enormous difference between the effect of chloride ion in bridging as compared to nonbridging positions. Reactions A and D feature the substitution of a bridging water by chloride ion, and are seen to be extremely favorable ( $Q > 10^5$ ). In contrast, reactions B, C, E, and F, which feature the substitution of a *cis* or *trans* nonbridging water by chloride ion, are relatively unfavorable ( $Q = 0.05$ – $0.1$ ). Moreover, the values  $Q = 1.0$  and  $0.8$  for reactions G and H, respectively, when compared with the statistically expected value of 4.0, clearly demonstrate the increased stability of the transition states containing *trans* chloride ions. Interestingly, an entirely similar observation regarding the relative stabi-

TABLE II

FORMAL EQUILIBRIUM CONSTANTS FOR SUBSTITUTION OF WATER BY CHLORIDE ION IN CHROMIUM-CONTAINING TRANSITION STATES

Reactions	$Q, M^{-1}{}^a$	Ref
A. $[(\text{NH}_3)_5\text{Co}(\text{OH}_2)\text{Cr}^{5+}]^\ddagger + \text{Cl}^- \rightleftharpoons [(\text{NH}_3)_5\text{CoClCr}^{4+}]^\ddagger + \text{H}_2\text{O}$	$5.8 \times 10^6$	<i>b, c, d</i>
B. $[(\text{NH}_3)_5\text{CoClCr}^{4+}]^\ddagger + \text{Cl}^- \rightleftharpoons [cis-(\text{NH}_3)_5\text{CoClCrCl}^{3+}]^\ddagger + \text{H}_2\text{O}$	0.1	This work
C. $[(\text{NH}_3)_5\text{CoClCr}^{4+}]^\ddagger + \text{Cl}^- \rightleftharpoons [trans-(\text{NH}_3)_5\text{CoClCrCl}^{3+}]^\ddagger + \text{H}_2\text{O}$	0.1	This work
D. $[(\text{H}_2\text{O})_5\text{Cr}(\text{OH}_2)\text{Cr}^{5+}]^\ddagger + \text{Cl}^- \rightleftharpoons [(\text{H}_2\text{O})_5\text{CrClCr}^{4+}]^\ddagger + \text{H}_2\text{O}$	$>10^5$	<i>e, f, g</i>
E. $[(\text{H}_2\text{O})_5\text{CrClCr}^{4+}]^\ddagger + \text{Cl}^- \rightleftharpoons [cis-(\text{H}_2\text{O})_5\text{CrClCrCl}^{3+}]^\ddagger + \text{H}_2\text{O}$	0.05	<i>f, h, i</i>
F. $[(\text{H}_2\text{O})_5\text{CrClCr}^{4+}]^\ddagger + \text{Cl}^- \rightleftharpoons [trans-(\text{H}_2\text{O})_5\text{CrClCrCl}^{3+}]^\ddagger + \text{H}_2\text{O}$	0.06	<i>f, h, i</i>
G. $[trans-(\text{NH}_3)_5\text{CoClCrCl}^{3+}]^\ddagger \rightleftharpoons [cis-(\text{NH}_3)_5\text{CoClCrCl}^{3+}]^\ddagger$	1.0	This work
H. $[trans-(\text{H}_2\text{O})_5\text{CrClCrCl}^{3+}]^\ddagger \rightleftharpoons [cis-(\text{H}_2\text{O})_5\text{CrClCrCl}^{3+}]^\ddagger$	0.8	<i>f, h, i</i>

<sup>a</sup> Values for reactions A–C and G at 25°. Values for reactions D–F and H at 0°. <sup>b</sup> A. M. Zwickel and H. Taube, *J. Am. Chem. Soc.*, **81**, 1288 (1959). <sup>c</sup> Reference 12. <sup>d</sup> D. A. Buckingham, I. I. Olsen, A. M. Sargeson, and H. Satrapa, *Inorg. Chem.*, **6**, 1027 (1967). <sup>e</sup> A. Anderson and N. A. Bonner, *J. Am. Chem. Soc.*, **76**, 3826 (1954). <sup>f</sup> Reference 21. <sup>g</sup> C. F. Hale and E. L. King, *J. Phys. Chem.*, **71**, 1779 (1967). <sup>h</sup> Reference 10. <sup>i</sup> Reference 14.

ties of *cis* and *trans* isomers was previously made for the  $\text{CrCl}_2^+$  system.<sup>13,14</sup>

CONTRIBUTION FROM THE OLIN RESEARCH CENTER, CHEMICALS GROUP, NEW HAVEN, CONNECTICUT 06504

## Reaction of Chlorine Monofluoride with Some Group VI Elements and Derivatives

BY JAMES J. PITTS AND ALBERT W. JACHE

Received January 26, 1968

As a fluorinating agent, chlorine monofluoride has been placed approximately below  $\text{F}_2$ ,  $\text{ClF}_3$ ,  $\text{BrF}_5$ , and  $\text{IF}_7$  in reactivity.<sup>1,2</sup> The available information for such an assignment is mostly qualitative. For example, relatively little is known about the reactivity of chlorine monofluoride with group VI elements. Sulfur is said to react slowly and without flame, but no products are specified.<sup>3</sup> With  $\text{SF}_4$  at 350°, chlorine monofluoride is said to react as a chlorofluorinating agent to yield  $\text{SF}_5\text{Cl}$ .<sup>4</sup>

Acting as a moderate fluorinating agent, chlorine monofluoride should have a number of attractive synthesis applications. This paper describes the investigation of reactions between chlorine monofluoride

(20) Espenson and Slocum<sup>10</sup> reported the values 0.22 and 0.28, respectively. However, they failed to recognize that the observed<sup>6</sup> rate constant of 0.50 needs to be multiplied by 2 to account for the fact that only half of the forward events in reactions 7 and 8 lead to chloride exchange. The reported<sup>10</sup> values of  $Q_{20}$  and  $Q_{21}$  also must be multiplied by 2 to correct for the statistical effect.

(21) D. L. Ball and E. L. King, *J. Am. Chem. Soc.*, **80**, 1091 (1958).

(22) J. H. Espenson, *Inorg. Chem.*, **4**, 1025 (1965).

(23) T. W. Newton, private communication to J. H. Espenson; quoted in ref. 22.

(24) A. Haim, *Inorg. Chem.*, **7**, 1475 (1968).

(1) H. C. Clark, *Chem. Rev.*, **58**, 869 (1958).

(2) H. Meinert, *Z. Chem.*, **7**, 41 (1967).

(3) H. S. Booth and J. T. Pinkston, Jr., *Chem. Rev.*, **41**, 421 (1947).

(4) F. Nyman and H. L. Roberts, *J. Chem. Soc.*, 3180 (1962).

and the group VI elements sulfur, selenium, molybdenum, tungsten, and several of their derivatives.

### Experimental Section

**Materials.**—Chlorine monofluoride was obtained from the Ozark-Mahoning Co. and was determined to be 99.4% pure by vapor-phase chromatography. Separation, using a column of Kel-F No. 1 oil on Fluoro-Pak support media in Teflon tubing, showed 0.4%  $\text{CF}_4$  (undoubtedly present in the starting fluorine) and 0.2%  $\text{Cl}_2$  as impurities. The presence of  $\text{CF}_4$  in the infrared spectrum of  $\text{ClF}$  was also confirmed by its strong sharp band at  $1283\text{ cm}^{-1}$ .<sup>5</sup>

The group VI elements and derivatives were obtained from a number of sources: tungsten (assay 99.9%), molybdenum (assay 99.7%), molybdenum disulfide, lead molybdate, and lead tungstate, K & K Laboratories, Inc.; selenium, molybdic anhydride (assay 99.5%), sodium molybdate, and sodium tungstate, Fisher Scientific Co.; tungstic anhydride, Fansteel Metallurgical Corp.; potassium molybdate, Amend Drug and Chemical Co., Inc.; sulfur (precipitated USP grade, assay 99.5%, Baker and Adamson label), Allied Chemical Corp.

All reagents were used as received with the exception of the dihydrated sodium molybdate and tungstate which were dried for several hours under vacuum at  $120^\circ$  to remove the water.

**Apparatus.**—The vacuum system used in this work consisted of a 0.25-in. o.d. copper tubing manifold equipped with Imperial Eastman Hi-Duty brass fittings and Hoke M327 Monel needle valves. Attached to the manifold were several Hoke high-pressure 4HSM150 Monel cylinders which served as the reaction vessel and for collection and storage of the gaseous products. The cylinders were fitted with the same type of Monel needle valves, the threaded joints being sealed with "Strip Teeze" brand Teflon pipe joint tape. The pressure within the vacuum system was recorded on an Ashcroft 1082AC test gauge which was calibrated to give a reading to within  $\pm 3\text{ mm}$ .

**Infrared Spectra.**—The infrared spectra of the gaseous products were obtained on a Perkin-Elmer Infracord spectrophotometer using an Isomet Corp. 5-cm Monel-body gas cell with  $\text{AgCl}$  windows sealed with Teflon gaskets. Extreme care was taken initially to pacify both the ir cell and the metal-transfer line with  $\text{ClF}$ . Failure to do so resulted in spectra showing  $\text{ClO}_2$ <sup>8</sup> and  $\text{ClO}_2\text{F}^7$  as impurities.

Identification (both qualitative and quantitative) of the gaseous reaction products was made by comparison to known spectra appearing in the literature and included the references cited here:  $\text{ClF}$ ,<sup>8</sup>  $\text{SF}_4$ ,<sup>9-10</sup>  $\text{SOF}_2$ ,<sup>10-11</sup>  $\text{SO}_2\text{ClF}$ ,<sup>10</sup>  $\text{SF}_6$ ,<sup>12,13</sup>  $\text{SOF}_4$ ,<sup>14</sup>  $\text{S}_2\text{F}_{10}$ ,<sup>15</sup>  $\text{SF}_5\text{Cl}$ ,<sup>16,17</sup>  $\text{SeF}_4$ ,<sup>18</sup>  $\text{SeF}_6$ ,<sup>19</sup>  $\text{WF}_6$ ,<sup>18,20</sup>  $\text{WF}_5\text{Cl}$ ,<sup>21</sup>  $\text{MoF}_6$ ,<sup>20</sup>  $\text{SiF}_4$ ,<sup>22</sup>  $\text{CF}_3\text{Cl}$ ,<sup>23,24</sup>  $\text{COF}_2$ ,<sup>25</sup> and  $\text{COCIF}$ .<sup>26</sup>

(5) P. J. H. Woltz and A. H. Nielsen, *J. Chem. Phys.*, **20**, 307 (1952).

(6) K. Hedberg, *ibid.*, **19**, 509 (1951).

(7) D. F. Smith, G. M. Begun, and W. H. Fletcher, *Spectrochim. Acta*, **20**, 1763 (1964).

(8) E. A. Jones, T. F. Parkinson, and T. G. Burke, *J. Chem. Phys.*, **18**, 235 (1950).

(9) R. E. Dodd, L. A. Woodward, and H. L. Roberts, *Trans. Faraday Soc.*, **52**, 1054 (1956).

(10) F. Seel and H. D. Göltz, *Z. Anorg. Allgem. Chem.*, **327**, 32 (1964).

(11) E. L. Pace and H. V. Samuelson, *J. Chem. Phys.*, **44**, 3682 (1966).

(12) R. T. Lagemann and E. A. Jones, *ibid.*, **19**, 534 (1951).

(13) J. Gaunt, *Trans. Faraday Soc.*, **49**, 1122 (1953).

(14) F. B. Dudley, G. H. Cady, and D. F. Eggers, Jr., *J. Am. Chem. Soc.*, **78**, 1553 (1956).

(15) R. E. Dodd, L. A. Woodward, and H. L. Roberts, *Trans. Faraday Soc.*, **53**, 1545 (1957).

(16) H. L. Roberts and N. H. Ray, *J. Chem. Soc.*, 665 (1960).

(17) L. H. Cross, H. L. Roberts, P. Goggin, and L. A. Woodward, *Trans. Faraday Soc.*, **56**, 945 (1960).

(18) E. E. Aynsley, R. E. Dodd, and R. Little, *Spectrochim. Acta*, **18**, 1005 (1962).

(19) T. G. Burke, *J. Chem. Phys.*, **25**, 791 (1956).

(20) T. G. Burke, D. F. Smith, and A. H. Nielsen, *ibid.*, **20**, 447 (1952).

(21) G. W. Fraser, M. Mercer, and R. D. Peacock, *J. Chem. Soc., A*, 1091 (1967).

(22) E. A. Jones, J. S. Kirby-Smith, P. J. H. Woltz, and A. H. Nielsen, *J. Chem. Phys.*, **19**, 242 (1951).

(23) H. W. Thompson and R. B. Temple, *J. Chem. Soc.*, 1422 (1948).

The presence of nonabsorbing chlorine was established by its diluent effect on the intensity of the absorption bands of the other gaseous products. Further confirmation was made by the 100 mm vapor pressure exhibited by liquid chlorine at  $-78^\circ$ .

**Reaction with Sulfur.**—A 150-ml Monel cylinder was first pacified with chlorine monofluoride and then charged with 1.0 g of pharmaceutical grade powdered sulfur. A Monel needle valve was attached and the cylinder evacuated on the vacuum line described above. Chlorine monofluoride was then introduced stepwise into the reaction cylinder from a storage cylinder until an equilibrium pressure of 1 atm was attained. The infrared spectrum of a 200-mm sample of the gaseous products revealed a number of compounds resulting from impurities in the sulfur. The main product proved to be  $\text{SF}_4$  exhibiting its characteristic bands at  $1744\text{ vvw}$ ,  $1281\text{ w}$ ,  $889\text{ vvs}$ ,  $738\text{ vvs}$ ,  $728\text{ vvs}$ , and  $715\text{ vs cm}^{-1}$  with lesser amounts of  $\text{SO}_2\text{ClF}$  at  $1475\text{ m}$  and  $1230\text{ m cm}^{-1}$ ,  $\text{SOF}_4$  at  $928\text{ s cm}^{-1}$ , and  $\text{SiF}_4$  at  $1030\text{ m cm}^{-1}$ .

The reaction system was then evacuated of all gaseous products; more chlorine monofluoride was introduced and again an infrared spectrum of the gaseous products was taken. This step was repeated until the system was "pacified" and the infrared spectrum was free from the effect of oxygen-containing impurities. At this point  $\text{SF}_4$  was the only product observed in the infrared spectrum. From band intensity *vs.* pressure measurements it was apparent that the  $\text{SF}_4$  was diluted with a non-absorbing gas by an approximate 1:2 ratio thus strongly suggesting chlorine from the stoichiometry. Further confirmation was made by the approximate 100 mm vapor pressure exhibited by the gaseous products when cooled to  $-78^\circ$  in a trichloroethylene-Dry Ice slush bath.

The products above were allowed to warm to room temperature and additional chlorine monofluoride was introduced to a pressure of 5 psi. The reaction cylinder was then suspended in a fluidized sand bed and heated at  $200^\circ$  for 2 hr. After cooling to room temperature, the infrared spectrum of a 250-mm sample of the gaseous products showed the presence of  $\text{SF}_6$  at  $940\text{ vvs}$ , possibly  $\text{S}_2\text{F}_{10}$  at  $827\text{ vvw}$ ,  $\text{CF}_4$  at  $1283\text{ s}$ ,  $\text{CF}_3\text{Cl}$  at  $1206\text{ vs}$  and  $1105\text{ vvw}$ ,  $\text{COCIF}$  at  $1095\text{ ms}$ , and  $\text{COF}_2$  at  $1248\text{ s cm}^{-1}$ . The latter four impurities are no doubt a result of interaction of the Teflon thread tape on the cylinder valve.

**Reaction with Selenium.**—In the same manner as above for sulfur, chlorine monofluoride was introduced into 0.5 g of powdered metallic selenium. The infrared spectrum of a 200-mm sample of the gaseous products showed the characteristic bands for  $\text{SeF}_4$  at  $745\text{ s}$  and  $735\text{ vs cm}^{-1}$  "diluted" in intensity approximately 1:2 by the presence of chlorine. The chlorine was removed from the  $\text{SeF}_4$  by vacuum distillation at  $0^\circ$  leaving solid  $\text{SeF}_4$  which melted at room temperature and exhibited a vapor pressure of 12 mm. This was further confirmed by the infrared spectrum of this vapor which showed  $\text{SeF}_4$  in high purity.

Additional chlorine monofluoride was introduced (to a pressure of 5 psi) onto the liquid  $\text{SeF}_4$  and the reaction cylinder was heated at  $200^\circ$  in a fluidized sand bed for 2 hr. The infrared spectrum of the gaseous products showed no  $\text{SeF}_6$  at either  $1484$  or  $1438\text{ cm}^{-1}$ , only unreacted  $\text{ClF}$  and  $\text{SeF}_4$ .

**Reaction of Tungsten and Its Derivatives.**—Chlorine monofluoride, 1 g (15.5 mmol), was introduced onto 1 g (5.5 mg-atom) of powdered tungsten in the same procedure as described previously for sulfur and selenium. After reaction, the cylinder was cooled to  $-78^\circ$  in a trichloroethylene-Dry Ice bath. The product liquid chlorine, as shown by its vapor pressure of about 100 mm, was then removed by pumping for several hours at  $-78^\circ$ . On warming to room temperature the infrared spectrum of the remaining gaseous products showed pure  $\text{WF}_6$  at  $1480\text{ m}$ ,  $1380\text{ m}$ ,  $795\text{ s}$ , and  $712\text{ vvs cm}^{-1}$ , but there was no indication for  $\text{WF}_5\text{Cl}$ .

Following the procedure above, 2.0 g (31.0 mmol) of  $\text{ClF}$  was allowed to react with 1.5 g (6.5 mmol) of tungstic anhydride,  $\text{WO}_3$ . After distilling off the chlorine formed at  $-78^\circ$ , the infra-


(24) E. K. Plyler and W. S. Benedict, *J. Res. Natl. Bur. Std.*, **47**, 202 (1951).

(25) A. H. Nielsen, T. G. Burke, P. J. H. Woltz, and E. A. Jones, *J. Chem. Phys.*, **20**, 596 (1952).

red spectrum of the remaining gaseous products showed  $\text{ClO}_2\text{F}$  at 1258 vs and 1100  $\text{s cm}^{-1}$  in addition to the  $\text{WF}_6$ .

As above 1.0 g (15.5 mmol) of chlorine monofluoride was allowed to react with 0.5 g (1.1 mmol) of lead tungstate,  $\text{PbWO}_4$ , at room temperature. However, only a trace of  $\text{WF}_6$  was formed as evidenced in the infrared spectrum of the product gases which showed mostly unreacted  $\text{ClF}$ . The reaction cylinder was then suspended in a vertical electric furnace, heated at  $300^\circ$  for 2 hr, and allowed to cool to room temperature. At this point the gaseous products were separated as before and proved to be  $\text{WF}_6$ ,  $\text{ClO}_2\text{F}$ , and chlorine. The cylinder was opened under a blanket of dry nitrogen and a pale yellow solid was recovered and transferred immediately into a nitrogen-filled Nalgene Teflon (FEP) 1600 bottle. The solid was found to hydrolyze rapidly in air turning brown and suggesting that it was  $\text{PbF}_4$ . *Anal.* Calcd for  $\text{PbF}_4$ : F, 26.8. Found: F, 26.1.

The reaction of 1.0 g (15.5 mmol) of  $\text{ClF}$  with 0.5 g (1.7 mmol) of anhydrous sodium tungstate again yielded only a trace of free  $\text{WF}_6$  at room temperature. After heating at  $300^\circ$  for 2 hr the only gaseous products proved to be  $\text{ClO}_2\text{F}$  and chlorine with again only a trace of  $\text{WF}_6$  showing up in the infrared region. A white solid was recovered from the reactor under a blanket of nitrogen. It was suspected to be the known  $2\text{NaF}\cdot\text{WF}_6$  complex.<sup>26</sup> Crushed ice was added to a weighed amount of the solid in a Teflon bottle and the bottle was capped. Hydrolysis of the material gave a clear acidic solution. The "active fluorine" in the product was determined by titrating the acidity generated from the hydrolysis with 0.5 N NaOH. *Anal.* Calcd



for  $2\text{NaF}\cdot\text{WF}_6$ : active F, 29.9. Found: active F, 28.0.

**Reaction of Molybdenum and Its Derivatives.**—In a manner similar to the procedure used above for sulfur, selenium, and tungsten, 1.5 g (22.3 mmol) of chlorine monofluoride was allowed to react with 1.0 g (10.4 mg-atom) of powdered molybdenum at room temperature. The reaction cylinder was cooled to  $-78^\circ$  and the liquid chlorine formed in the reaction was pumped off as before. The infrared spectrum of the gaseous products remaining showed the presence of high-purity  $\text{MoF}_6$  at 1475 vvw, 1385 vvw, and 741 vvs  $\text{cm}^{-1}$ .

Reaction of 1.5 g (22.3 mmol) of chlorine monofluoride with 0.5 g (3.1 mmol) of  $\text{MoS}_2$  yielded, in addition to chlorine and  $\text{MoF}_6$ ,  $\text{SF}_4$  as evidenced by its infrared absorption at 889 vs, 738 sh, 728 sh, and 715  $\text{s cm}^{-1}$ .

In addition to the gaseous products described, traces of blue-green solids were recovered from the two reactions above and, when exposed to moist air, rapidly hydrolyzed to dark blue inks. This suggests lower molybdenum fluorides which are known to hydrolyze to the blue oxyfluorides.<sup>27</sup>

Chlorine monofluoride, 1.5 g (22.3 mmol), was added to 0.5 g (2.1 mmol) of potassium molybdate utilizing the same general procedure. At room temperature chlorine and  $\text{ClO}_2\text{F}$  proved to be the only major gaseous products with only a trace of  $\text{MoF}_6$  observed in the infrared spectrum. The reaction cylinder was placed in the electric furnace and heated at  $300^\circ$  for 2 hr. Then, after cooling the cylinder in a  $-78^\circ$  trichloroethylene-Dry Ice bath, the product chlorine and a trace of unreacted  $\text{ClF}$  were pumped off. The infrared analysis of the remaining trace of gaseous products showed high-purity  $\text{MoF}_6$  at 1475, 1385, and 741  $\text{cm}^{-1}$ . After the cylinder was completely evacuated of all gases at room temperature, a pale gray solid remained. This appeared to be the known  $2\text{KF}\cdot\text{MoF}_6$  complex,<sup>28</sup> and the hydrolysis in ice water gave an acidic solution which was titrated with 0.5 N NaOH for "active fluorine." *Anal.* Calcd for





$2\text{KF}\cdot\text{MoF}_6$ : active F, 35.0. Found: active F, 34.8.


The reaction of  $\text{ClF}$  with anhydrous sodium molybdate proved virtually identical with the reaction of  $\text{ClF}$  with  $\text{K}_2\text{MoO}_4$ . The reaction of 2.0 g (31.0 mmol) of  $\text{ClF}$  with 0.5 g (2.4 mmol) of  $\text{Na}_2\text{MoO}_4$  at  $300^\circ$  resulted in gaseous chlorine,  $\text{ClO}_2\text{F}$ , and a light gray-brown solid again suggesting the known complex  $2\text{NaF}\cdot\text{MoF}_6$ .<sup>28</sup> *Anal.* Calcd for  $2\text{NaF}\cdot\text{MoF}_6$ : F, 55.7. Found: F, 55.1.

## Results and Discussion


The reaction of chlorine monofluoride with sulfur and selenium was found to proceed quite smoothly at room temperature to yield only the respective tetrafluorides




The complete absence of  $\text{SF}_5\text{Cl}$  in the products from the reaction at  $200^\circ$  between  $\text{SF}_4$  and  $\text{ClF}$  in the presence of chlorine does prove to be somewhat of an enigma.



Nyman and Roberts<sup>4</sup> have claimed that chlorine monofluoride prepared *in situ* reacts with  $\text{SF}_4$  to form  $\text{SF}_5\text{Cl}$  in 90% yield






The reaction of  $\text{SF}_4$  and cesium fluoride with chlorine has also been reported to yield  $\text{SF}_5\text{Cl}$ <sup>29</sup>



The failure of chlorine monofluoride to fluorinate  $\text{SeF}_4$  further to the hexafluoride even up to  $200^\circ$  is also somewhat surprising but perhaps indicative of its more moderate reactivity as a fluorinating agent.

With elemental tungsten and molybdenum, as well as molybdenum disulfide, fluorination to the respective hexafluorides proceeded smoothly at room temperature



The reaction of sodium and potassium molybdates and sodium tungstate with  $\text{ClF}$  at  $300^\circ$  yielded the  $2\text{MF}\cdot\text{MoF}_6$  and  $2\text{MF}\cdot\text{WF}_6$  type of complexes. However, with  $\text{PbWO}_4$  the products were  $\text{PbF}_4$  and  $\text{WF}_6$ . The failure of a similar  $\text{PbF}_4\cdot\text{WF}_6$  complex to form is best explained on the basis of the more polarizing character of  $\text{Pb}^{4+}$  vs.  $\text{Na}^+$  or  $\text{K}^+$ .

**Acknowledgment.**—The authors wish to thank the Olin Central Analytical Department and specifically Mr. W. A. Nichols for the vpc separation work.

(26) S. Katz, *Inorg. Chem.*, **3**, 1598 (1964).

(27) D. E. LaValle, R. M. Steele, M. K. Wilkinson, and H. L. Yakel, Jr., *J. Am. Chem. Soc.*, **82**, 2433 (1960).

(28) G. B. Hargreaves and R. D. Peacock, *J. Chem. Soc.*, 4390 (1958).

(29) C. W. Tullock, D. D. Coffman, and E. L. Muetterties, *J. Am. Chem. Soc.*, **86**, 357 (1964).