structure factor calculations at their calculated positions. The positional and anisotropic thermal parameters were refined by block-matrix/fullmatrix procedures.

The atomic scattering factors were those from ref 35 modified for the real and imaginary parts of anomalous dispersion.

(35) Cromer, D. T.; Waber, J. T. International Tables for X-Ray Crystallography; Kynoch Press: Birmingham, England, 1974; Vol. IV, Table 2.2A, pp 71-79. Templeton, D. H. International Tables for X-Ray Crystallography; Kynoch Press: Birmingham, England, 1962; Vol. III, Table 3.3.2.c, pp 215-216.

Acknowledgment. This material is based upon work supported by the National Science Foundation under Grant CHE-8401844 (R.J.A.). X-ray diffraction studies were supported by the U.S. Department of Energy, Office of Basic Energy Sciences, Materials Sciences Division, Under Contract W-7405-ENG-82. We thank Dr. N. Kostić for helpful discussions.

Supplementary Material Available: Tables of positional parameters, anisotropic thermal parameters, all bond distances and angles, and structure factors (32 pages). Ordering information is given on any current masthead page.

Coreceptor Molecules. Synthesis of Metalloreceptors Containing Porphyrin Subunits and Formation of Mixed Substrate Supermolecules by Binding of Organic Substrates and of Metal Ions

Andrew Hamilton,[†] Jean-Marie Lehn,^{*} and Jonathan L. Sessler[‡]

Contribution from the Institut Le Bel, Université Louis Pasteur, 67000 Strasbourg, France. Received January 2, 1986

Abstract: Three coreceptor molecules, two macrotetracycles 1 and 2 and one macropentacycle 3, containing [18]-N₂O₄ macrocyclic and porphyrinic subunits, have been synthesized and characterized. They form metalloreceptors by complexation of transition-metal ions like Zn(II) or Cu(II) with the porphyrin groups. Binding of α,ω -diammonium cations to the [18]-N₂O₄ units occurs with substrate inclusion, yielding η^2 -cryptates like 27, 28, and 31, as indicated by very large upfield shifts of the CH₂ proton NMR signals of the substrates. Similarly, these organic cations also bind to the Zn(II)-containing metalloreceptors 19, 20, and 22, giving organic/inorganic mixed substrate supermolecules, such as 29, 30, and 32. In these and similar species, the simultaneous complexation of several substrates opens the way to the study of mutual interactions within supramolecular structures and of regulation effects on physical and chemical properties.

Beyond receptor molecules possessing a single binding unit, the chemistry of molecular receptors enters the domain of polytopic coreceptors, macropolycyclic molecules incorporating several binding subunits which may cooperate for the complexation of either several singly bound substrates or of multiply bound po-lyfunctional substrates.^{1,2} Homotopic coreceptors which bind terminal diammonium cations and dicarboxylate anions by means of two identical subunits have been described (see references in ref 1 and below).

Metalloreceptors are heterotopic coreceptors which combine suitable substrate selective binding subunits, such that they can complex both organic substrate(s) and metal cation(s), thus forming mixed substrate supermolecules. As a consequence, interactions and reactions may take place between metal-centered reactive sites and co-bound molecular substrate(s). Such metalloreceptors, -reagents, and -catalysts may also mimic features of metalloproteins.

Earlier examples of a process which may be considered to be of the metalloreceptor type were the formation of molecular cation-anion pairs by a cascade mechanism which involved first complexation of one or two metal cations to a neutral aporeceptor followed by ion pairing between the resulting holoreceptor cation and a molecular anion.^{3,4}

We have designed and studied in recent years several metalloreceptor systems (for brief descriptions see ref 1 and 2). We

now describe the synthesis of the three coreceptors 1-3 and report their ability to complex both metal cations and primary diammonium ions, by means of porphyrin and macrocyclic [18]- N_2O_4 (12) subunits, respectively.^{5,6}

(5) Preliminary communication, see: Hamilton, A. D.; Lehn, J. M.; Sessler, J. L. J. Chem. Soc., Chem. Commun. 1984, 311-313.

0002-7863/86/1508-5158\$01.50/0 © 1986 American Chemical Society

Present address: Chemistry Department, Princeton University.

^tChemistry Department, University of Texas, Austin. [⊥] UA 422 of the CNRS.

For a general presentation of the concepts and definitions see: Lehn, J. M. In *Biomimetic Chemistry*; Yoshida, Z. I., Ise, N., Eds.; Kodansha Ltd.: Tokyo; Elsevier Scientific Publishing Co.: Amsterdam, 1983; pp 163-187. See also: Lehn, J. M. Science 1985, 227, 849-856.
 (2) Lehn, J. M. In *IUPAC Frontiers of Chemistry*; Laidler, K. J., Ed.;

⁽²⁾ Lehn, J. M. In *IUPAC Frontiers of Chemistry*; Laidler, K. J., Ed.;
Pergamon Press: Oxford and New York, 1982; pp 265-272.
(3) Lehn, J. M.; Simon, J.; Wagner, J. Angew. Chem. 1973, 85, 622-624;
Angew. Chem., Int. Ed. Engl. 1973, 12, 579-580. Dietrich, B.; Lehn, J. M.;
Simon, J. Angew. Chem. 1974, 86, 443-444; Angew. Chem., Int. Ed. Engl.
1974, 13, 406-407. Lehn, J. M.; Simon, J.; Moradpour, A. Helv. Chim. Acta
1978, 61, 2407-2418.

⁽⁴⁾ For metalloreceptors derived from α -cyclodextrins see: Breslow, R.; Overman, L. E. J. Am. Chem. Soc. 1970, 92, 1075-1077. Tabushi, I.;
 Shimizu, N.; Sugimoto, T.; Shokuza, M.; Yamamura, K. J. Am. Chem. Soc.
 1977, 99, 7100-7102. For multiple binding effects see also: Rebek, J., Jr.
 Acc. Chem. Res. 1984, 17, 258-264. Willner, I.; Goren, Z. J. Chem. Soc., Chem. Commun. 1983, 1469-1470.

<sup>sler, J. L. J. Chem. Soc., Chem. Commun. 1984, 311-313.
(6) Other systems containing porphyrin and polyhetero macrocyclic groups have appeared: Chang, C. K. J. Am. Chem. Soc. 1977, 99, 2819-2822.
Thanabal, V.; Khrisnan, V. Ibid. 1982, 104, 3643-3650. For a recent review on variously bridged porphyrins, see: Baldwin, J. E.; Perlmutter, P. Top. Current Chem. 1984, 121, 181-220. For non-porphyrinic systems see for instance: Busch, D. H.; Christoph, G. C.; Zimmer, L. L.; Jackels, S. C.; Grzybowski, J. J.; Callahan, R. C.; Kojima, M.; Holter, K. A.; Mocak, J.; Herron, N.; Chavan, M.; Schammel, W. P. J. Am. Chem. Soc. 1981, 103, 5107-5114</sup> 5107-5114.

Scheme I



Scheme II



Results

Synthesis of Metalloreceptors 1-3. The synthetic route used to obtain the macrotetracycles 1 and 2 is shown in Scheme I. It follows the strategy that allows introduction of different bridges between the two macrocyclic subunits of a cylindrical macrotricycle (see Figure 7, route A, in ref 7). The porphyrin dicarboxylic acids 4 and 5 were prepared according to the literature^{8,9} and subsequently activated by conversion to the bis(p-nitrophenyl) esters 6 and 7^{10} which by condensation with the biphenylenebridged bis-[18]-N₂O₄ macrocycle 9 in warm pyridine^{10,11} under

conditions of moderate dilution led to the tetraamides 10 and 11, respectively. Good yields were obtained for these condensations provided long reaction times were employed (up to 48 h were required to reach completion); because of the slow nature of the coupling reaction, the reactants were simply mixed together at the outset of the reaction. The tetraamides 10 and 11 could be obtained in pure form following careful chromatography on silica gel and recrystallization from a dichloromethane-hexane mixture. In general, however, the crude products could be used directly for further synthetic work.



The tetraamines 1 and 2 were obtained from the tetraamides 10 and 11 by using a three-step sequence which involved (i) preparation of the zinc(II) derivatives, (ii) reduction with diborane, and (iii) treatment with concentrated HCl to effect hydrolysis and demetalation. The reduction of the amides proceeded smoothly if this three-step sequence was employed; the crude yields were nearly quantitative. Only poor yields were obtained, however, if the free-base porphyrins were treated directly with diborane. Formation of the zinc complex serves to protect the porphyrin macrocycle from reaction with diborane. Compounds 1 and 2 were obtained in analytically pure, crystalline form following chromatography on alumina and slow recrystallization from a dichloromethane-hexane mixture. A minimum of alumina was used as poor recoveries were observed on this support. Overall, yields of 54% and 45% were obtained for 1 and 2 (based on 6 and 7).

The bis-macrocycle compound 9 was prepared by two methods (Scheme II). The first utilized a direct reaction between dichloroformyl-4,4'-biphenyl (13) and the [18]-N₂O₄ macrocycle 12. The second involved treatment of the monoprotected [18]-

⁽⁷⁾ Lehn, J. M. Pure Appl. Chem. 1980, 52, 2441-2459.
(8) Fischer, H.; Orth, H. Die Chemie des Pyrrols; Akademischer Verlag:

Leipzig, 1937.

 ^{(9) (}a) Chang, C. K. J. Am. Chem. Soc. 1977, 99, 2819-2822. (b) Chang,
 (c) (a) Chem. Ser. 1979, 173, 162-177; (c) Ogoshi, H.; Sugimoto, H.;
 Yoshida, Z. Tetrahedron Lett. 1977, 169-172. (d) Paine, J. B., III In The
 Porphyrins; Dolphin, D., Ed.; Academic Press: New York, 1978; Vol. IA, Porphyrins; Dolphin, D., Ed.; Academic Press: New York, 1978; Vol. 1A, Chapter 4. (e) Bullock, E.; Johnson, A. W.; Markahm, E.; Shaw, K. B. J. Chem. Soc. 1958, 1430-1440. (f) Battersby, A. R.; Buckley, D. G.; Hartley, S. G.; Turnbull, M. D. J. Chem. Soc., Chem. Commun. 1976, 879-881. (g) Smith, K. J. Chem. Soc., Perkin Trans. 1 1972, 1471-1475. (10) (a) Collman, J. P.; Anson, F. C.; Barnes, C. E.; Bencosme, C. S.; Geiger, T.; Evitt, E. R.; Kreh, R. P.; Meier, K.; Pettman, R. B. J. Am. Chem. Soc. 1983, 105 2604-2609. (b) Collman, L. P.: Deniewich, P.: Konsi, Y.;

Soc. 1983, 105, 2694-2699. (b) Collman, J. P.; Denisevich, P.; Konai, Y.; Marrocco, M.; Koval, C.; Anson, F. C. J. Am. Chem. Soc. 1980, 102, 6027-6036.

⁽¹¹⁾ Alberts, A. H.; Annunziata, R.; Lehn, J. M. J. Am. Chem. Soc. 1977, 99, 8502-8504.

Scheme III



Table I. Electronic Spectral Data for Porphyrin-Containing Heterotopic Coreceptors in Chloroform $[\lambda_{max} \text{ in nm } (\log \epsilon)]$ and a Reference Compound

metal	receptor	soret	visible			
2H	1	375 (sh), 397 (5.15)	497 (4.04)	532 (3.90)	567 (3.69)	619 (3.53)
	2	374 (sh), 397 (5.21)	497 (4.12)	531 (3.98)	567 (3.72)	619 (3.61)
	3	388 (4.86)	502 (4.21)	536 (4.03)	571 (3.90)	623 (3.71)
	8 ^a	400	498	536	566	620
Zn	19	405 (5.41)		536 (4.17)	572 (4.17)	
	20	402 (5.39)		536 (4.17)	572 (4.22)	
	22	385 (5.39)		537 (4.07)	572 (4.32)	
Cu	21	399		525	563	
	23	390		530	556	
Cu	22 21 23	385 (3.39) 399 390	, <u></u> ,,,	537 (4.07) 525 530	572 (4.32) 563 556	

^a From ref 15.

 N_2O_4 macrocycle 14¹² with 13 to give 15, which, after hydrogenolysis, afforded 9. Although both routes proceeded in good yield, we favor the latter stepwise approach in that the product 9 was obtained in higher purity.

Two strategies were employed to synthesize the macropentacyclic tetraamine 3. Both involved the generation and subsequent reduction of the tetraamide 16, as outlined in Scheme III. In method A, the key tetraamide 16 (along with numerous impurities) is produced directly by condensing the activated porphyrin 6^{10} with the [18]-N₂O₄ macrocycle 12 under conditions of high dilution. When this reaction was carried out on small scale, the intermediate 16 could be purified by careful chromatography on silica gel. On larger scales, however, the low solubility of 16 in common organic solvents precluded such purifications. In practice, therefore, the crude reaction mixture was subjected directly to the three-step reducing procedure outlined above. Complete purification was then effected at the terminal tetraamine stage. The tetraamine 3 is far more soluble than the tetraamide precursor 16 and could be purified by repeated chromatography on alumina (using a minimum of support to avoid substantial material losses) followed by a recrystallization from dichloromethane/hexane. Unfortunately, the yields of 3 obtained by this direct condensation approach (method A, Scheme III) were low (ca. 10% based on 6).

Satisfactory yields of 3 were obtained by using a stepwise procedure (method B, Scheme III). In this approach, the monoprotected [18]-N₂O₄ compound 14^{12} is first condensed with the activated porphyrin 6^{10} to yield the porphyrin-bridged, bismacrocyclic compound 17 in good yield. Standard methods (e.g., H₂/Pd or HBr/HOAc) failed to cleave the carbobenzoxy protecting group cleanly. Deprotection could be effected smoothly by treatment with a mixture of concentrated hydrochloric and trifluoroacetic acids. The reaction was monitored carefully as under these conditions the rate of amide hydrolysis is apparently close to that of carbamate cleavage. In favorable cases, the crude diamide-diamine 18 was obtained in sufficient purity for use in the subsequent steps. When required, purification was effected by chromatography on alumina. A second condensation reaction with 6^{10} then gave the intermediate tetraamide 16, in a relatively pure state. After reduction, the final purifications of 3 were far easier than those required when the direct condensation approach was employed. The overall yield of 3 with use of the stepwise procedure was ca. 25% (based on the monoprotected macrocycle 14).

Formation of the Metalloporphyrin Complexes. In general, as metal binding to a porphyrin site occurs, the four absorptions in

^{(12) (}a) Lehn, J. M.; Simon, J.; Wagner, J. Angew. Chem. 1973, 85, 621-622; Angew Chem., Int. Ed. Engl. 1973, 12, 578-579. (b) Lehn, J. M.; Simon, J.; Wagner, J. Nouv. J. Chim. 1978, 1, 77-84.



Figure 1. Spectrophotometric titration of 1 with zinc acetate: generation of metalloreceptor 19. Curve A; 1. Curve B; 19, obtained by the addition of 1.05 equiv of zinc(II). Intermediate curves obtained by adding increasing quantities of zinc acetate in CH₃OH to the initial solution of 1 in CHCl₃. Spectra were recorded with use of a gas-tight cuvette maintained at 40 °C for 30 min between runs.

the 450-700-nm region disappear and are replaced by one or, more frequently, two bands.¹³ Electronic spectroscopy was therefore used to demonstrate the formation of the zinc(II) and copper(II) derivatives 19-23 of the free-base porphyrins 1-3. Spectral data for these complexes are given in Table I.

The zinc(II) porphyrin complexes were produced cleanly. For instance, when a solution of the tetraamine 1 in chloroform, maintained at 40 °C in a gas-tight cuvette, was titrated over the course of several hours with 1 equiv of zinc(II) acetate in methanol, complete conversion to the zinc(II) derivative 19 was achieved (Figure 1). That clean isosbestic behavior was observed during the course of the titration suggests that metal insertion into the porphyrin unit occurred quantitatively without competing complexation into the [18]-N2O4 macrocyclic sites. Similar isosbestic behavior was observed when either 2 or 3 was titrated under similar conditions with 1 equiv of zinc(II) acetate per porphyrin group.

When analogous titrations of 1 and 3 were performed with copper(II) acetate, isosbestic behavior was not observed. Apparently copper binding into the [18]-N₂O₄ sites competes with incorporation into the porphyrin unit. This finding is consistent with the observation that the [18]-N₂O₄ macrocycle, 12, displays an ca. 10^4 higher affinity for Cu(II) than for Zn(II).¹⁴ The copper(II) cation bound in the [18]-N₂O₄ sites is more labile than that held in the porphyrin macrocycle. It thus proved possible to prepare the pure copper porphyrin complexes, 21 and 23, by adding an excess of copper(II) acetate dissolved in MeOH to solutions of the porphyrins 1 and 3 in boiling CHCl₃ and then washing the solutions with aqueous sodium cyanide.

Characterization of the Receptors 1-3 and of Their Metal Complexes 19-23. UV-Visible Absorption Spectra. UV-visible spectral data for the three porphyrin-containing receptors and for their metal complexes described in this report are summarized in Table I. The electronic spectra of the free-base compounds 1 and 2 display etio-type features which are typical of monomeric porphyrins with 6 or more alkyl groups in the β -pyrrolic positions.¹³ Specifically, an intense Soret at λ_{max} 397 nm and four satellite bands of decreasing intensity in the region between 500 and 700 nm were observed. An essentially etio-type spectrum was also displayed by the dimeric porphyrin compound 3. In this case,

Table II. Representative ¹H NMR Data for Porphyrin-Containing Receptors and Reference Compounds

	chemical shift," ppm					
compd	NH (pyrrole)	CH ₃ (ring)	H (meso)	H (phenyl)		
8 ^b	-3.75	3.67	10.10, 10.16			
24 ^c				7.35, 7.40		
1	-3.74	3.54	10.02, 10.09	5.90, 6.41		
2	-3.69	3.62, 3.64	10.06, 10.13	6.41, 6.66		
3	-4.39	2.85, 3.04	8.30, 9.76			

^a In CDCl₃ at 200 MHz, unless otherwise noted. ^b In CDCl₃ at 100 MHz, from ref 10. 'In CDCl₃ at 250 MHz, from ref 17.



Figure 2. Proton NMR spectrum of receptor 1 at 400 MHz $(CD_2Cl_2:CD_3OD 9:1 (v/v))$. Due to exchange, no signals for the internal pyrrole N-H protons are observed in this solvent system. Peaks marked with x and S are due to impurities and solvent, respectively. For assignments of the other peaks, see Experimental Section.

however, the Soret band was broadened and blue shifted by 11 nm and the visible bands slightly red-shifted as compared to the monomers 1 and 2, or to the diethyl ester 8.15 Such spectral shifts have previously been observed for "face-to-face" dimeric porphyrin systems in which the two prophyrin subunits are covalently constrained to adopt a close orientation.^{10,15,16} Both the zinc(II) and copper(II) derivatives of the dimeric porphyrin receptor (22 and 23) also had blue-shifted Soret bands as compared to the corresponding monoporphyrin complexes (i.e., 19, 20, and 21).

¹H NMR Spectra. ¹H NMR spectroscopy was used to characterize the porphyrin-containing receptors 1, 2, and 3. With exception of the tetraamides 10, 11, and 16 which displayed poorly resolved spectra, all intermediates reported in this study were also characterized by ¹H NMR spectroscopy. Peak assignments were made on the basis of integrations, spectral intercomparisons of similar compounds, and homodecoupling studies. Selected data are summarized in Table II; complete tabulations for each compound are included in the Experimental Section.

In the spectra of 1 and 2 signals due to both the biphenyl and porphyrin subunits are clearly discernable (Figure 2). Of particular interest are the peaks ascribable to the two nonequivalent biphenyl protons. As compared to the analogous bis-biphenyl

^{(13) (}a) White, W. I. In *The Porphyrins*; Dolphin, D., Ed.; Academic Press: New York, 1978; Vol. V, Chapter 7. (b) Gouterman, M. In *The Porphyrins*; Dolphin, D., Ed.; Academic Press: New York, 1978; Vol. IIIA, Chapter I. (c) Smith, K. M. In Porphyrins and Metalloporphyrins; Smith,
K. M., Ed.; Elsevier: Amsterdam, 1975.
(14) (a) Spiess, B.; Arnaud-Neu, F.; Schwing-Weill, M. J. Helv. Chim. Acta 1979, 62, 1531-1542. (b) Anderegg, G. Helv. Chim. Acta 1981, 64,

^{1790-1799.}

⁽¹⁵⁾ Collman, J. P.; Bencosme, C. S.; Durand, R. R., Jr.; Kreh, R. P.;

⁽¹⁵⁾ Collman, J. P.; Bencosme, C. S.; Durand, R. R., Jr.; Kreh, R. P.; Anson, F. C. J. Am. Chem. Soc. 1983, 105, 2699-2703.
(16) (a) Collman, J. P.; Chong, A. O.; Jameson, G. B.; Oakley, R. T.; Rose, E.; Schmittou, E. R.; Ibers, J. A. J. Am. Chem. Soc. 1981, 103, 516-533. (b) Collman, J. P.; Elliott, C. M.; Halbert, T. R.; Tovrog, B. S. Proc. Natl. Acad. Sci. U.S.A. 1977, 74, 18-22. (c) Chang, C. K.; Kuo, M. S.; Wang, C. B. J. Heterocycl. Chem. 1977, 14, 943-945. (c) Chang, C. K. J. Chem. Soc., Chem. Commun. 1977, 800-801. (e) Ward, B.; Wang, C. G.; Chang, C. K. J. Am. Chem. Soc. 1981, 103, 5236-5238. (f) Chang, C. K.; Abdalmuhdi, I. J. Org. Chem. 1983, 48, 5380-5390. (g) Chang, C. K.; Abdalmuhdi, I. J. Org. Chem. 1983, 96, 154-155; Angew. Chem., Int. Ed. Engl. 1984, 23, 164-165. (h) Collman, J. P.; Anson, F. C.; Bencosme, S.; Chong, A.; Collins, T.; Denisevich, P.; Evitt, E.; Geiger, T.; Ibers, J. A.; Jameson, G.; Konai, Y.; Koval, C.; Meier, K.; Oakley, R.; Pettman, R.; Schmittou, E.; Sessler, J. L. In Organic Synthesis Today and Tomorrow (IUPAC); Hutchinson, C. R., Ed.; Pergamon: Oxford, 1981, pp 29-45. (IUPAC); Hutchinson, C. R., Ed.; Pergamon: Oxford, 1981, pp 29-45.



Figure 3. Proton NMR spectrum of receptor 3 at 200 MHz ($CDCl_3$); the signal of the internal NH protons is observed upfield at -4.39 ppm.

receptor 24,¹⁷ these signals are shifted upfield by 0.94 and 0.74 ppm for 2 and 1.45 and 0.99 ppm for 1. These upfield shifts suggest that the biphenyl protons of 1 and 2 are shielded by the porphyrin macrocycle and imply that in solution, as in the crystalline state (vide infra), the biphenyl subunits lie above the porphyrin rings.



Shielding effects are also observed in the case of the dimeric porphyrin receptor 3 (Figure 3). For instance, the internal pyrrole protons resonate at 0.55 ppm higher field in the dimer 3 than in the monomer 1, a behavior which is qualitatively consistent with that observed for several covalently linked, cofacial porphyrin dimers.^{10,16} This suggests that conformations which allow for complete or partial overlap of the two porphyrin macrocycles occur in solution. In the dimer 3 several other signals, including those of the meso and porphyrin methyl protons, are also shifted to higher field (Table II).

As is the case for the members of the β -linked "face-to-face" dimeric porphyrin series, ^{9,10,16c,d,e,h} compound 3 could exist as a mixture of two diastereomers in which the relative orientation of the two rings is different. Space filling molecular models suggest, however, that ring flipping and interconversion between the two forms is possible in the case of 3. For a mixture of two diastereomeric β -linked porphyrins, up to eight signals are possible for the meso protons. Although somewhat broadened as compared to the monomers 1 and 2, only two peaks are observed in this region for the dimer 3. At present it is difficult to ascertain if only one diastereomer is produced in the coupling reaction, or if, more likely, the two diastereomers which might be formed interconvert rapidly on the NMR time scale.¹⁸

The zinc(II) derivatives 19, 20, and 22 are diamagnetic and their ¹H NMR spectra were recorded. In general, the basic spectral features were similar to those of the free-base compounds 1–3. Although no efforts were made to make detailed assignments,



Figure 4. ¹³C NMR spectrum of 1 at 50 MHz (CDCl₃). Peaks marked with S are due to solvent. For peak assignments, see Experimental Section.



Figure 5. ¹H NMR spectrum at 200 MHz (CD₂Cl₂:CD₃OD 9:1 (v/v)) of the supramolecular complex 27 formed from the receptor 1 and the S_9^{2+} substrate ⁺H₃N-(CH₂)₉-NH₃⁺ dipicrate.

it is to be noted that the spectra of these complexes displayed no signals corresponding to the internal pyrrole protons and hence no peaks upfield of Me_4Si .

¹³C NMR Spectra. Well-resolved ¹³C NMR spectra, which are consistent with the proposed structures, were obtained for the receptors 1-3 and several key intermediates. For instance, the ¹³C NMR of **1** is shown in Figure 4. Tentative peak assignments for these compounds were made on the basis of intercomparisons and by analogy to simpler compounds; these data are included in the Experimental Section. In general, the ¹³C signals of the receptors 1 and 2 lie within 3 ppm of those for appropriate reference compounds (e.g., 8 and 26). Exceptions are the two signals of carbons 1 and 4 of the biphenyl moiety. Compared to the bis-biphenyl receptor, 24, ^{17b,19} these signals are split and shifted to both higher and lower field. These shifts may again derive from the fact that the biphenyl subunit lies above the porphyrin plane. A comparison between the porphyrin dimer 3 and the monomer 1 (or the "crowned" porphyrin of Chang^{9a}) reveals that the signals of the pyrrole α and β carbons have been shifted to *lower* field by ca. 8-10 ppm; no other signals are shifted by more than ca. 3 ppm.

Solid-State Structure. An X-ray structural determination of complex **21** was performed.²⁰ It confirmed the structure of this compound and, by extension, those of the other compounds synthesized in the course of this work. It also showed that one phenyl group of the biphenyl unit lies face to face above the porphyrin site.

Formation of the Supermolecular Species 27, 28, 31 by Binding of Organic Substrate to the Receptors 1–3. The qualitative proton

^{(17) (}a) Kotzyba-Hibert, F.; Lehn, J. M.; Vierling, P. Tetrahedron Lett. 1980, 941-944. (b) Kotzyba-Hibert, F. Ph.D. Thesis, Université Louis Pasteur de Strasbourg, 1983.

⁽¹⁸⁾ Similarly, compounds 1 and 2 present *ansa*-type asymmetry which would be observable provided that the rate of ring flipping were slow on the NMR time scale.

⁽¹⁹⁾ Kintzinger, J. P.; Kotzyba-Hibert, F.; Lehn, J. M.; Pagelot, A.; Saigo, K. J. Chem. Soc., Chem. Commun. 1981, 833-836.

⁽²⁰⁾ Chevrier, B.; Moras, D., unpublished results.

Table III. ¹H NMR Chemical Shifts of the Bound ⁺H₃N-[CH₂]_n-NH₃⁺, S_n^{2+} Substrates in the Molecular Cryptates of Coreceptors 1-3 and Metalloreceptors 19, 20, and 22 and Chemical Shift Data for Reference Systems^a

		substrate chemical shifts					
cryptand	substrate	α	β	γ	δ	é	
1 1 1	$\frac{S_8^{2+}}{S_9^{2+}}$	0.60 b b	$\begin{array}{c} -0.67, -1.25^{c} \\ -0.08, -0.28^{c} \\ 0.4 \end{array}$	-1.83 -1.27, -1.53° -0.85	-2.40 -2.00, -2.20 ^c -1.88	-2.69 -2.45, -2.99°	
2 2 2 3	S_8^{2+} S_9^{2+} S_{10}^{2+} S_8^{2+}	ь ь ь	b 0.25 b -0.35, -0.94 ^c	b -1.03 -0.62 -2.13, -2.57 ^c	-1.5 -1.60 -1.53 -4.53	-2.38 -2.19	
3 3	S_{9}^{2+} S_{10}^{2+}	b b	b -0.96	-2.2 -2.43	-4.0 -3.43	-5.5 -4.75	
24 ^d 24 ^d 24 ^d	${S_6}^{2+}$ ${S_7}^{2+}$ ${S_8}^{2+}$	b 2.19 2.51	0.32 0.04 0.48	0.32 -0.56 0.32	0.04 0.32		
26 ^d 26 ^d 26 ^d	${S_8}^{2+}$ ${S_9}^{2+}$ ${S_{10}}^{2+}$	2.78 2.74 2.73	1.50 1.55 1.55	1.34 1.26 1.28	1.34 1.26 1.28	1.26 1.28	
8	S ₉ ²⁺	1.83	3.18	3.21	3.21	3.21	
19 19 19	S_8^{2+} S_9^{2+} S_{10}^{2+}	b 0.32 b	-0.83, -1.37 ^c 0.02, -0.44 ^c 0.42	-1.91 -1.24, -1.48° -0.98	-2.43 -2.11 -1.91	-2.60 -2.48, -3.00 ^c	
20 20 20	S_8^{2+} S_9^{2+} S_{10}^{2+}	ь ь ь	b 0.28 b	-1.4 -1.03 -0.69	-1.6 -1.69 -1.48	-2.47 -2.32	
22 22 22	S_8^{2+} S_9^{2+} S_{10}^{2+}	0.79 0.48 b	-2.31 -1.64 -0.60	-3.12 -2.88 -2.3	-4.36 -4.88 -4.3	-6.80 -5.3	

^a 200-MHz ¹H NMR spectra in CD₂Cl₂:CD₃OD 9:1 (v/v) at 18 °C. Shifts δ given vs. Me₄Si. Peak assignments were based in part on comparisons with "strapped" porphyrins²² and by analogy to earlier work.^{17,23} ^b Not assigned. Peak is not resolved or is hidden under resonances of the receptor. ^cDoublet observed. ^d 250-MHz ¹H NMR spectra in CD₂Cl₂:CD₃OD 9:1 (v/v) at 18 °C. Shifts δ given vs. Me₄Si. From ref 17.

NMR method described earlier^{17,21} was used to study the binding of organic dications, derived from α,ω -diamines, to the receptors 1-3 and to the metalloreceptors 19, 20, and 22. In general, the ¹H NMR spectra of the receptor in CD₂Cl₂:CD₃OD 9:1 (v/v) was recorded; slightly over 1 equiv of the appropriate diammonium dipicrate was then added and the spectrum recorded again. In



cases where substrate binding occurred, peaks ascribable to the CH₂ signals of the substrate were observed at high field, usually upfield of Me₄Si. The results of these studies are summarized in Table III. Figure 5 shows the ¹H NMR spectrum of the complex formed by the binding of ⁺H₃N-(CH₂)_n-NH₃⁺, n = 9

(substrate S_9^{2+}), in receptor 1. Peaks upfield of Me₄Si are clearly visible for the β , γ , δ , and ϵ CH₂ groups of the organic substrate. No such upfield peaks are seen in the ¹H NMR of 1 in the absence of the substrate. Nor are they observed when S_9^{2+} is bound by 2 equiv of the dimethyl [18]- N_2O_4 macrocycle 26¹⁷ (Table III). In fact, as compared to this reference system, the ϵ CH₂ peak of S_9^{2+} bound to 1, at -2.60 ppm, is shifted upfield by ca. 3.9 ppm. The upfield shifts observed here are also considerably greater than those shown by related complexes formed from 24,¹⁷ the bis-biphenyl bridged analogue of 1 (Table III). They may be attributed to the shielding effect of the bridging porphyrin and, to a lesser extent, of the biphenyl bridging groups. Upfield shifts are also found for various "strapped" porphyrins²² in which an alkyl chain is covalently constained to lie above the porphyrin plane. Consequently, as with the other macrocyclic diammonium η^2 -cryp-tates,^{17,19,21,23,24} the S₉²⁺ substrate is bound as a 1:1 molecular complex within the central cavity of receptor 1 by an anchoring of the primary alkyl ammonium groups to the two [18]-N₂O₄ subunits. Structures 27-30 show a schematic representation of the supramolecular complex formed from 1 and S_9^{2+} ; it is derived by analogy to earlier spectroscopic^{17,19,21,23} and structural²⁴ studies.

Receptor 1 similarly complexes substrates S_8^{2+} and S_{10}^{2+} . Substrates S_7^{2+} and S_{11}^{2+} , on the other hand, are not bound, and no upfield shifts are observed under the same conditions. Substrate binding to 1 is therefore selective.^{17a,19,21,23,24} The 1/1 stoichiometry

⁽²¹⁾ See also: Jones, N. F.; Kumar, A.; Sutherland, I. O. J. Chem. Soc., Chem. Commun. 1981, 990-992.

^{(22) (}a) Wijesekera, T. P.; Paine, J. B., III; Dolphin, D.; Eisenstein, F. N. B.; Jones, T. J. Am. Chem. Soc. **1983**, 105, 6747-6749. (b) Chang, C. K.; Kuo, M. S. J. Am. Chem. Soc. **1979**, 101, 3413-3415. (c) Baldwin, J. E.; Klose, T.; Peters, M. J. Chem. Soc., Chem. Commun. **1976**, 881-883. Baldwin, J. E.; Crossley, M. J.; Klose, T.; O'Rear, E. O., III; Peters, M. K. Tetrahedron **1982**, 38, 27-39. (d) Momenteau, M.; Mispelter, D.; Loock, B.; Bisagni, E. J. Chem. Soc., Perking Trans. 1 **1983**, 189-200. (e) Ogoshi, H.; Sugimoto, H.; Yoshida, Z.-i. Tetrahedron Lett. **1976**, 4477-4480. (23) Kotzyba-Hibert F.: Lehn I. M.; Saigo K. J. Am. Chem. Soc. **1981**.

⁽²³⁾ Kotzyba-Hibert, F.; Lehn, J. M.; Saigo, K. J. Am. Chem. Soc. 1981, 103, 4266-4268.

⁽²⁴⁾ Pascard, C.; Riche, C.; Cesario, M.; Kotzyba-Hibert, F.; Lehn, J. M. J. Chem. Soc., Chem. Commun. 1982, 557-560.



of the complexes formed was established by integration studies. Broadened signals for the substrate CH_2 protons resulted when an excess of substrate was present; apparently under these conditions, substrate not only was complexed as an η^2 -cryptate but was also bound in various η^1 external orientations. So far, the relative energetics associated with these two binding modes and with the substrate binding selectivity have not been studied quantitatively.

The homologue 2 also functions as a molecular receptor. Supramolecular complexes of 1/1 stoichiometry are formed selectively with the substrates S_8^{2+} , S_9^{2+} , and S_{10}^{2+} (Table III). As with 1, marked upfield shifts are observed, indicative of the formation of η^2 -cryptates with the substrates held within the central molecular cavity (cf. structure 28). Similar substrate binding occurs for 3 (see below).

An interesting feature of the ¹H NMR spectrum of 27 is that the β , γ , and δ CH₂ signals are split (Figure 5), reflecting the diastereotopic nature of these protons. These splittings suggest that rotation of the ⁺H₃N-(CH₂)₉-NH₃⁺ substrate about teh *N*,*N*-axis and/or rotation of the porphyrin subunits are/is slow. In the latter case, the supramolecular complex 27 would be seen chiral on the NMR time scale. Complexes derived from the larger homologue 2 do not show such splittings but present broad signals, suggesting that greater degrees of freedom pertain in these systems.

Formation of the Mixed Substrate Supermolecules 29, 30, 32 by Binding of Organic Substrate to the Metalloreceptors 19, 20, 22. The heterotopic apo-receptors 1 and 2 can be converted into metal-containing receptors by specifically binding a metal cation within the porphyrin ring; when Zn(II) is used, the diamagnetic complexes 19 and 20 are obtained cleanly from 1 and 2 (see above).²⁵ As for the free-base congeners, ¹H NMR may be used to study organic substrate binding to these metalloreceptors. Figure 6 shows the ¹H NMR spectrum of the complex which derives from the binding of S_9^{2+} into the metalloreceptor 19. As was true for the free-base system, signals ascribable to the β , γ , δ , and ϵ CH₂ protons are observed upfield of Me₄Si, giving peaks which are not present in the absence of substrate. Upfield shifts are also displayed in complexes derived from receptor 19 and substrates S_8^{2+} and S_{10}^{2+} (but not S_7^{2+} or S_{11}^{2+}) as well as from receptor 20 and substrates S_8^{2+} , S_9^{2+} , and S_{10}^{2+} (Table III). Significantly, the complexation of molecular substrates by the metalloreceptors 19 and 20 did not result in demetalation; the visible portion of the electronic spectra was essentially identical both before and after the addition of the diammonium salts. The organic substrates are therefore bound within the central cavities of frameworks which contain bound inorganic centers. The resulting species are organic/inorganic mixed substrate supramo-



Figure 6. ¹H NMR spectrum at 200 MHz (CD₂Cl₂:CD₃OD 9:1 (v/v)) of the supramolecular complex 29 formed from the metalloreceptor 19 and the S₉²⁺ substrate ⁺H₃N-(CH₂)₉-NH₃⁺ dipicrate.



Figure 7. ¹H NMR spectrum at 200 MHz (CD₂Cl₂:CD₃OD 9:1 (v/v)) of the sparingly soluble supramolecular complex 32 formed from the metalloreceptor 22 and the S₉²⁺ substrate ⁺H₃N-(CH₂)₉-NH₃⁺ dipicrate; because of the low solubility of 32, the spectrum also contains the signals of the metalloreceptor 22 and of uncomplexed substrate; the high-field signals (insert) are assigned to the -CH₂- protons of the substrate.



Figure 8. CPK model representation of the possible structure of the mixed substrate supramolecular species 32 formed by binding of ${}^{+}H_{3}N-(CH_{2})_{9}-NH_{3}^{+}$ to the metalloreceptor 22 (see text).

lecular complexes. Structures 29 and 30 are schematic representations of the species which result when both Zn(II) and S_9^{2+} are co-complexed within the heterotopic receptors 1 and 2.

When slightly over 1 equiv of the S_9^{2+} or S_{10}^{2+} substrate was added to either the free-base receptor 3 or its bis-zinc derivative

⁽²⁵⁾ The bis-Co¹¹ complex of 3 has also been obtained recently.²⁷

22, complexes formed which were sparingly soluble.²⁶ Proton NMR analysis of these species is thus complicated by the additional presence of both free receptor and uncomplexed substrate in the solution. Since, however, only the complexed substrate displays CH₂ signals upfield of Me₄Si, the assignment of these peaks remains straightforward. As expected, the upfield shifts of the substrate \mathbf{CH}_i signals, when bound in the porphyrin dimers 3 or 22, are approximately twice that they are when bound in the monomers 1, 2, 19, or 20 (Table III). In fact, the ϵ CH₂ protons, observed at -6.80 ppm for the complex formed from S₉²⁺ and 22 (Figure 7, Table III) are shifted by over 8 ppm as compared to the reference complex formed from 26. Structures 31 and 32 are schematic representations of the organic/inorganic substrate supramolecular species obtained respectively from the heterotopic receptor 3 and from its Zn(II) complex 22. The CPK molecular model shown in Figure 8 provides a possible picture of the structural features of species 32 based on present and earlier^{1,17,19,24} results.

Conclusion

The results described above show that suitably designed macropolycyclic molecules function as heterotopic coreceptors allowing one to bring together within the same supramolecular structure inorganic and organic substrates complexed by means of selective subunits. Mutual interactions between the co-bound substrates should provide means for regulating the physical properties and chemical reactivity of the supramolecular species. This applies in particular to the photochemical and electrochemical properties of the complexes (such as 19-23 and 27-32) formed by the tritopic and tetratopic receptors 1-3 described here, since these species contain photoactive and electroactive subunits; such results have been obtained and will be described in another report.²⁷ As pointed out earlier,¹ polytopic receptors may possess properties such as regulation, cooperativity, and allostery, which are not present in monotopic receptors. Furthermore, the ability of porphyrins to form stacking complexes with flat substrates and intercalation species with nucleic acids together with their photoactivity, confer to receptors 1-3 and to their metal complexes properties of cyclointercalands, potentially capable of performing photochemical reactions on selectively bound substrates, in particular photocleavage of DNA. Studies along these lines are in progress.

Experimental Section

Reagents and Solvents. All solvents and reagents were of reagent grade quality, purchased commercially, and used without further purification, except as noted below. Dichloromethane and THF were refluxed with and distilled from CaH2 under argon. Benzene and toluene were distilled from sodium metal under argon. Pyridine was refluxed with and distilled from KOH pellets and stored over 4 Å molecular sieves under an argon atmosphere. Benzyl chloroformate was distilled at reduced pressure and stored at -70 °C. The [18]-N₂O₄ macrocycle, 12 (1,10-diaza-4,7,13,16-tetraoxacyclooctadecane), was obtained from Merck, Inc. Merck type 60 (TLC grade) silica gel or Merck activity III neutral alumina were used for column chromatography. Thin-layer chromatography was performed with commercially prepared, plasticbacked, precoated alumina or silica plates.

Physical and Spectroscopic Methods. Melting points were obtained on a Thomas-Hoover capillary apparatus and are uncorrected. Electronic spectra were recorded on either a Cary 219 or a Cary 118 spectrophotometer. Proton NMR spectra was recorded on either a Varian EM 360A (60 MHz), Bruker SY 200 (200 MHz), or Bruker WP 400 (400 MHz) spectrometer. Carbon-13 NMR spectra were recorded at 50 MHz on a Bruker SY 200 spectrometer. Chemical shifts are reported in ppm vs. Me₄Si. Mass spectrometry was performed either by Chemical Ionization at the Laboratoire de Spectrométrie de Masse, Strasbourg, or by Fast Atom Bombardment at the Service Central de Microanalyse du CNRS, Lyon. Microanalyses were obtained at the Centre de Microanalyses du CNRS, Institut de Chimie de Strasbourg.

Techniques for Handling Unstable Compounds. The free-base porphyrins were treated as if light sensitive and handled in subdued light since similar porphyrinic compounds have been reported to be light sensitive.^{10,15,16} The final tetraamine products 1-3 were stored under argon

Biphenyl-Bridged Bis-[18]-N2O4 Macrocycle 9. Direct Procedure. To a solution of the [18]-N₂O₄ macrocycle 12 (3 g, 11.5 mmol) and triethylamine (3 mL) in CH₂Cl₂ (25 mL), a solution of 4,4'-dichloroformylbiphenyl (13) (1 g, 3.57 mmol) in CH₂Cl₂ (100 mL) was added dropwise over 2 h with vigorous stirring. The mixture was stirred for an additional 16 h. Toluene (50 mL) was then added, and the solution was reduced in volume to 50 mL on a rotary evaporator. The organic solution was extracted with three 100-mL portions of water. The aqueous washings were combined and extracted with CH_2Cl_2 (3 × 100 mL). The dichloromethane solutions were combined, dried over Na₂SO₄, and evaporated to dryness to yield the diamine diamide 9 as an oil (2.2 g, 85% evaporation of the solvent afforded the macrotricyclic tetraamide 25 (0.43 g).¹⁷ Basification of the final aqueous phase to a U to a standard to the final aqueous phase to a U to a standard to the final aqueous phase to a U to a standard to the final aqueous phase to a U to a standard to the final aqueous phase to a U to a standard to the final aqueous phase to a U to a standard to the final aqueous phase to a U to a standard to the final aqueous phase to a U to a standard to the final aqueous phase to a standard to the final aqueous phase to a U to a standard to the final aqueous phase to a sta Basification of the final aqueous phase to pH 10 with 50% aqueous NaOH, extraction into CH₂Cl₂, drying with MgSO₄, and evaporation gave the starting [18]-N₂O₄ compound **12** (0.52 g). For compound **9**: ¹H NMR (CDCl₃) 2.85 (m, 8 H, CH_2 -NH), 3.20 (m, 2 H, NH), 3.70 (m, 40 H, CH₂–O, CH₂–NCO), 7.60 (dd, J = 14, 10 Hz, 8 H, aromatic

Biphenyl-Bridged Bis[18]-N2O4 Macrocycle 9. Indirect Procedure. Step 1, Preparation of 15. To a solution of 4.4'-dichloroformylbiphenyl (0.37 g, 1.32 mmol) in CH_2Cl_2 (30 mL) was added a solution f the N-benzyloylcarbonyl monoprotected [18]-N₂O₄ macrocycle 14¹² (1.04 g, 2.62 mmol) in CH₂Cl₂ (60 mL) and triethylamine (0.36 mL) over a period of 4 h with vigorous stirring. After the reaction mixture was washed with 100 mL of 10% aqueous NaOH, the organic layer was dried over Na₂SO₄ and evaporated to dryness to yield the diprotected diamide 15 as an oil (1.31 g, quantitative). This product was used directly without further purification. ¹H NMR (CDCl₃) 3.62 (m, 48 H, C H_2 -O, C H_2 -N), 5.11 (s, 4 H, Ar-C H_2), 7.40 (m, 18 H, aromatic H). Step 2, Preparation of 9. The diprotected diamide 15 (1.31 g, 1.31 mmol) in CH_3OH (150 mL) and palladium on carbon catalyst (5%, 0.2 g) were stirred under hydrogen overnight, or until no further hydrogen uptake was observed. The reaction mixture was filtered through Celite and evaporated to dryness. The crude product was dissolved in CH₂Cl₂:C- H_3OH 1:1 (v/v) and subject to a second filtration through a small plug of alumina. The diamide diamine 9, identical to that obtained above, was isolated as an oil (0.95 g, quantitative).

Biphenyl-C2-porphyrin Bis-[18]-N2O4 Macrocyclic Tetraamide 10. Dry pyridine (600 mL) was placed in a 1 L round-bottomed flask equipped with a Vigreux column and an argon inlet, and the column was warmed to 55 °C. Solutions of the *p*-nitrophenyl ester 6¹⁰ (336 mg, 0.43 mmol) and diamide diamine 9 (292 mg, 0.40 mmol) in dry pyridine (50 mL $\,$ each) were made up and quickly poured into the warm pyridine. The reaction was stirred for 2 days at 55 °C under argon in the dark. The pyridine was then removed in vacuo, and the residue was taken up in CHCl₃ and washed with 5% aqueous NaOH to remove *p*-nitrophenol. The aqueous phase was further extracted with $CHCl_3$ (3 × 100 mL). The combined CHCl₃ solutions were dried over Na₂SO₄, and the solvent was removed on a rotary evaporator to yield the crude product, which is contaminated with an unidentified porphyrinic impurity of lower R_f (silica, 5% CH₃OH in CHCl₃ eluent). It is nonetheless of sufficient purity that it may be used directly for the next step. When necessary, chromatography was carried out on a 1.5×40 cm column of silica gel with 2.5% CH₃OH in CH₂Cl₂ as the eluent. The first major band was collected and recrystallized from CH_2Cl_2 -hexane to yield 285 mg (54.6%) of the analytically pure tetraamide 10. ¹³C NMR (CDCl₃): 11.3, 11.9 (porphyrin–CH₃), 17.4 (CH₂–CH₃), 19.6 (CH₂–CH₃), 34.8 (CH₂–CO), 49.4, 49.7 (N–CH₂–CH₂–O), 70.2, 70.5, 70.8, 71.0 (N–C-H₂-CH₂-O), 96.3, 97.6 (meso C), 126.6, 126.8, (biphenyl C₂, C₃, C₅, C₆) 13.9, 134.6, 135.4, 138.7 (pyrrole β C), 138.4, 139.3, 149.1, 149.9 (pyrrole α C), 140.2, 140.4 (biphenyl C₁, C₄) 171.3, 172.3 (CO). Anal. Calcd for C₇₀H₈₈N₈O₁₂: C, 68.16; H, 7.19; N, 9.08. Found: C, 68.19; H, 7.31; N, 8.83. m/e 1233 (calcd 1233).

Biphenyl-C₃-porphyrin Bis-[18]-N₂O₄ Macrocyclic Tetraamide 11 was prepared in 50% yield from the *p*-nitrophenyl ester 7^{10} (261.8 mg, 0.32 mmol) and the diamine diamide 9 (251.7 mg, 0.34 mmol) with use of the same synthesis and purification procedure as above. Again, the crude product was generally sufficiently pure to be used directly for further work. m/e 1261 (calcd 1261).

Biphenyl-C2-porphyrin Bis-[18]-N2O4 Macrocyclic Tetraamine 1. The crude tetraamide 10 (165 mg, 0.13 mmol) was dissolved in 80 mL of boiling CH₂Cl₂. A solution of zinc acetate (0.5 g) in CH₃OH (10 mL) was then added, and the mixture was stirred at 40 °C for 5 min. At this

⁽²⁶⁾ When substrate S_8^{2+} for which appreciable but not optimal binding is to be expected (Table III) is added to either 3 or 22, complexes are obtained which are significantly more soluble than those obtained with S_9^{2+} or S_{10}^{2+} . With either the S_7^{2+} or S_{11}^{2+} substrates there is no evidence of complexation and the solution remains free of precipitate. (27) Gubelmann, M.; Lehn, J. M., unpublished results.

point it was confirmed by visible spectroscopy that the zinc insertion was complete (only bands at 537 and 575 nm were observed in the 700-450-nm range). The reaction mixture was then poured into 100 mL of H₂O. The organic layer was separated off, and the aqueous layer was washed with more CH_2Cl_2 (3 × 30 mL). The organic layers were combined, dried over Na2SO4, and evaporated to dryness. After being flushed with argon, the red residue was treated with diborane (30 mL of a 1 N solution in THF) and heated at reflux for 1 h. After the mixture was cooled, H₂O was added to quench the excess diborane and the solution was taken to dryness on a rotary evaporator. The solids were treated with 8 N HCl, and the resulting solution was heated for 8 h at reflux. After being allowed to cool, the solution was made basic by the addition of 1 N aqueous LiOH and extracted with CH_2Cl_2 (3 × 50 mL). The CH_2Cl_2 extracts were combined, dried over MgSO4, and evaporated to dryness. The crude yield was virtually quantitative. Purification involved chromatography on alumina (1.5×40 cm column) with 2.75% CH₃OH in CH₂Cl₂ as the eluent. Collection of the major band followed by recrystallization from CH₂Cl₂-hexane yielded 115 mg (73% based on 10, 54% based on 9) of the analytically pure tetraamine 1. ¹H NMR (CDCl₃): -3.75 (s, 2 H, pyrrole), 1.79 (br, 8 H, macrocyclic N-CH₂), 1.82 (t, 6 H, CH₂-CH₃), 2.18 (br, 4 H, N-CH₂-biphenyl), 2.85 (br, 8 H, macrocyclic N-CH₂), 3.25 (br, 16 H, macrocyclic O-CH₂), 3.54 (s, 12 H, CH₃), 3.59 (br, 4 H, N-CH₂-CH₂-porphyrin), 3.61 (s, 8 H, macrocyclic O-CH₂), 3.86 (t, 8 H, macrocyclic O-CH₂), 4.11 (q, 4 H, CH₂-CH₃), 4.22 (br, 4 H, N-CH₂-CH₂-porphyrin), 5.90 (br, 4 H, aromatic H), 6.41 (d, 4 H, aromatic H), 10.02 (s, 2 H, meso H), 10.09 (s, 2 H, meso H). ¹³C NMR (CDCl₃): 11.6, 12.0 (CH₃), 17.6 (CH₂-CH₃), 19.6 (CH₂-CH₃), 24.4 (porphyrin-CH₂-CH₂), 52.6, 54.0 (N-C-H₂-CH₂-O), 56.0 (porphyrin-CH₂-CH₂-N), 57.2 (biphenyl-CH₂-N), 68.0, 69.2, 69.3, 70.0 (N-CH₂-CH₂-0), 94.8, 95.2 (meso C), 123.6, 136.4 (biphenyl C₂, C₃, C₅, C₆), 124.0, 127.0 (pyrole β C), 132.8, 134.3, 136.7 136.7, 139.8 (pyrrole α C), 135.6, 136.0 (biphenyl C₁, C₄). λ_{max} 397 nm. Anal. Calcd for C₇₀H₉₆N₈O₈: C, 71.39; H, 8.22; N, 9.52. Found: C, 71.13; H, 8.14; N, 9.37. m/e 1177 (calcd 1177).

Biphenyl-C3-porphyrin Bis-[18]-N2O4 Macrocyclic Tetraamine 2. With use of the same procedure, the tetraamide 11 was reduced to the tetraamine 2. Purified 11 (550 mg, 0.44 mmol) gave analytically pure 2 (480 mg) in 91% yield after chromatography and recrystallization as above. ¹H NMR (CDCl₃): -3.69 (s, 2 H, pyrrole), 1.82 (t, 6 H, CH₂-CH₃), 2.06 (br, 8 H, macrocyclic N-CH₂), 2.53 (m, 4 H, CH₂-CH₂-CH₂), 2.78 (s, 4 H, N-CH2-biphenyl), 2.85 (br, 8 H, macrocyclic N-CH2), 3.01 (br, 8 H, macrocyclic O-CH₂), 3.25 (br, 8 H, macrocyclic eo-CH₂), 3.46 (br, 8 H, macrocyclic O-CH₂), 3.53 (m, 4 H, CH₂-CH₂-CH₂-N), 3.61 (t, 8 H, macrocyclic O-CH₂), 3.62 (s, 6 H, CH₃), 3.64 (s, 6 H, CH₃), 4.05 (q, 4 H, CH₂-CH₃), 4.15 (t, 4 H, porphyrin-CH₂-CH₂-CH₂), 6.41 (m, 4 H, aromatic H), 6.66 (d, 4 H, aromatic H), 10.06 (s, 2 H, meso H), 10.13 (s, 2 H, meso H). ¹³C NMR (CDCl₃): 11.7, 12.1 (CH₃), 17.8 (CH₂-CH₃), 20.0 (CH₂-CH₃), 24.7 (porphyrin-CH₂-CH₂-CH₂), 31.3 (CH2-CH2-CH2), 53.9, 54.8 (N-CH2-CH2-O), 56.0 (CH2-CH2-C-H₂-N), 59.1 (biphenyl-CH₂-N), 69.6, 70.4, 70.6, 71.0 (N-CH₂CH₂-O), 96.5, 96.8 (meso C), 125.9, 128.6 (biphenyl C₂, C₃, C₅, C₆), 126.9, 129.4 (pyrrole β C), 135.2, 135.9, 140.7, 142.3 (pyrrole α C), 138.1, 138.7 (biphenyl C₁, C₄). λ_{max} 397 nm. Anal. Calcd for C₇₂H₁₀₀N₈O₈: C, 71.73; H, 9.29; N, 8.36. Found: C, 71.66; H, 9.10; N, 8.22.

Bis-C₂-porphyrin Bis-[18]-N₂O₄ Macrocyclic Tetraamide 16. Direct Procedure. The *p*-nitrophenyl ester 6^{10} (83.9 mg, 1.1 mmol) and the [18]-N₂O₄ macrocycle 12 (26.3 mg, 1.0 mmol) were heated together for 48 h in pyridine (200 mL) at 55 °C in the dark under argon. The pyridine was then removed in vacuo, and the residues were taken up in CHCl₃ (ca. 400 mL) and washed with 5% aqueous NaOH. After being dried over Na₂SO₄, the solvent was removed on a rotary evaporator to yield the very crude tetraamide product. Chromatography on TLC grade silica (1.5 × 40 cm column) eluting with 2.75% CH₃OH in CHCl₃ gave, after collection of the first major band and subsequent recrystallization from CH₂Cl₂-hexane, the purified tetraamide 16 (10 mg, 12%). *m/e* 1529 (calcd 1529).

C₂-Porphyrin-Bridged Bis-(M-CBz protected [18]-N₂O₄) Macrocycle 17. The C₂ p-nitrophenyl ester 6¹⁰ (233 mg, 0.30 mmol) and the monoprotected [18]-N₂O₄ macrocycycle 14¹² (236 mg, 0.60 mmol) were added to 400 mL of dry pyridine and stirred for 48 h at 55 °C in the dark under an argon atmosphere. The pyridine was then removed in vacuo, and the residue was taken up in CHCl₃ and washed with 5% aqueous NaOH. After being dried over Na₂SO₄ and evaporated to dryness on a rotary evaporatory, the crude product was purified by chromatography on TLC grade silica with first 1% CH₃OH in CH₂Cl₂ and then 2% CH₃OH in CH₂Cl₂-hexane to yield the analytically pure product 17 (250 mg, 66%). ¹H NMR (CDCl₃): 1.85 (t, 6 H, CH₂-CH₃), 3.09 (m, 8 H, macrocyclic CH₂), 3.21 (m, 8 H, macrocyclic CH₂), 3.29 (m, 8 H, macrocyclic CH₂), 3.33 (m, 8 H, macrocyclic CH₂), 3.39 (m, 8 H, macrocyclic CH₂), 3.64 (m, 16 H, macrocyclic O–CH₂), 3.94 (br, 4 H, CH₂–CON), 4.15 (q, 4 H, CH₂–CH₃), 5.13 (s, 4 H, O–CH₂–phenyl), 7.38 (s, 10 H, aromatic H), 10.12 (s, 2 H, meso H), 10.46 (s, 2 H, meso H). (No peaks ascribable to the pyrrole N–H signal were observed for this compound.) Anal. Calcd for $C_{72}H_{94}N_8O_{14}$: C, 66.75; H, 7.31; N, 8.64. Found: C, 66.36; H, 7.22; N, 8.50.

C2-Porphyrin-Bridged Bis-[18]-N2O4 Macrocyclic Diamine Diamide 18. The diprotected porphyrin-bridged compound 17 (400 mg, 0.31 mmol) was dissolved at room temperature in 10 mL of concentrated HCl. If necessary, trifluoroacetic acid (ca. 5 mL) was then added to solubilize the porphyrin. The solution was stirred for several hours while the progress of the reaction was closely monitored by TLC. This was done by removing small aliquots from the reaction mixture, basifying with aqueous LiOH, extracting into CHCl₃, and drying over MgSO₄. When the spot at $R_f 0.30$ became dominant (and before spots with lower R_f values began to appear), the reaction was quenched by the slow addition of 1 N aqueous LiOH. After the solution had reached pH 10, it was extracted with CHCl₃ ($3 \times 100 \text{ mL}$). The CHCl₃ extracts were combined, dried over MgSO₄, and taken to dryness in vacuo to yield 313 mg (98%) of the crude diamine diamide 18. In general, the crude material is of sufficient purity that it may be used directly in the subsequent macrocycle forming reaction. As necessary, however, chromatography on activity III alumina, with 1% CH₃OH in CHCl₃, as eluent, was used to obtain pure material. ¹H NMR (CDCl₃): -3.81 (s, 2 H, pyrrole), 1.87 (t, 6 H, CH₂-CH₃), 1.99 (s, 2 H, macrocyclic NH), 2.34 (m, 8 H, macrocyclic N-CH₂), 2.96 (s, 8 H, macrocyclic CH₂), 3.08 (m, 8 H, macrocyclic CH₂), 3.29 (s, 6 H, CH₃), 3.30 (s, 6 H, CH₃), 3.42 (s, 8 H, macrocyclic CH₂), 3.64 (t, 16 H, macrocyclic O-CH₂), 3.93 (m, 4 H, CH2-CON), 4.15 (q, 4 H, CH2-CH3), 10.01 (s, 2 H, meso H), 10.38 (s, 2 H, meso H). ¹³C NMR (CDCl₃): 12.2, 13.2 (CH₃), 18.7 (CH₂-CH₃), 21.1 (CH₂-CH₃), 35.44 (CH₂-CON), 47.8, 50.1 (N-CH₂-CH₂-O), 70.7, 71.3, 71.4 (N-CH2-CH2-O), 97.5, 98.7, (meso C), 134.8, 135.6, 139.6, 141.4 (pyrrole β C), 145.4, 147.2, 149.2, 150.8 (pyrrole α C), 171.1 (CO). Anal. Calcd for $C_{56}H_{82}N_8O_{10}$ -1.5H₂O: C, 63.97; H, 7.86; N, 10.65. Found: C, 63.90; H, 7.90; N, 10.56. m/e 1026 (calcd 1026).

Bis-C₂-porphyrin Bis-[18]-N₂O₄ Macropentacyclic Tetraamide 16. Stepwise Procedure. The diamine diamide 18 (138 mg, 0.11 mmol) was mixed with the *p*-nitrophenyl ester 6 (110 mg, 0.11 mmol) in pyridine (300 mL) and stirred for 48 h at 55 °C in the dark under an argon atmosphere. The pyridine was then removed in vacuo, and the residue was taken up in CHCl₃ (300 mL). After being washed with 1 N aqueous LiOH (2 × 100 mL), the organic phase was dried over Na₂SO₄ and taken to dryness on a rotary evaporator to yield the crude tetraamide 16 (93 mg, 45%). The major migrating spot of this crude product proved by TLC (silica, 5% CH₃OH in CHCl₃) to be identical with the purified macrocyclic material obtained by the direct procedure. A small scale chromatographic separation (silica, 2.75% CH₃OH in CHCl₃ eluent) yielded purified tetraamide 16 identical with that prepared by the direct procedure. m/e 1529 (calcd 1529).

Bis-C₂-porphyrin Bis-[18]-N₂O₄ Macropentacyclic Tetraamine (3). The crude tetraamide 16 (180 mg, 0.12 mmol), prepared from 18, was dissolved in 100 mL of CHCl₃, and a solution of 0.5 g of zinc acetate in CH₃OH (10 mL) was added. The solution was heated at reflux for 0.5 h, the resulting pink solution was poured into H_2O , and the metalloporphyrin complex was extracted into CHCl₃. After the organic phase was dried over Na₂SO₄, the solvent was removed in vacuo and the flask was flushed with argon. Diborane (25 mL of a 1 N solution in THF) was added, and the resulting solution was heated at reflux for 8 h. After the mixture was cooled, excess diborane was quenched cautiously with water. The reaction mixture was then taken to dryness on a rotary evaporator. Aqueous 8 N hydrochloric acid (50 mL) was added to the solid residue, and the resulting mixture was heated at reflux for 8 h. After being cooled, the solution was made basic to pH 10 or 11 by the addition of 1 N aqueous LiOH and extracted with $CHCl_3$ (3 × 100 mL). The organic extracts were combined, dried over MgSO4, and taken to dryness on a rotary evaporator. The resulting crude product was chromatographed on alumina, eluting with 1% CH₃OH in CHCl₃, to yield, after recrystallization from CH₂Cl₂-hexane, 150 mg of the tetraamine 3 (86.5% based on 16, 25% based on 6). By the same procedure, compound 3 can also be prepared in 10% yield (based on 6) from the crude tetraamide 16 obtained by the direct condensation procedure. In this case, at least two chromatographic separations on alumina are required before pure product is obtained. ¹H NMR (CDCl₃): -4.39 (s, 4 H, pyrrole), 1.71 (br, 8 H, N- CH_2 - CH_2 -porphyrin), 1.90 (t, 12 H, CH_2 - CH_3), 2.25 (br, 16 H, macrocyclic N- CH_2), 2.72 (br, 8 H, N- CH_2 -CH2-porphyrin), 2.85 (s, 12 H, CH3), 3.04 (s, 12 H, CH3), 3.45 (br, 16 H, macrocyclic O-CH₂), 3.59 (s, 16 H, macrocyclic O-CH₂), 4.10 (q, 8 H, CH₂-CH₃), 8.30 (br, 4 H, meso H), 9.76 (s, 4 H, meso H). ¹³C NMR (CDCl₃): 10.9, 11.1 (CH₃), 17.9 (CH₂-CH₃), 20.1 (porphyrin-CH2-CH3), 21.3 (porphyrin-CH2-CH2-N), 53.6 (N-CH2-CH2-O), 55.8 (porphyrin-CH₂-CH₂-N), 70.8, 71.2 (N-CH₂-CH₂O), 95.5, 95.7 (meso C), 134.3, 135.2, 136.8, 141.9 (pyrrole \$\beta\$ C), 143.0, 145.1 (pyrrole α C). λ_{max} 388 nm. Anal. Calcd for C₈₈H₁₂₀N₁₂O₈: C, 71.71; H, 8.21; N, 11.40. Found: C, 71.90; H, 8.12; N, 11.39. *m/e* 1473 (calcd 1473).

Zn Biphenyl-C2-porphyrin Bis-[18]-N2O4 Macrotetracyclic Complex 19. The free-base porphyrin 1 (10 mg, 0.0085 mmol) was dissolved in 10 mL of hot CH₂Cl₂, and 0.89 mL of a 0.010 N solution of zinc acetate in CH₃OH was added. When, as judged by the absence of bands at 497 nm 619 nm in the visible spectrum, the reaction was considered complete, the pink solution was poured into a very dilute aqueous solution of LiOH. The organic layer was separated off, and the aqueous phase was extracted with $CHCl_3$ (2 × 20 mL). The organic phases were combined, dried over $MgSO_4,$ and taken to dryness in vacuo to yield 19 (10.2 mg, 97%). $^1\mathrm{H}$ NMR (CDCl₃): no peaks associated with the pyrrole N-H protons are observed; other features were similar to those observed for the free-base 1. λ_{max} 405 nm.

Zn Biphenyl-C3-porphyrin Bis-[18]-N2O4 Macrotetracyclic Complex 20 was prepared from 2 by the same procedure used to prepare 19 from 1. For 10 mg (0.0083 mmol) of the free-base, 0.90 mL of the stock 0.010 N zinc acetate solution was used to give 10 mg (95%) of the zinc complex 20. ¹H NMR (CDCl₃): no signals ascribable to a pyrrolic N-H are detected; other features much as for the free-base. λ_{max} 402 nm.

Zn₂ Bis-C₂-porphyrin Bis-[18]-N₂O₄ Dinuclear Macropentacyclic Complex 22 was prepared in 97% yield from 3 by using a similar procedure. For 10 mg of the free-base 3, 1.42 mL of the 0.010 N zinc acetate stock solution was used to give 10.5 mg of the bis-zinc complex 22. ¹H NMR (CDCl₃): no signals ascribable to a pyrrolic N-H were observed; other features much as for the free-base. λ_{max} 385 nm.

Cu Biphenyl-C2-porphyrin Bis-[18]-N2O4 Macrotetracyclic Complex 21. The purified free-base compound, 1 (50 mg, 0.043 mmol), was dissolved in CHCl₃ (20 mL) at 40 °C, and 5 mL of a saturated solution of copper acetate in CH₃OH were added. After approximately 10 min, when visible spectroscopy indicated the absence of starting material (no bands were observed at 497 and 619 nm), the reaction mixture was poured into a 0.1 N aqueous solution of NaCN, made slightly basic by the addition of LiOH. The organic layer was extracted out, washed with water (3 \times

20 mL), dried over Na₂SO₄, and stripped to dryness on a rotary evaporator. Recrystallization from CH2Cl2-hexane yielded the copper complex **21** as large purple crystals (50 mg, 95%). λ_{max} 399 nm. Anal. Calcd for $C_{70}H_{99}N_8O_8Cu\cdot 2H_2O$: C, 65.94; H, 7.75. Found: C, 66.24; H, 7.81. This compound was also characterized by an X-ray crystallographic structure determination.²⁰

Cu₂ Bis-C₂-porphyrin Bis-[18]-N₂O₄ Dinuclear Macropentacyclic Complex 23 was prepared in an identical manner as above from the free-base 3 in 92% yield. λ_{max} 390 nm. The crystals obtained were kept in view of an X-ray crystallographic analysis. Anal. Calcd for C₈₈H₁₁₆N₁₂O₈Cu₂·5H₂O: C, 62.65; H, 7.53; N, 9.96. Found: C, 62.69; H, 7.22; N, 9.62.

Acknowledgment. The authors thank Merck A.G., Darmstadt, for the gift of the [18]-N₂O₄ macrocycle 12 used in this work. A.D.H. and J.L.S. thank both the NATO and NSF-CNRS scientific exchange programmes for postdoctoral research fellowships.

90633-76-4; 10 (Zn complex), 103201-95-2; 11, 90633-77-5; 12, 23978-55-4; 13, 2351-37-3; 14, 42031-79-8; 15, 90633-73-1; 16, 90653-20-6; 16 55-4; 13, 2351-37-3; 14, 42031-79-8; 15, 90633-73-1; 16, 90653-20-6; 16 (Zn complex), 103201-96-3; 17, 90633-78-6; 18, 90653-19-3; 19, 90603-80-8; 19.⁺H₃N(CH₂)₈NH₃⁺, 90818-88-5; 19.⁺H₃N(CH₂)₉NH₃⁺, 90818-90-9; 19.⁺H₃N(CH₂)₁₀NH₃⁺, 90818-89-5; 20, 90603-81-9; 20.⁺H₃N(CH₂)₈NH₃⁺, 90822-43-8; 20.⁺H₃N(CH₂)₉NH₃⁺, 90818-90-9; 20.⁺H₃N(CH₂)₁₀NH₃⁺, 90818-91-8; 21, 103201-97-4; 22, 90603-82-0; 22.⁺H₃N(CH₂)₁₀NH₃⁺, 90818-92-1; 22.⁺H₃N(CH₂)₉NH₃⁺, 90818-93-2; 22.⁺H₃N(CH₂)₁₀NH₃⁺, 90818-94-3; 23, 103201-98-5; ⁺H₃N-(CH₂)₈NH₃⁺, 49745-06-4; ⁺H₃N(CH₂)₉NH₃⁺, 103201-88-3; ⁺H₃N-(CH₂)₁₀NH₃⁺, 49745-07-5.

The Mechanism of Nickel-Catalyzed Ethylene Hydrocyanation. Reductive Elimination by an Associative Process

Ronald J. McKinney* and D. Christopher Roe*

Contribution No. 3765 from the Central Research and Development Department, E. I. du Pont de Nemours and Company, Experimental Station, Wilmington, Delaware 19898. Received February 13, 1986

Abstract: The complex $(C_2H_4)L(CN)(C_2H_5)Ni^{11}$ [L = P(O-o-tolyl)₃] has been identified at -40 °C by ¹H, ³¹P, and ¹³C NMR spectroscopy as the primary intermediate in the catalytic hydrocyanation of ethylene. Reaction of this intermediate with L causes reductive elimination of propanenitrile and produces $(C_2H_4)L_2N_i$ which reacts with ethylene and hydrogen cyanide to regenerate the intermediate. Measurements of second-order rate constants at -50 to -10 °C result in $\Delta H^* = 8.8 \pm 0.9$ kcal/mol and $\Delta S^* = -34 \pm 4$ eu ($\Delta G^* = 16.7 \pm 0.1$ kcal/mol at -40 °C). At higher L concentrations, these nickel species are also in equilibrium with $(C_2H_4)L_3Ni$, L_4Ni , and $HNi(CN)L_3$ which remove nickel from the productive catalytic cycle. Equilibrium constants relating these species and corresponding thermodynamic parameters have been determined. The rates of dissociation of L from L₄Ni and HNi(CN)L₃ have also been determined. The former reaction is very slow with $\Delta H^* = 20.1 \pm 1.6$ kcal/mol and $\Delta S^* = 7 \pm 7$ eu ($\Delta G^* = 18.5$ kcal/mol at -40 °C) whereas the latter is about 10³ faster with ΔH^* = 18.8 ± 2.7 kcal/mol and $\Delta S^* = 17 \pm 10$ eu ($\Delta G^* = 14.8$ kcal/mol at -40 °C).

The development of the Du Pont adiponitrile process,¹ i.e., the dihydrocyanation of butadiene, has provided on the one hand a remarkable example of industrial application of homogeneous catalysis and on the other an excellent opportunity for detailed mechanistic studies of a complex practical system. A series of recent reports from our laboratories.²⁻¹¹ as well as others,¹²⁻¹⁷ is developing a general picture of nickel-catalyzed olefin hydro-cyanation.^{18,19} However, understanding of many of the funda-

5167

Chem. Eng. New 1971, 49, 30.
 (2) (a) Tolman, C. A.; Seidel, W. C.; Gosser, L. W. J. Am. Chem. Soc.
 1974, 96, 53. (b) Gosser, L. W.; Tolman, C. A. Inorg. Chem. 1970, 9, 2350.

⁽³⁾ Druliner, J. D.; English, A. D.; Jesson, J. P.; Meakin, P.; Tolman, C.

⁽a) Dramer, J. D., English, A. D., Sesson, J. F., Meakin, F., Tohnan, C. A. J. Am. Chem. Soc. 1976, 98, 2156.
(4) Tolman, C. A.; Seidel, W. C. J. Am. Chem. Soc. 1974, 96, 2774.
(5) Tolman, C. A. J. Am. Chem. Soc. 1974, 96, 2780.
(6) Tolman, C. A. Organometallics 1983, 2, 614.
(7) Tolman, C. A.; Seidel, W. C.; Gosser, L. W. Organometallics 1983, 2, 2001. 2, 1391

⁽⁸⁾ Tolman, C. A. Inorg. Chem. 1971, 10, 1540.