

S_{RN}1 Substitutions of Halogenoperfluoroalkanes (CF₃Br or CF₂Cl₂) under Pressure

Claude Wakselman* and Marc Tordeux


CNRS-CERCOA, 2 à 8 rue Henri Dunant, 94320 Thiais, France

Radical chain fluoroalkylation of arenethiolates by CF₃Br or CF₂Cl₂, performed under slight pressure in a glass apparatus, gives aryl polyfluoromethyl sulphides.



Halogenoperfluoroalkanes (Freons) are noted for their inertness.¹ In the case of CF₃Br and CF₂Cl₂, some reactivity has been observed when powerful nucleophilic reagents such as organolithium derivatives are used. These condensations often lead to partial (or even total) decomposition of the perfluoroalkyl group.^{2,3†} We describe here the reactivity of these perhalogenomethanes with arenethiolates under special conditions.

The gaseous trifluoromethyl bromide does not react if it is simply bubbled through a solution of arenethiolates in dimethylformamide (DMF). We obtained only very small amounts of the sulphides (**1**) by carrying out the reaction in an autoclave at 80 °C. We report here that the condensation can be achieved in glass apparatus at room temperature when a pressure of about 2 atm is used, and under these conditions the trifluoromethyl aryl sulphides (**1**) were obtained in 62% yield

for Ar = Ph, 75% for Ar = *p*-MeC₆H₄, and 83% for Ar = *p*-MeOC₆H₄ [reaction (1)].



Inhibition experiments with nitrobenzene (0.025–0.1 mol of inhibitor for 0.1 mol of arenethiolate in 100 ml of DMF) are clear cut‡ and show that a radical chain mechanism, similar to the one described by Kornblum and Bunnett for benzyl and aryl halides, is involved.⁴ The chain process can be written as in reactions (2)–(5).




† It has been possible to introduce a fluoroalkyl group (–CF₂Br) from the more reactive CF₃Br₂ or CF₂BrCl by capture of the intermediate difluorocarbene (ref. 7). The introduction of a halogenoperfluoroalkyl group can also be achieved with these reagents through a radical chain process similar to the one observed with iodoperfluoroalkanes (D. Cantacuzene, C. Wakselman, and R. Dorme. *J. Chem. Soc., Perkin Trans. 1*, 1977, 1365) as shown in the condensation with uncharged nucleophiles (I. Rico, D. Cantacuzene, and C. Wakselman, *Tetrahedron Lett.*, 1981, **22**, 3405).

‡ We have also observed a clear-cut inhibition by nitrobenzene in the condensation of benzenethiolate with perfluorohexyl bromide in DMF at atmospheric pressure. The inhibition by nitrobenzene of the similar condensation with perfluorohexyl iodide is not so efficient. However in that case inhibition by styrene has been reported recently (A. E. Feiring, *J. Fluorine Chem.*, 1984, **24**, 191).

The initiation step is likely to be the electron transfer (2) from the arenethiolate to the trifluoromethyl bromide. Radical anions $[\text{CF}_3\text{X}]^\cdot$ (with $\text{X} = \text{Cl}, \text{Br}, \text{or I}$) have been observed as stable species by e.s.r. spectroscopy at very low temperature; their breakdown to give a trifluoromethyl radical and a halide ion has also been followed by e.s.r.⁵ The slight pressure used for the condensation can easily be obtained in a Parr apparatus. This pressure seems to increase the concentration of CF_3Br in the solution and the chain process can occur.

Therefore, the use of the expensive CF_3I and u.v. irradiation⁶ is not necessary for the trifluoromethylation of arenethiolates.

The behaviour of CF_2Cl_2 has also been examined under these conditions. § The major compound obtained is the chlorodifluoromethyl aryl sulphide (2) [yield: (2) 62%, (3) 8%, (4) 7% with PhSK; yield: (2) 44%, (3) 8%, (4) 6% with *p*-MeC₆H₄SK; reaction (6)].



Inhibition of the substitution by nitrobenzene shows that the mechanism with CF_2Cl_2 is similar to the one proposed for CF_3Br . The radical anion $[\text{CF}_2\text{CL}_2]^\cdot$ has also been observed.⁵ The formation of the side product (4) can be explained by the decomposition of the radical anion $[\text{ArSCF}_2\text{Cl}]^\cdot$ to the radical ArSCF_2^\cdot which can react further with the arenethiolate anion.⁷

§ We have observed previously a partial condensation of CF_2Cl_2 with benzenethiolate under u.v. irradiation; however, the main product obtained was the disubstituted product (4) (ref. 7).

The condensation of arenethiolates with CF_3Br or CF_2Cl_2 gives aryl polyfluoromethyl sulphides in fair to good yields. The method used is cheap and can be applied to polyhalogenosulphides in the agrochemical field,⁸ the properties of which are due to their high lipophilicity.⁹

This type of condensation is the first example of monosubstitution of these poorly reactive reagents by a radical chain process.

We thank Rhône-Poulenc for financial support.

Received, 10th February 1984; Com. 184

References

- 1 R. E. Banks, 'Organofluorine Chemicals and their Industrial Applications,' Soc. Chem. Ind., London, 1979, p. 68, 89.
- 2 V. Franzen and L. Fikentscher, *Chem. Ber.*, 1962, **95**, 1958; M. Schlosser, Le Van Chau, and B. Spahic, *Helv. Chim. Acta*, 1975, **58**, 2575.
- 3 The recent synthesis of trifluoromethylphosphanes is an exception explained by an ionic mechanism, W. Volbach and I. Ruppert, *Tetrahedron Lett.*, 1983, **24**, 5509.
- 4 N. Kornblum, *Angew. Chem., Int. Ed. Engl.*, 1975, **14**, 734; J. F. Bunnett, *Acc. Chem. Res.*, 1978, **11**, 413.
- 5 A. Hasegawa, M. Shiotani, and F. Williams, *Faraday Discuss. Chem. Soc.*, 1977, **63**, 157; M. C. R. Symons, *Pure Appl. Chem.*, 1981, **53**, 223.
- 6 V. I. Popov, V. N. Boiko, and L. M. Yagupolskii, *J. Fluorine Chem.*, 1982, **21**, 365.
- 7 I. Rico, D. Cantacuzene, and C. Wakselman, *J. Org. Chem.*, 1983, **48**, 1979.
- 8 E. Kuhle and E. Klauke, *Angew. Chem., Int. Ed. Engl.*, 1977, **16**, 735.
- 9 C. Hansch, A. Leo, S. H. Unger, K. H. Kim, O. Nikaitani, and E. J. Lien, *J. Med. Chem.*, 1973, **16**, 1207.