Zwitterionic, Ring-Borylated Vanadium(III) Complexes from [Cp₂VCO] and B(C₆F₅)₃

Robert Choukroun,* Christian Lorber, and Bruno Donnadieu

Laboratoire de Chimie de Coordination, CNRS, 205 Route de Narbonne, 31077 Toulouse Cedex, France

Received December 18, 2003

The reaction of the vanadium(II) carbonyl $[Cp_2V(CO)]$ and $B(C_6F_5)_3$ resulted in formation of zwitterionic, ring-borylated vanadium(III) complexes $[(Cp)(C_5H_4B(C_6F_5)_3)V]$ (**3**) and $[(Cp)(C_5H_4B(C_6F_5)_3)V(CO)_2]$ (**4**) and of the salt $[Cp_2V(CO)_2][HB(C_6F_5)_3]$ (**5**). All were characterized by crystallography. Formation of a hydride vanadium(IV) $[(Cp)(C_5H_4B(C_6F_5)_3)VH(CO)]$ obtained by the electrophilic addition of the borane at the Cp-ring followed by redox and disproportionation reaction is suggested to account for these results.

Introduction

The reactivity of the borane $B(C_6F_5)_3$ toward organic and organometallic substrates is of current interest. New organic reactions, new catalytic processes, and new reactivity pathways have been observed for this borane.¹ In our studies of the reactivity of $B(C_6F_5)_3$ (1) with organometallic complexes of the groups 4 and 5 elements,^{2,3} in which catalytic application in olefin polymerization is well known, we have explored its reactivity with $[Cp_2Ti(CO)_2]$. The product of this reaction was the acylborane derivative $[Cp_2Ti(CO)(\eta^2-OCB(C_6F_5)_3]$.³ The preferential attack of borane 1 at the carbon atom of a carbonyl ligand is puzzling if we consider that attack at the more nucleophilic oxygen atom of the carbonyl ligand should be favored.⁴ Thus we decided to extend our investigation of the reactivity of 1 toward the readily available $[Cp_2VCO]$ (2).

Results and Discussion

The carbonyl complex [Cp₂VCO] (**2**) reacted with **1** in pentane to give a yellow-brown precipitate that contained a crystalline air-sensitive blue product. The latter, formed in low yield, was isolated by careful separation by hand. This product is a paramagnetic high-spin V^{III} complex ($\mu_{eff} = 2.85 \,\mu_B$). Its structure was shown by X-ray crystallography to be that of zwitterionic [(η^5 -Cp)(η^5 -C₅H₄B(C₆F₅)₃)V] (**3**) (Figure 1). The borane is linked to one of the cyclopentadienyl rings, and an



Figure 1. Molecular structure of **3**, showing 50% probability thermal ellipsoids and partial atom-labeling schemes. Selected bond distances (Å) and angles (deg): $V(1)\cdots F(36) 2.1568(14)$, C(5a)-B(1) 1.639(3), (Cp)-V(1) 1.923, $(Cp^B)-V(1) 1.913$, $(Cp)-V(1)-(Cp^B) 144.43$. (Cp^B) and (Cp) are the centroids of C(1a)-C(5a) and C(6a)-C(10a) rings, respectively.

ortho-fluorine atom of one perfluoro phenyl group of the borane is coordinated to the vanadium center with a V^{···} F distance of 2.1568(14) Å, a separation that is in the same range as those reported in the literature⁵ for fluorine atoms that bridge neighboring vanadium atoms (2.044–2.173 Å). The paramagnetic vanadium center is ⁵¹V NMR silent, whereas a ¹¹B NMR resonance at –14.8 ppm confirmed the presence of a tetracoordinated anion (we assume that the boron atom is not affected by the proximity of the paramagnetic vanadium center). Electrophilic substitution of a hydrogen atom of the Cp ring by **1** has been observed previously in group 4.⁶ In particular, the interaction of the bis(trimethylsilyl)acetylene complex of titanocene [Cp₂Ti(Me₃SiC₂SiMe₃)] with **1** gave a zwitterionic complex of titanium(III),

^{*} Corresponding author. E-mail: choukrou@lcc-toulouse.fr. (1) (a) Piers, W. E.; Chivers, T. Chem. Soc. Rev. 1997, 26, 345-354.
(b) Stoddard, J. M.; Shea, K. J. Organometallics 2003, 22, 1124-1131.
(c) Parks, D. J.; Piers, W. E.; Parvez, M.; Atencion, R.; Zaworotko, M. J. Organometallics 1998, 17, 1369-1377. (d) Parks, D. J.; Blackwell, J. M.; Piers, W. E. J. Org. Chem. 2000, 65, 3090-3098. (e) Mountford, A. J.; Hughes, D. L.; Lancaster, S. Chem. Commun. 2003, 2148-2149.
(f) Watson, I. D.; Yudinn, A. J. Org. Chem. 2003, 68, 5160-5167. (g) Chen, E. Y.-X.; Marks, T. J. Chem. Rev. 2000, 100, 1391. (h) Bochmann, M. J. Chem. Soc., Dalton Trans. 1996, 225. (i) Britovsek, G. J. P.; Gibson, V. C.; Wass, D. F. Angew. Chem., Int. Ed. 1999, 38, 428.

 ⁽²⁾ Wolff, F.; Choukroun, R.; Lorber, C.; Lepetit, C.; Donnadieu, B.
 Eur. J. Inorg. Chem. 2003, 628–632.

⁽³⁾ Choukroun, R.; Lorber, C.; Lepetit, C.; Donnadieu, B. Organometallics **2003**, *22*, 1995–1997.

⁽⁴⁾ Braunschweig, H.; Klinkhammer, K. W.; Koster, M.; Radacki, K. *Chem. Eur. J.* **2003**, *9*, 1030–1309.

^{(5) (}a) Fairhurst, S. A.; Hughes, D. L.; Leigh, G. J.; Snaders, J. R.; Weisner, J. *J. Chem. Soc., Dalton Trans.* **1994**, 2591–2598. (b) Bukovec, P.; Milicev, S.; Demsar, A.; Golic, L. *J. Chem. Soc., Dalton Trans.* **1981**, 1802–1806. (c) Buchholz, N.; Leimkühler, M.; Kiriazis, L.; Mattes, R. *Inorg. Chem.* **1988**, *27*, 2035–2039.



Figure 2. Molecular structure of **4**, showing 50% probability thermal ellipsoids and partial atom-labeling schemes. Selected bond distances (Å) and angles (deg): V(C1) 1.9484-(17), V-C(2) 1.9686(17), C(1)-O(1) 1.1369(19), C(2)-O(2) 1.133(2), B-C(21) 1.655(2), C(1)-V-C(2) 85.44(7), (Cp)-V(1) 1.930, (Cp^B)-V(1) 1.924, (Cp)-V(1)-(Cp^B) 139.15. (Cp^B) and (Cp) are the centroids of C(21)-C(25) and C(11)-C(15) rings, respectively.

 $[(\eta^5-Cp)(\eta^5-C_5H_4B(C_6F_5)_3)Ti]$, in which *ortho*-fluorine atoms of two C_6F_5 groups are coordinated to the titanium center in a pseudotetrahedral geometry.^{6b} The difference in the reactivity between the titanocene and vanadocene units could be the result of a steric effect: the larger atomic radius of Ti versus V may allow two fluorine atoms to coordinate at the Ti center.

Two other products were contained in the yellowbrown crude precipitate as observed by ¹H NMR spectroscopy (the characteristic cyclopentadienyl signals) as well as by the ⁵¹V and ¹¹B NMR spectra, in which two resonances were observed, respectively. A magnetic measurement made on the crude product (after careful and tedious separation of 3) indicated that it consisted of diamagnetic species. We were able to isolate separately each complex by dissolving the crude product in a small amount of toluene or THF followed by pentane diffusion. Formation of crystalline, air-sensitive complexes suitable for an X-ray structure determination were obtained from these selective crystallizations. In the crystals obtained from THF/pentane solution, a dicarbonylvanadium(III) species in which the borane is linked to one of the cyclopentadienyl rings of the vanadocene unit in the zwitterionic $[(\eta^5-Cp)(\eta^5-C_5H_4B (C_6F_5)_3)V(CO)_2$ (4) (Figure 2) was shown to be present. Crystals were obtained from toluene/pentane solution, and the X-ray structure determination established the formation of a salt, $[Cp_2V(CO)_2][HB(C_6F_5)_3]$ (5) (Figure 3).⁷ The isolated **4** often was contaminated by **5** in low



Figure 3. Molecular structure of **5**, showing 50% probability thermal ellipsoids and partial atom-labeling schemes. Selected bond distances (Å) and angles (deg): V(C1) 1.965-(3), V-C(2) 1.964(2), C(1)-O(1) 1.132(3), C(2)-O(2) 1.135-(3), (Cp)-V (or (Cp')-V) 1.901, (Cp)-V-(Cp') 138.01, B-H(1B) 1.14(2), C(1)-V-C(2) 86.63(11). (Cp) and (Cp') are the centroids of C(11)-C(15) and C(21)-C(25) rings, respectively.

ratio, as judged by ¹H NMR (0-5%), and isolated 5 often was contaminated by small amounts of 4. In another procedure, the synthesis carried out in a small amount of toluene as solvent gave 5 as an immediate orange crystalline product in high yield. The filtrate, after 2-3days at low temperature (-30 °C), afforded blue crystals of **3**. After filtration, the filtrate was layered with pentane to give a brown precipitate, which was analyzed by ¹H NMR as **4** contaminated with **5**. Both complexes 4 and 5 are diamagnetic (confirmed by Faraday balance experiments and multinuclear NMR). The ¹¹B NMR shift of 4 (-14.9 ppm) is in the same range as that of 3, as expected for a tetracoordination of the boron atom. The resonance of the ¹¹B NMR spectrum of 5 shows a doublet at -25.5 ppm (${}^{1}J_{B-H} = \hat{8}9$ Hz), in agreement with the presence of the $[HB(C_6F_5)_3]^-$ anion,⁸ which is also supported by the ¹H NMR spectrum, which shows a quadruplet at 3.63 ppm with ${}^{1}J_{BH} = 91$ Hz. The ${}^{51}V$ NMR spectra of 4 and 5 show very low field peaks at -1652 and 1662 ppm, respectively, due to the high basicity of the cationic $[V]^+$ moiety.⁹ Both compounds have two infrared C-O strechting vibrations (Nujol mull) at 2050 and 2004 cm^{-1} and 2038 and 1990 cm^{-1} for 4 and 5, respectively (to be compared with the bands at 2050 and 2010 cm⁻¹ reported for the $[Cp_2V(CO)_2]^+$ cation¹⁰).

From these different results, we can draw a scheme showing the possible formation of **3**, **4**, and **5**. All these compounds seem to arise from the same parent mol-

(10) Calderazzo, F.; Bacciarelli, S. Inorg. Chem. 1963, 2, 721-723.

^{(6) (}a) Ruwwe, J.; Erker, G.; Frölich, R. Angew. Chem., Int. Ed. Engl. **1996**, 35, 80–82. (b) Burlakow, V. V.; Troyanov, S. I.; Letov, A. V.; Strunkina, L. I.; Minacheva, M. Kh.; Furin, G. G.; Rosenthal, U.; Shur, V. B. J. Organomet. Chem. **2000**, 598, 243–247. (c) Burlakow, V. V.; Pellny, P.-M.; Arndt, P.; Baumann, W.; Spannenberg, A.; Shur, V. B.; Rosenthal, U. Chem. Commun. **2000**, 241–242. (d) Burlakow, V. V.; Arndt, P.; Baumann, W.; Spannenberg, A.; Rosenthal, U.; Letov, A. V.; Lyssenko, K. A.; Korlyukov, A. A.; Strunkina, L. I.; Minacheva, M. Kh.; Shur, V. B. Organometallics **2001**, 20, 4072–4079.

⁽⁷⁾ Compound **5** is related to the carbonyl salt [Cp₂V(CO)₂][BPh₄], whose structure has been reported with a disorder that prevents the accurate location of the atom involved. Atwood, J. L.; Rogers, R. D.; Hunter, W. E.; Floriani, C.; Fachinetti, G.; Chiesi-Villa, A. *Inorg. Chem.* **1980**, *19*, 3812–3817.

⁽⁸⁾ Temme, B.; Erker, G. J. Organomet Chem. 1995, 488, 177–182.
(9) (a) Devore, D. D.; Lichtenhan, J. D.; Takusagawa, E.; Maatta, A. J. Am. Chem. Soc. 1987, 109, 7408–7416. (b) Rehder, D. Coord. Chem. Rev. 1991, 110, 161–210. (c) Priebsch, W.; Rehder, D. Inorg. Chem. 1985, 24, 3058–3062.

Table 1. Summary of Crystal Data, Data Collection, and Structure Refinement Parameters for 3-5

	3	4	5
fw	1384.20	748.12	750.14
temperature (K)	180	180	160
wavelength (Å)	0.71073	0.71073	0.71073
cryst syst	monoclinic	monoclinic	monoclinic
space group	$P2_{1}/c$	$P2_1/c$	$P2_1/n$
a (Å)	17.010(5)	10.819(5)	14.594(5)
b (Å)	15.787 (5)	13.785(5)	16.127(5)
<i>c</i> (Å)	18.810(5)	18.532(5)	11.890(5)
β (deg)	94.282 (5)	105.417(5)	97.046(5)
$V(Å^3)$	5037(3)	2668.7(17)	2777.3(17)
Z, density (g cm ^{-3})	4, 1.825	4, 1.862	4, 1.794
F(000)	2720	1472	1480
abs coeff (mm $^{-1}$)	0.525	0.509	0.489
2θ range (deg)	3.36 - 31.79	2.91 - 30.51	2.44 - 28.13
no. of reflns collected/unique	46 805/15892	54 611/7749	26 433/6555
-	$(R_{\rm int} = 0.0483)$	$(R_{\rm int} = 0.0369)$	$(R_{\rm int} = 0.0492)$
completeness to 2θ (%)	63.58 (92.5)	61.02 (95.1)	79.8
no. of data/restraints/params	15 892/0/811	7749/0/442	6555/0/446
GOF on F^2	0.926	1.099	1.015
final R indices $(I > 2\sigma(I))$	R1 = 0.0483,	R1 = 0.0369,	R1 = 0.0369,
	wR2 = 0.1094	wR2 = 0.0951	wR2 = 0.0799
R indices (total)	R1 = 0.0936,	R1 = 0.0469,	R1 = 0.0663,
	wR2 = 0.1269	wR2 = 0.1001	wR2 = 0.0911
largest diff peak and hole (e·Å $^{-3}$)	0.370 and - 0.456	0.326 and -0.355	0.266 and -0.249

ecule. Electrophilic addition of 1 at the Cp-ring occurred and afforded the suggested intermediate vanadium(IV) hydride $[(\eta^5-Cp)(\eta^5-C_5H_4B(C_6F_5)_3)VH(CO)]^{.11}$ The reductive elimination of the hydride leads to simultaneous formation of 0.5 equiv of the monocarbonylvanadium-(III) $[(Cp)(C_5H_4B(C_6F_5)_3)V(CO)] \{A\}$ and 0.5 equiv of the monocarbonyl cationic [Cp₂V(CO)]⁺ species associated with the anion $[HB(C_6F_5)_3]^-$ {**B**}. Disproportionation of **{A}** gives the borane-substituted cyclopentadienyl complexes 3 and 4, whereas {B} disproportionates to 5 and a suggested $[Cp_2VH][HB(C_6F_5)_3]$ species that we have not been able to observe due to its probable decomposition (Scheme 1). Gas phase analysis by MS of the gas above the reaction mixture, in a closed vessel, did not show evidence of H₂ evolution during the synthesis of the crude product in pentane.

Conclusion

In summary, combination of 1 with [Cp₂VCO] gives new vanadium complexes. In the same reaction, three vanadium complexes were found showing the complexity of the vanadium chemistry, resulting from the oxidation of the metal center toward the divalent vanadocene **2**. Complexes **3** and **4** represent the first structurally characterized ring-borylated vanadium zwitterionic complexes.

Experimental Section

All manipulations were carried out using standard Schlenk line or drybox techniques under an atmosphere of argon. Solvents were refluxed and dried over appropriate drying agents under an atmosphere of argon, collected by distillation,



and stored in a drybox over activated 4 Å molecular sieves. Deuterated solvents were degassed and dried over activated 4 Å molecular sieves. NMR data were recorded using Bruker AMX-400 and AC-200 spectrometers, referenced internally to residual protio solvent (¹H) resonances, and are reported relative to tetramethylsilane ($\delta = 0$ ppm). ¹⁹F NMR (188.298 MHz) spectra were recorded on a Bruker AC-200 spectrometer (reference CF₃CO₂H). ⁵¹V NMR (105.24 MHz; reference VOCl₃ in C₆D₆, 9:1) and ¹¹B NMR (128.37 MHz; reference BF₃·Et₂O) spectra were recorded on a Bruker AMX-400 spectrometer. Elemental analyses (C, H) were performed at the Laboratoire de Chimie de Coordination (Toulouse, France). [Cp₂VCO] and B(C₆F₅)₃ were prepared according to the literature.^{12,13}

Preparation of $[(\eta^5-Cp)(\eta^5-C_5H_4B(C_6F_5)_3)V]$ (3), $[(\eta^5-Cp)-(\eta^5-C_5H_4B(C_6F_5)_3)V(CO)_2]$ (4), and $[Cp_2V(CO)_2][HB(C_6F_5)_3]$ (5). To a stirred pentane solution (5 mL) of $[Cp_2VCO]$ (2) (36 mg, 0.2 mmol) was added in small portions solid $B(C_6F_5)_3$ (102 mg, 0.2 mmol). The resulting yellow-brown slurry was left one week at room temperature until blue crystals of 3 appeared. After careful hand-separation of 3, the yellow-brown solid was dissolved in THF (1–2 mL) and layered with pentane to give

⁽¹¹⁾ V–H bonds were reported in the literature: (a) Jonas, K.; Wiskamp, V. J. Am. Chem. Soc. **1983**, *105*, 5480–5481. (b) Bansemer, J. R. L.; Huffman, C.; Caulton, K. G. J. Am. Chem. Soc. **1983**, *105*, 6163–6164. (c) Berno, P.; Gambarotta, S. Angew. Chem. **1995**, *107*, 871–873; Angew. Chem., Int. Ed. Engl. **1995**, *34*, 822–824. (d) Clancy, G. P.; Clark, H. C. S.; Clentsmith, G. K. B.; Cloke, F. G. N.; Hitchcock, P. B. J. Chem. Soc., Dalton Trans. **1999**, 3345–3347. Very recently, the zwitterionic hydride ring-borylated ansa-chromocene [Me4C2($\eta^5-C_5H_4$]($\eta^5-C_5H_3$ B(C₆F₅)₃)CrH(CO)] has been isolated from [Me4C2($\eta^5-C_5H_4$)($q^5-C_5H_4$)($q^5-C_5H_4$), Sinnema, P.-J.; Shapiro, P. J.; Foo, D. M. J.; Twamley, B. J. Am. Chem. Soc. **2002**, *124*, 10996–10997.

 ⁽¹²⁾ Massey, A. G.; Park, A. J. J. Organomet. Chem. 1966, 5, 218.
 (13) Calderazzo, F.; Fachinetti, G.; Floriani, C. J. Am. Chem. Soc.
 1974, 96, 3695–3696.

4 as a crystalline, orange solid suitable for X-ray structure determination. In another experiment (using the same experimental conditions), the crude yellow-brown product was dissolved in toluene (1 mL) and layered with pentane to give 5 as a crystalline, orange-yellow solid suitable for X-ray structure determination. Note that 4 is very often contaminated by some proportion of 5 and similarly 5 by some amount of 4, as indicated by ¹H NMR spectroscopy. C, H analytical results are in the same range for $\boldsymbol{4}$ ($C_{30}H_9BF_{15}O_2V)$ as for $\boldsymbol{5}$ ($C_{30}H_{11}$ $BF_{15}O_2V$) and fall within the experimental error for both compounds, and they are not indicative of the purity of the product. Analysis of 4 and 5 was attempted for characterization and gave reproducible analysis. In another procedure, the crude yellow product can be obtained directly by admission of CO gas in a stoichiometric mixture of [VCp₂] and 1 in pentane.

3: $\mu_{\text{eff}} = 2.85 \ \mu_{\text{B}}$. Anal. Calcd for C₂₈H₉BF₁₅V: C 48.59, H 1.31. Found: C 48.40, H 1.22. Yield: 22 mg (50%). 4: 1H NMR (200 MHz, CD₃CN) 5.19 (5H, Cp), 5.36 (b, 2H, C₅H₄) 5.49 (t, 2H, C₅H₄); ¹⁹F NMR (CD₃CN) -52.9; -86.2; -91.0 (*o*, *p*, *m*-F, C₆F₅); ¹¹B (CD₃CN) -14.9; ⁵¹V (CD₃CN) -1652; IR (Nujol mull) ν_{CO} 2050, 2004. Anal. Calcd for C₃₀H₉BF₁₅O₂V: C 48.16, H 1.21. Found: C 48.4, H 1.35. Yield: 21 mg (59%). 5: ¹H NMR (200 MHz, CD₃CN) 5. 59 (10H, Cp), 3.63 (q (${}^{1}J_{(BH)} = 91$ Hz), B-*H*); ¹⁹F (CD₃CN) -58.9, -88.8, -92.0 (*o*, *p*, *m*-F, C₆F₅); ¹¹B (CD₃-CN) -25.5 (d, $({}^{1}J_{(BH)} = 87$ Hz), B-H); ${}^{51}V$: -1661; IR (Nujol mull) $\nu_{(CO)}$ 2038, 1990; $\nu_{(BH)}$ 3123. Anal. Calcd for C₃₀H₁₁-BF15O2V: C 48.03, H 1.48. Found: 48. 38, C 1.30. Yield: 30 mg (83%).

Crystallographic Data for 3, 4, and 5. Selected crystals, sensitive to air and moisture, were suspended in oil on a glass slide. Under a microscope, a single block was isolated. For structures 3, 4, and 5 (Table 1), data were collected using a Stoe Imaging Plate Diffraction System (IPDS). The final unit cell parameters were obtained by least-squares refinement of a set of 5000 reflections, and crystal decay was monitored by measuring 200 reflections by image. No fluctuations of the intensity were observed over the course of the data collection.

A semiempirical correction absorption¹⁴ was applied to the data. The structure was solved by direct methods using SIR92¹⁵ and refined by least-squares procedures on F^2 with the aid of SHELXL97,¹⁶ included in WinGX (version 1.63).¹⁷ The atomic scattering factors were taken from International Tables for X-Ray Crystallography.¹⁸ All hydrogen atoms were located on a difference Fourier map. All the remaining nonhydrogen atoms were anisotropically refined, and in the last refinement cycles a weighting scheme was used where weights were calculated from the following formula: $w = 1/[\sigma^2(F_0^2) +$ $(aP)^2 + bP$, where $P = (F_0^2 + 2F_c^2)/3$. Drawing of the molecules was performed using the program ORTEP32.¹⁹ Criteria for a satisfactory, complete analysis were the ratio of rms shift to standard deviation being less than 0.1 and no significant features in final difference maps.

Acknowledgment. We thank the CNRS for financial support.

Supporting Information Available: Crystallographic data for compounds **3**, **4**, and **5** including ORTEP diagrams and tables of crystal data and data collection parameters, atomic coordinates, anisotropic displacement parameters, and all bond lengths and bond angles. This material is available free of charge via the Internet at http://pubs.acs.org.

OM034383W

435.

Hamilton, W. C., Eds.; Kynoch Press: Birmingham, England, 1974; Vol. IV

⁽¹⁴⁾ Walker, N.; Stuart, D. Acta Crystallogr., Sect. A 1983, 39, 158. (15) Altomare, A.; Cascarano, G.; Giacovazzo, G.; Guagliardi, A.; Burla, M. C.; Polidori, G.; Camalli, M. J. Appl. Crystallogr. **1994**, *27*,

⁽¹⁶⁾ Sheldrick, G. M. SHELX97, Program for Crystal Structure (16) Sheidrick, G. M. STELLAY, Frogram for Crystal Sciences Analysis; University of Göttingen: Germany, 1998.
(17) Farrugia, L. J. J. Appl. Crystallogr. 1999, 32, 837.
(18) International Tables for X-Ray Crystallography, Ibers, J. A.,

⁽¹⁹⁾ Farrugia, L. J. J. Appl. Crystallogr. 1997, 30, 565.