



Enantioselective Preparation of Allenecarboxylates by Asymmetric Horner-Wadsworth-Emmons Reaction


Kiyoshi Tanaka, Kenji Otsubo and Kaoru Fuji*

Institute for Chemical Research, Kyoto University, Uji 611, Japan

Abstract: Optically active 4,4-disubstituted conjugated allenecarboxylates were enantioselectively prepared via C=C bond formation through an asymmetric Horner-Wadsworth-Emmons reaction with an optically active phosphonoacetate reagent. Copyright © 1996 Elsevier Science Ltd

Conjugated allenecarboxylates (alka-2,3-dienoates) are interesting building blocks showing a versatile reactivity, and hence have been widely used in a variety of synthetic transformations.¹ More interesting is that the selective transfer of the axial chirality of the optically active allenic molecules to the central chirality by appropriate chemical or biological transformation might result in an efficient method for construction of optically active compounds.² Among various methods for the synthesis of allenecarboxylic acid derivatives,³ a relatively limited number of approaches to optically active 4,4-disubstituted alka-2,3-dienoates are known. The latter involves optical resolution of the allenic acids,⁴ displacement from optically active propargylic compounds,^{3a,5} enantioselective elimination of chiral selenoxides⁶ or asymmetric Horner-Wadsworth-Emmons (HWE) reaction of unsymmetrical ketenes or their equivalents.⁷ However, these transformations often suffered from tedious and multistage operational steps^{5,6} as well as low chemical and optical yields.⁷ We recently reported a convenient one-pot procedure for the preparation of allenecarboxylates based on an HWE reaction of *in situ* generated ketenes⁸ from enolates of 2,6-di-*tert*-butyl-4-methylphenyl (BHT) esters.⁹ We wish to report herein an enantioselective version of the above reaction.


Beside (*S*)-1,¹⁰ analogous reagents, (*S*)-2, -3 and -4, possessing substituents at 3,3' positions in the binaphthyl auxiliary moiety,^{11,12} were newly prepared and the reaction of ethyl phenyl ketene derived from the BHT ester of 2-phenylbutyric acid **5a** with these reagents was examined (Table 1). Our previous experiments⁹ suggested that an additive ZnCl₂ or SnCl₂ is essential for high chemical yield in production of allenic esters. Therefore, a solution of ZnCl₂ (2 eq) in ether¹⁴ was added after 1 h to a solution of the ketene generated from 2-phenylbutyrate (2 eq) and *n*-BuLi (2 eq) at -78 °C, and then the solution of the anion of the HWE reagent (1 eq) was added. Subsequently, the whole reaction mixture was allowed to room temperature with stirring for the indicated hours in Table 1. (*Ra*)-Allenecarboxylate **6** was obtained in all cases. The chemical yield of methyl allenecarboxylate **6** was generally high in the presence of an additive ZnCl₂ as expected; however, optical yields with (*S*)-1 were disappointing



(*S*)-1 : X = H
 (*S*)-2 : X = Me
 (*S*)-3 : X = SiMe₃
 (*S*)-4 : X = Ph

(entries 1 - 3). Using LDA as a base, a better result was obtained with (*S*)-2 (entry 5) and it turned out that the prolonged reaction time caused no significant racemization (entry 6). Use of Cp_2ZrCl_2 , SnCl_2 , $\text{CITi}(\text{O}^i\text{Pr})_3$ or $n\text{-Bu}_3\text{SnF}$ in place of ZnCl_2 as an additive under the reaction conditions of entry 5 gave no improvement. Among the bases employed, KHMDS provided more satisfactory results than LDA (entries 4-6, 9 and 11) and the optimum ee up to 67% was observed with the methyl substituted reagent (*S*)-2 (entry 7). Compared with the reported value of 22.8% ee observed in a similar approach by asymmetric HWE reactions,⁷ this value is remarkable. The usage of the increased amount of ZnCl_2 did not affect the optical yield but the chemical yield (entry 8). Neither DME nor Et_2O -toluene as solvent improved the results.

Table 1. Enantioselective Preparation of Allenecarboxylates **6**.^a



entry	phosphonoacetate	ZnCl_2 (eq)	base	time (h)	yield (%) ^b	% ee ^c
1	(<i>S</i>)-1	none	LDA	15	22	0
2	(<i>S</i>)-1	2	LDA	15	88	17
3	(<i>S</i>)-1	3	KHMDS	3	80	22
4	(<i>S</i>)-2	2	LDA	15	78	5
5	(<i>S</i>)-2	3	LDA	3	65	47
6	(<i>S</i>)-2	3	LDA	15	51	45
7	(<i>S</i>)-2	3	KHMDS	4	87	67
8	(<i>S</i>)-2	6	KHMDS	4	14	64
9	(<i>S</i>)-3	3	LDA	15	77	10
10	(<i>S</i>)-3	3	KHMDS	4	83	50
11	(<i>S</i>)-4	3	LDA	5	71	17
12	(<i>S</i>)-4	3	KHMDS	4	78	62

^a **5a** (2.0 eq) and each 1.0 eq of base and the indicated amount of ZnCl_2 were employed toward 1.0 eq.

of (*S*)-1 - 4. ^b Isolated yield based on (*S*)-1 - 4. ^c Determined by HPLC analysis (Chiralpak AS).

Using the optically active reagent (*S*)-2 and the established conditions discussed above, i.e., in THF in the presence of ZnCl_2 (3 eq) with KHMDS as a base for 4 h, the asymmetric HWE reactions were examined with a variety of *in situ* generated ketenes from the BHT esters of 2,2-disubstituted acetic acid derivatives. The results are summarized in Table 2.¹⁵ Generally, the reaction proceeded smoothly to give the corresponding allenecarboxylates in satisfactory chemical yields of 71-94% as well as enantioselectivities of 61-84% ee except for a few cases (entries 7 and 8). The absolute configuration of **6**, **7** and **10** was referred to the reported specific rotation⁷ of these compounds and the CD analysis¹⁶ allowed us to propose absolute configuration for the other optically active allenic carboxylates. These assignments are in full agreement with configurations predicted by the Lowe rule.¹⁷ Of particular interest is that the considerably high optical yields were obtained even when the difference in bulkiness between two substituents on the ketene moiety is not so significant (entries 2, 3 and 4).

The observed enantioselectivity with (*S*)-2 is best understood by considering the favorable transition state rather than the stability of the phosphoxetane intermediate. Thus, the addition of ZnCl_2 causes the rigidly


Table 2. Preparation of Optically Active Allenecarboxylates 7 - 14 with the Anion (*S*)-2.

entry	BHT esters			allenecarboxylate			
	5	R ¹	R ²	yield(%) ^a	% ee ^b (config.) ^c	[α] _D ²⁰ (solv, conc)	
1	5b	Ph	Me	7	94	62 (<i>Ra</i>) ^d	-168.1 (CCl ₄ , 0.5)
2	5c	<i>i</i> -Pr	Ph	8	71	81 (<i>Sa</i>)	+150.2 (CHCl ₃ , 0.25)
3	5d	cyclo-Hex	Ph	9	81	79 (<i>Sa</i>)	+106.0 (CHCl ₃ , 0.5)
4	5e		Ph	10 ^e	74	84 (<i>Ra</i>) ^d	-9.5 (CCl ₄ , 0.25)
5	5f	2-Naphthyl	Et	11	91	72 (<i>Ra</i>)	-213.5 (CHCl ₃ , 0.25)
6	5g	Ph ₂ CH	Me	12	85	61 (<i>Ra</i>)	-4.5 (CHCl ₃ , 0.18)
7	5h	1-Naphthyl	Me	13	83	4 (<i>Ra</i>)	-5.6 (CHCl ₃ , 0.5)
8	5i	2,4,6-trimethyl-phenyl	Me	14	60	10 (<i>Ra</i>)	-23.1 (CHCl ₃ , 0.2)

^aIsolated yield based on (*S*)-2. ^bDetermined by HPLC analysis on chiral column. ^cDeduced from CD spectra (entries 1-6)¹⁶ and assumed on the basis of the Lowe rule¹⁷ (entries 7 and 8). ^dRef 7. ^eThe product 10 had a tendency to decompose during isolation and purification procedures.

chelated phosphate anion bound by Zn²⁺ as depicted in Figure 1, where the axially dissymmetric binaphthyl group dictates the orientation of the approach to the electrophile from the less hindered *si* face of the reagent (route a). Assuming R¹ > R² in bulkiness, it is likely that the nucleophile approaches the LUMO of the ketene carbonyl from the face, including the less bulky R², to avoid the severer steric repulsive interaction with R¹. Consequently, the methyl group at 3-position on the naphthalene molecule exerts its efficiency as the bulkiness of R² group increases. No reasonable explanation for the low optical yields obtained with (*S*)-3, -4 as well as in reactions starting from the BHT esters of an aryl substituent bearing the substituent at the ortho position (entries 7 and 8 in Table 2), is available at the moment.

In conclusion, the present study demonstrates a simple one-flask preparation of 4,4-disubstituted allenecarboxylates with a high degree of optical purity. The increasing synthetic availability of optically active allenes must enhance their attractiveness as versatile chiral building blocks. Since both the starting BHT esters and the allenic products are rather stable compounds, this transformation might provide an alternative and practical methodology for construction of useful optically active compounds such as natural products.

Figure 1. Possible Mechanistic Explanation for the Observed Enantioselectivity in Asymmetric HWE Reaction with (*S*)-2.

References and Notes

- 1) a) Schuster, H. F.; Coppola, G. M. "Allenenes in Organic Synthesis" Wiley: New York, 1984; b) Jung, M. E. In *Comprehensive Organic Synthesis*; Trost, B. M., Fleming, I., Eds.; Pergamon: Oxford, 1991; Vol. 4, pp 53 - 58; c) Naruse, Y.; Kakita, S.; Tsunekawa, A. *Synlett* **1995**, 711-713; d) Trost, B. M.; Kottirsch, G. *J. Am. Chem. Soc.* **1990**, *112*, 2816-2818; e) Ishar, M. P. S.; Gandhi, R. P. *Tetrahedron* **1993**, *49*, 6729-6740; f) Tsuboi, S.; Kuroda, H.; Takatsuka, S.; Fukawa, T.; Sakai, T.; Utaka, M. *J. Org. Chem.* **1993**, *58*, 5952-5957; g) Tanaka, H.; Sumida, S.; Sorajo, K.; Torii, S. *J. Chem. Soc. Chem. Commun.* **1994**, 1461-1462; h) Jung, M. E.; Zimmerman, C. N.; Lowen, G. T.; Khan, S. I. *Tetrahedron Lett.* **1993**, *34*, 4453-4456; i) Gillmann, T.; Weeber, T. *Synlett* **1994**, 649-650; j) Oppolzer, W.; Chapuis, C.; Dupuis, D.; Guo, M. *Helv. Chim. Acta* **1985**, *68*, 2100-2114; k) Kanda, Y.; Fukuyama, T. *J. Am. Chem. Soc.* **1993**, *115*, 8451-8452; l) Zhang, C.; Lu, X. *Synlett* **1995**, 645-646.
- 2) a) Musierowicz, S.; Wróblewski, A. E. *Tetrahedron* **1978**, *34*, 461-465; b) Ikeda, I.; Kanematsu, K. *J. Chem. Soc. Chem. Commun.* **1995**, 453-454; c) Agosta, W. C. *J. Am. Chem. Soc.* **1964**, *86*, 2638-2642; d) Shingu, K.; Hagishita, S.; Nakagawa, M. *Tetrahedron Lett.* **1967**, 4371-4374; e) Crombie, L.; Jenkins, P. A. *J. Chem. Soc. Chem. Commun.* **1967**, 870-872; f) Rambeck, B.; Simon, H. *Angew. Chem. Int. Ed. Engl.* **1974**, *13*, 609; g) Schwab, J. M.; Lin, D. C. T. *J. Am. Chem. Soc.* **1985**, *107*, 6046-6052.
- 3) a) Tsuji, J.; Mandai, T. *Angew. Chem. Int. Ed. Engl.* **1995**, *34*, 2589-2612; b) Rossi, R.; Diversi, P. *Synthesis* **1973**, 25-36; c) Wadsworth, Jr., W. S.; Emmons, W. D. *J. Am. Chem. Soc.* **1961**, *83*, 1733-1738; d) Lang, R. W.; Hansen, H. J. *Org. Synth. Col. Vol. 7* **1990**, 232-235; e) Franck-Neumann, M.; Neff, D.; Nouali, H.; Martina, D.; de Cian, A. *Synlett* **1994**, 657-659; f) Sleeman, M. J.; Meehan, G. V. *Tetrahedron Lett.* **1989**, *30*, 3345-3348.
- 4) a) Runge, W.; Kresze, G. *Liebigs Ann. Chem.* **1975**, 1361-1378; b) Kresze, G.; Runge, W.; Ruch, E. *ibid.* **1972**, *756*, 112-127.
- 5) a) Marshall, J. A.; Pinney, K. G. *J. Org. Chem.* **1993**, *58*, 7180; b) Marshall, J. A.; Wallace, E. M. *J. Org. Chem.* **1995**, *60*, 796-797.
- 6) Nishibayashi, Y.; Singh, J. D.; Fukuzawa, S.; Uemura, S. *J. Org. Chem.* **1995**, *60*, 4114-4120.
- 7) a) Musierowicz, S.; Wróblewski, A. E. *Tetrahedron* **1980**, *36*, 1375-1380; b) Tömöskösi, I.; Bestmann, H. J. *Tetrahedron Lett.* **1964**, 1293-1295.
- 8) a) Häner, R.; Laube, T.; Seebach, D. *J. Am. Chem. Soc.* **1985**, *107*, 5396-5403; b) Baigrie, L. M.; Seiklay, H. R.; Tidwell, T. T. *ibid.* **1985**, *107*, 5391-5396.
- 9) a) Tanaka, K.; Otsubo, K.; Fuji, K. *Synlett* **1995**, 933-934; b) *Idem Tetrahedron Lett.* **1995**, *36*, 9513-9514.
- 10) Tanaka, K.; Ohta, Y.; Fuji, K.; Taga, T. *Tetrahedron Lett.* **1993**, *34*, 4071-4074.
- 11) Cox, P. J.; Wang, W.; Snieckus, V. *Tetrahedron Lett.* **1992**, *33*, 2253-2256.
- 12) (S)-2 (100 %ee, $[\alpha]^{20}_D +414$ (CHCl₃, c 1.0)) was prepared from (S)-3,3'-dimethyl-1,1'-bi-2-naphthol (1.0 eq) with Cl₂POCH₂CO₂Me (1.2 eq) in the presence of Et₃N (2.3 eq) in CH₂Cl₂ at 0 °C to r. temp. for 3 h in 72 % yield.¹³ (S)-3 (98 %ee, 73 % yield, $[\alpha]^{18}_D +418$ (CHCl₃, c 1.0)) and (S)-4 (100 %ee, 84 % yield, $[\alpha]^{18}_D +381$ (CHCl₃, c 1.0)) were prepared from the corresponding binaphthol (1.0 eq) with Cl₂POCH₂CO₂Me (1.1 eq) in the presence of *n*-BuLi (2.1 eq) in THF at -78 °C to r. temp. for 15 h.
- 13) Still, W. C.; Gennari, C. *Tetrahedron Lett.* **1983**, *24*, 4405-4408.
- 14) House, H. O.; Crumrine, D. S.; Teranishi, A. Y.; Olmstead, H. D. *J. Am. Chem. Soc.* **1973**, *95*, 3310-3324.
- 15) All new compounds in this paper provided satisfactory spectroscopic and analytical data.
- 16) CD (in MeOH) spectral data of allenecarboxylates **7** - **14** of the optical purity indicated in Table 2: **7**; λ_{ext} 238nm ($\Delta\epsilon$ -5.83), **8**; λ_{ext} 232nm ($\Delta\epsilon$ +7.07), **9**; λ_{ext} 232nm ($\Delta\epsilon$ +7.65), **10**; λ_{ext} 249nm ($\Delta\epsilon$ -6.48), **11**; λ_{ext} 253nm ($\Delta\epsilon$ -10.5), **12**; λ_{ext} 228nm ($\Delta\epsilon$ -2.60).
- 17) a) Lowe, G. J. *J. Chem. Soc. Chem. Commun.* **1965**, 411-413; b) Brewster, J. H. *Topics in Stereochemistry*, **1967**, *2*, pp 1 - 72; c) Crabbé, P.; Velarde, E.; Anderson, H. W.; Clark, S. D.; Moore, W. R.; Drake, A. F.; Mason, S. F. *J. Chem. Soc. Chem. Commun.* **1971**, 1261-1264.