


An Efficient Synthesis of Selenocarbonyl Compounds by the Treatment of Carbonyl Compounds with Bis[1,5-cyclooctanediylboryl] Selenide

Kazuaki SHIMADA, Norikazu JIN, Manabu FUJIMURA, Yumi NAGANO,
Eiichi KUDOH, and Yuji TAKIKAWA*

Department of Applied Chemistry and Molecular Science,
Faculty of Engineering, Iwate University, Morioka, Iwate 020

An efficient synthesis of selenoaldehydes, selenoketones, and selenoamides was achieved by the treatment of carbonyl compounds with bis[1,5-cyclooctanediylboryl] selenide.

One-step conversion of carbonyl compounds to the corresponding selenocarbonyl compounds by using selenating reagents is one of the most important concerns in organic heteroatom chemistry.¹⁻⁵⁾ However, among the reagents possessing reactive metal-Se bonds, boryl selenides and boryl metal selenides have been investigated mainly from structural interests⁶⁾ in spite of their selenating potential resulting from the significant affinity of B-Se bonds to oxygen functionalities. Actually, some boryl selenide was thought to play an essential role in the selenation of carbonyl compounds by the treatment of (Me₃Si)₂Se-BF₃·OEt₂⁵⁾ or [(c-C₆H₁₁)₃Sn]₂Se-BCl₃.⁶⁾ It was naturally supposed that structural modification of boryl selenides by so-called steric protection was much effective in reducing their air-lability, as is widely known to be the case with general dialkylborane compounds. Herein, we wish to report a novel and efficient conversion of carbonyl compounds to the corresponding selenocarbonyl compounds by using bis[1,5-cyclooctanediylboryl] selenide 1.



Bis[1,5-cyclooctanediylboryl] selenide 1 was prepared by treating elemental selenium with 9-BBN in mesitylene at refluxing temperature under N₂ atmosphere based on the Köster's method.⁷⁾ Subsequently, reagent 1 was treated with aldehydes and ketones under N₂ atmosphere at room or a higher temperature in the presence of 2,3-dimethyl-1,3-butadiene. After usual work-up and chromatographic separation, [4+2] cycloadducts of selones with the

diene were isolated in modest to high yields besides the recovery of small amounts of aldehydes and ketones. N,N-Dialkyl carboamides and an ester were similarly treated with reagent 1 in the absence of a diene, and the corresponding selenoamides and a selenoester were isolated in modest yields. All results are shown in Table 1 and Table 2.

In particular, the excellent selenating activity of reagent 1 was exhibited by the reaction with benzaldehyde 2 and sterically crowded ketones, 3 and 4, in the absence of 2,3-dimethyl-1,3-butadiene. By treating with reagent 1, 2 was converted to dibenzyl diselenide 5 (44%) derived by the further reduction of transiently-generated selenobenzaldehyde 6.⁸⁾ Fenchone 3 and santonine 4 also afforded selones, 7⁹⁻¹¹⁾ and 8¹²⁾, in 35% and 23% yields, respectively, besides inseparable stereoisomeric mixtures of diselenides in 20-40% yields.^{8,13,14)} Subsequent NaBH₄ or LiAlH₄ reduction of the resulting selones in MeOH or Et₂O at 0 °C to room temperature followed by an aerobic work-up afforded sole diselenides, 9 (quant.) and 10 (67%) respectively.¹²⁾ Orientation of the C-Se bonds of each product was supposed to be endo and exo, respectively, as were similarly mentioned in the hydride reduction of 3¹⁵⁾ and 4.¹⁶⁾ These results demonstrated that the strong electronic interaction of boron atom with carbonyl oxygen atom behaved as a substantial controlling factor of the reactions and steric factors of substrates and the reagent played a less important role in the reactions. In contrast, selenation of cinnamic aldehyde gave an unexpected compound 11 (37%)¹²⁾ which may be produced by selenium-extrusion of a dimeric cycloadduct of α , β -unsaturated selenoaldehyde.^{17,18)}

Conclusively, efficient synthesis of selenocarbonyl compounds was achieved by treating carbonyl compounds with bis[1,5-cyclooctanediylboryl] selenide 1. Further synthetic expansions of reagent 1 to tellurium-transfer reagents as well as the application of diselenides 9 and 10 to novel chiral auxiliaries for asymmetric syntheses are now in progress in our laboratory.







Table 1. Trapping of selenoaldehydes and selenoketones with 2,3-dimethyl-1,3-butadiene^{a)}

Substrate		<u>1</u>	Temp	Time	Yield
R ¹	R ²	equiv.	°C	h	%
C ₆ H ₅	H	1.2	110	18	79
4-CH ₃ -C ₆ H ₄	H	1.2	110	8	82
CH ₃	H	1.2	150	18	32
C ₆ H ₅	CH ₃	2.0	80	15	54
C ₆ H ₅	C ₂ H ₅	2.0	80	24	81
C ₆ H ₅	C ₆ H ₅	2.0	110	24	96
C ₂ H ₅	CH ₃	2.0	110	48	38
(CH ₃) ₂ CH	CH ₃	2.0	110	48	53
-(CH ₂) ₅ -		2.0	130	48	50
9-fluorenone		2.0	110	24	82

a) All reactions were carried out in a sealed tube.

Table 2. Synthesis of selenocarbonyl compounds by treating carboxylic acid derivatives with reagent 1^{a)}

Substrate		Solvent	Temp	Time	Yield
R ¹	R ²		°C	h	%
H	N(CH ₃) ₂	toluene	r.t.	24	39
H	N[CH(CH ₃) ₂] ₂	benzene/CH ₂ Cl ₂	110	1	16
CH ₃	N(CH ₃) ₂	mesitylene/CHCl ₃	63	5	34
C ₆ H ₅	OC ₂ H ₅	benzene/CH ₂ Cl ₂	130	10	14 ^{b)}

a) All reactions were carried out in a sealed tube. b) Starting ester was recovered in 81% in this case.

We are grateful to Saneyoshi Scholarship Foundation for financial support in this work.

References

- 1) M. Segi, T. Nakajima, S. Suga, S. Murai, I. Ryu, A. Ogawa, and N. Sonoda, *J. Am. Chem. Soc.*, **110**, 1976 (1988).
- 2) M. Segi, M. Takahashi, T. Nakajima, S. Suga, S. Murai, and N. Sonoda, *Tetrahedron Lett.*, **29**, 6965 (1988).
- 3) M. Segi, T. Koyama, T. Nakajima, S. Suga, S. Murai, and N. Sonoda, *ibid.*, **30**, 2095 (1989).
- 4) M. Segi, A. Kojima, T. Nakajima, and S. Suga, *Synlett*, **1991**, 105.
- 5) Y. Takikawa, A. Uwano, H. Watanabe, M. Asanuma, and K. Shimada, *Tetrahedron Lett.*, **30**, 6047 (1989).
- 6) K. Steliou and M. Mrani, *J. Am. Chem. Soc.*, **104**, 3104 (1984).
- 7) R. Köster, G. Seidel, and M. Yalpani, *Chem. Ber.*, **122**, 1815 (1989), and references cited therein.
- 8) J. W. Lewicki, W. H. H. Günther, and J. Y. C. Chu, *J. Org. Chem.*, **43**, 2672 (1978).
- 9) T. G. Back, D. H. R. Barton, M. R. Britten-Kelly, and F. S. Guziec, Jr., *J. Chem. Soc., Chem. Commun.*, **1975**, 539.
- 10) F. S. Guziec, Jr. and C. A. Moustakis, *J. Org. Chem.*, **49**, 189 (1984).
- 11) A. Ishii, R. Okazaki, and N. Inamoto, *Bull. Chem. Soc. Jpn.*, **61**, 861 (1988).
- 12) **8**: A deep blue oil; bp 130 °C(7 mmHg); MS(m/e) 270(M⁺,15%,⁸⁰Se); IR(neat) 2990, 1450, 1080, 800 cm⁻¹; ¹H NMR(CDCl₃) δ 0.83(3H,s), 1.04(3H,s), 1.20(3H,s), 1.30-2.30(13H,m); **9**: Yellow needles; mp 50.0-51.0 °C; MS(m/e) 434(M⁺,2%,⁸⁰Se), 137(bp); IR(KBr) 2860, 1440, 1360, 720 cm⁻¹; ¹H NMR(CDCl₃) δ 1.02(6H,s), 1.08(6H,s), 1.20(6H,s), 0.95-1.88 (14H,m), 3.10(2H,d,J=3.0 Hz); **10**: Yellow oil; MS(m/e) 542(M⁺,21%,⁸⁰Se); IR(neat) 2950, 1390, 1380, 880, 780 cm⁻¹; ¹H NMR(CDCl₃) δ 0.83(6H,s), 0.94(6H,s), 1.04(6H,s), 1.17-2.32(24H,m), 1.18-1.29(2H,m), 3.20-3.53(2H,m); **11**: Colorless needles; mp 105.5-106 °C; MS(m/e) 312(M⁺,bp,⁸⁰Se), 310(M⁺,42%,⁷⁸Se), 231(88%,M⁺-PhCH=CH); IR(KBr) 1570, 1470, 1440, 1270, 1240, 1140, 1100, 1070, 1010, 960, 730, 670 cm⁻¹; ¹H NMR (CDCl₃) δ 4.13(1H,ddd,J=7.5,3.0,2.5 Hz), 4.67(1H,dd,J=8.5,7.5 Hz), 6.07(1H,dd,J=6.5, 3.0 Hz), 6.28(1H,d,J=15.5 Hz), 6.57(1H,dd,J=15.5,8.5 Hz), 6.83(1H,dd,J=6.5,2.5 Hz), 7.01-7.49(10H,m). Found: C,68.73; H,5.14%. Calcd for C₁₈H₁₆Se: C,69.45; H,5.18%.
- 13) Trapping of selones with 2,3-dimethyl-1,3-butadiene was not successful in these cases.
- 14) NaBH₄ reduction of diselenides in ethanol followed by the treatment of an excess amount of methyl iodide afforded inseparable stereoisomeric mixtures of methylseleno derivatives in modest yields.
- 15) H. C. Brown and H. R. Deck, *J. Am. Chem. Soc.*, **87**, 5620 (1965).
- 16) Y. Fujikura, T. Kitsuki, and M. Nakajima, *Jpn Kokai Tokkyo Koho*, JP 63 99,028 (1988).
- 17) Treatment of cinnamic aldehyde with **1** at 110 °C for 18 h in the presence of an excess amount of 2,3-dimethyl-1,3-butadiene in a sealed tube also afforded **11** in 51% yield.
- 18) The stereochemistry of **11** was not characterized by ¹H NMR data.

(Received June 22, 1992)