

254. Synthesis of Cycloprop[*b*]anthracenes via Trapping of *o*-Naphthoquinodimethane

by Paul Müller* and Domingo Rodriguez

Département de Chimie Organique, Université de Genève, CH-1211 Genève 4

(5.X.1983)

Summary

Cycloprop[*b*]anthracenes **1**, **1a**, **1b** are prepared in a 3-step synthesis starting from *o*-di(iodomethyl)benzene. The key step consists of trapping of *o*-naphthoquinodimethane (**9**) with 1,2-di- and tetrahalogenocyclopropenes (**3**, **3a**, **4b**).

In previous communications we have reported the synthesis of 1,1-dihalogenocycloprop[*b*]anthracenes **1** and **1a** [1]. The key step involved cycloaddition of 2,3-dimethylidene-1,2,3,4-tetrahydronaphthalene (**2**) to tetrahalogenocyclopropene **3**. The cycloadduct **4** was first dehydrogenated with 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ) to **5** and subsequently aromatized by treatment with *t*-BuOK. Several approaches were directed towards preparation of the parent compound, cycloprop[*b*]anthracene (**1b**), for example *via* reduction of **1** with LiAlH₄/AlCl₃ [2] or cyclopropanation of suitably 2,3-disubstituted-1,4-dihydroanthracenes **5**, but none were successful. A new scheme was then developed based on the trapping of *o*-quinodimethanes with cyclopropenes. The only precedent for this approach is the electrocyclic ring opening of *trans*-1,2-diphenylbenzocyclobutene followed by addition to tetrachlorocyclopropene which was used in the synthesis of 1,1-dichloro-2,7-diphenylcyclopropa[*b*]naphthalene [3]. However, the reaction is not general; in particular it fails with unsubstituted benzocyclobutene, mainly because of the decomposition of tetrachlorocyclopropene at the temperature required for benzocyclobutene ring opening.

Since we considered 1,2-dihalogenocyclopropenes potentially explosive [4] it appeared necessary to generate the *o*-quinodimethane at low temperature. An elegant method for this has been used by *Sondheimer et al.* [5] and worked out in all details by *Roth & Bartmann* [6]. *o*-Di(iodomethyl)benzene (**7**) is converted to *o*-di(2-propynyl)benzene (**8**) [6]. Upon treatment with *t*-BuOK in THF/*t*-BuOH **8** isomerized to the *bis*-allene at -78° . The latter is extracted with hexane. It cyclizes upon standing to *o*-naphthoquinodimethane (**9**) at *ca.* 30° . The latter can be trapped by addition of dienophiles to the cooled solution, which is allowed to warm up to 25° . *Roth & Bartmann* [6] obtained yields of 90–100% for trapping with *trans*-1,2-dichloroethene, O₂ and SO₂. The reaction of **8** with O₂ has also been reported by *Gisin & Wirz* [7].

While this work was in progress we were informed by *Billups et al.* [9] that a newly developed synthon, 1-chloro-2-bromocyclopropene (**3b**) undergoes smooth cycloaddition to **2**. Aromatization of the adduct, first with DDQ, then with *t*-BuOK afforded cycloprop[*b*]anthracene (**1b**). The successful synthesis of **1b** terminates speculations that it might not be capable of existence because of bond fixation [10].

We have repeated the preparation of **3b** according to *Billups et al.* [9] and confirmed its structure by reaction with diphenylisobenzofuran. Compound **3b** reacted with **9** to form the adduct **6b**, the structure of which was verified by comparison of the spectral data with those reported by *Billups*. Additional structural proof was obtained by dehydrohalogenation of **6b** to **1b** with *t*-BuOK under the usual conditions [1] [9].

Experiments involving the trapping of *o*-benzoquinodimethane, prepared by several procedures [5] [11] have so far given no positive results.

Financial support by the Swiss National Science Foundation (Project No. 2.236-0.81) is gratefully acknowledged. One of the authors (*D.R.*) thanks the sponsors of the 'Bourse Givaudan' for a scholarship. We are indebted to Prof. *W.R. Roth*, University of Bochum, for the detailed experimental procedure for preparation of **9** and to Prof. *E. Billups* for sending us a preprint of his article for synthesis of **1b**.

REFERENCES

- [1] *P. Müller & M. Rey*, *Helv. Chim. Acta* **65**, 1157 (1982); **64**, 354 (1981).
- [2] *P. Müller*, *Helv. Chim. Acta* **57**, 704 (1974).
- [3] *A.R. Browne & B. Halton*, *J. Chem. Soc., Chem. Commun.* **1972**, 1341.
- [4] *R. Breslow, G. Ryan & J.T. Groves*, *J. Am. Chem. Soc.* **92**, 988 (1970).
- [5] *D.A. Ben-Efraim & F. Sondheimer*, *Tetrahedron Lett.* **1963**, 313; *C.M. Bowes, D.F. Montecalvo & F. Sondheimer*, *Tetrahedron Lett.* **1973**, 3181.
- [6] *M. Bartmann*, Ph.D. Thesis, University of Bochum.
- [7] *M. Gisin & J. Wirz*, *Helv. Chim. Acta* **59**, 2273 (1976).
- [8] *R.P. Thummel, W.E. Cravey & Nutalene*, *J. Org. Chem.* **43**, 2473 (1978); *A.T. Blomquist & J.A. Verdol*, *J. Am. Chem. Soc.* **77**, 1806 (1955); *R.O. Angus & R.P. Johnson*, *J. Org. Chem.* **48**, 273 (1983).
- [9] *W.E. Billups, E.W. Casserly & B.E. Arney, Jr.*, *J. Am. Chem. Soc.* **1983**, submitted for publication.
- [10] *D. Davalian, P.J. Garratt, W. Koller & M.M. Mansuri*, *J. Org. Chem.* **45**, 4183 (1980).
- [11] *B.H. Han & P. Boudjouk*, *J. Org. Chem.* **47**, 751 (1982); *Y. Ito, Y. Amino, M. Nakatsuka & T. Saegusa*, *J. Am. Chem. Soc.* **105**, 1586 (1983).