Synthesis of 1-Oxidopyridinium-2-thiolato- and Pyridine-2-thiolatocomplexes of Nickel, Palladium, and Platinum

Jack L. Davidson, Peter N. Preston, and Maria V. Russo

Department of Chemistry, Heriot-Watt University, Riccarton, Currie, Edinburgh EH14 4AS

Quinoline *N*-oxide reacts with K[PtCl₃(CH₂=CH₂)] to give [PtCl₂(C₂H₄)(ONC₉H₇)] whereas 1-oxidopyridinium-2-thiol (1a) gives [Pt(SONC₅H₄)₂] (3a) and an insoluble oligomeric material containing at least four Pt atoms; complex (3a) and the oligomer can also be synthesised from K₂[PtCl₄]. The oligomeric material can be cleaved by PPh₃ and PMe₂Ph to give [PtCl(SONC₅H₄)(PPh₃)] (5) and [PtCl(SONC₅H₄)(PMe₂Ph)₂] (6) respectively, and compound (3a) is converted by PPh₃ into the monophosphine complex [Pt(SONC₅H₄)₂(PPh₃)] (7a). The complex (7a) and an analogue [Pd(SONC₅H₄)₂(PPh₃)] (7b) have been prepared from [MCl₂(PPh₃)₂] and the thallium salt of (1a), and related complexes [M(SC₅H₄N)₂(PPh₃)₂] [M = Pt (8a) or Pd (8b)] have been similarly synthesised from the thallium salt of 2-mercaptopyridine. Evidence for intramolecular scrambling in (7) and (8) has been adduced from variable temperature ¹H n.m.r. spectra.

A variety of metallic complexes of pyridine N-oxides have been described ¹ but particular attention has been focused on metal-containing compounds of the tautomeric ligand, 1oxidopyridinium-2-thiol [(1a) \implies (2)] because of their useful



biological activity (see for example, the use of Sn-,² Al-,³ and Zn⁴-containing compounds as antifungal ² and antibacterial ^{3,4} agents). We recently described ⁵ the powerful ligating properties of 1-oxidopyridinium-2-thiol in its reaction with $[WCl(CF_3C_2CF_3)_2(\eta-C_5H_5)]$ and now report further reactions of this, and the related pyridine-2-thiolato-ligand, in the Ni, Pd, and Pt series.

Results and Discussion

Pyridine N-oxide has been shown to react with Zeise's salt, K[PtCl₃(C₂H₄)], to give the complex [PtCl₂(C₂H₄)L] (L = ONC_5H_5)⁶ and we have now demonstrated that quinoline Noxide behaves in similar fashion to give a related complex $(L = ONC_9H_7)$ in 54% yield. In contrast to this behaviour, the tautomeric 1-oxidopyridinium-2-thiol $[(1a) \implies (2)]^{7}$ reacts with Zeise's salt in acetone at room temperature to give an orange soluble compound in poor yield and a yellow insoluble material. Analytical data, including ' direct ' measurement of oxygen content, for the soluble material provide an empirical formula of C10H8N2O2PtS2. From spectral analyses we suggest structure (3a) † for this compound although we cannot exclude the alternative cis form. The N-O stretching frequency occurs as a single band at 1 095 cm⁻¹ [cf. 1 083 and 1 115 cm⁻¹ in the free ligand ^{8,9} and single bands in the region 1 075-1 085 cm⁻¹ in analogues (3b)-(3d) ⁹ of (3a)]. The ¹H n.m.r.



parameters of (3a) are very close to those of the heteroaromatic component of the free ligand. ‡ In subsequent reactions of 1-oxidopyridinium-2-thiol with dipotassium tetrachloroplatinate it was shown that the complex (3a) was obtained using acetone as solvent, whereas the yellow insoluble material was again formed in water. Samples of the latter obtained from different sources (i.e. from Zeise's salt or K₂[PtCl₄]) gave differing results for elemental analysis (C, 19.10-21.05; H, 1.35-1.70; Cl, 9.10-13.50%). Since this material is very insoluble, has a high melting point (>350 $^{\circ}$ C), and has an i.r. spectrum which is almost identical to the complex (3a), we assume that it is oligomeric and contains bridging 1-oxidopyridinium-2-thiolato-ligands; from analytical data, it seems likely that the number of platinum atoms would be >4[cf. (4)]. The nature of bonding in the bridging ligand is unclear since we are unable to distinguish between a simple sulphur bridge and a situation where the SONC₅H₄ acts as a bidentate bridging ligand (cf. unusual bidentate bridge-bonding in the quinoline-8-oxo-complex $[{PtMe_2(ONC_{9}H_6)}_2]$.¹⁰

The oligomeric nature of (4) was demonstrated by allowing it to react with phosphine derivatives. It was converted by triphenylphosphine in low yield into a product believed to be (5), of undefined stereochemistry. The infrared spectrum of complex (5) was very similar to that of (3a) and the ¹H n.m.r. low- (δ 8.31) and high-field (δ 6.87) resonances were also comparable (*cf.* 8.24, 6.85). The ³¹P and ¹⁹⁵Pt n.m.r. spectra indicated the presence of one nuclear environment with a ³¹P-¹⁹⁵Pt coupling constant (*ca.* 3 870 Hz) in accord with a

⁺ For recent studies of the chromatographic separation of 1oxidopyridinium-2-thiol complexes of Group 8 metals, see K.-H. Konig, B. Steinbrach, G. Schneeweis, P. Chaudhuri, and H.-U. Ehancke, *Fresenius Z. Anal. Chem.*, 1979, **297**, 411.

[‡] For a detailed study of the ¹H n.m.r. spectra of pyridine N-oxide derivatives, see R. A. Abramovitch and J. B. David, J. Chem. Soc. B, 1966, 1137.









(A)

J. CHEM. SOC. DALTON TRANS. 1983

was apparent. The ³¹P and ¹⁹⁵Pt n.m.r. spectra are unexceptional indicating one phosphorus [δ 6.30, t, $J({}^{31}P{}^{-195}Pt)$ ca. 3 926 Hz] and one platinum $[\delta - 1 964, d, J(^{195}Pt^{-31}P) ca.$ 3 295 Hz] environments. We assign structure (7a) to this complex and present evidence for the fluxional process (A) (A') from high resolution ¹H n.m.r. spectral data. Four pairs of heterocyclic ligand resonances are observed in the 360 MHz ¹H n.m.r. spectrum of complex (7a) at 220 K. By analogy with literature data,* the following tentative assignments can be made: δ 8.25 and 7.98 (H² and H^{2'}), 8.06 and 7.57 (H⁵ and H^{5'}), 7.31 and 6.65 (H^{3 or 4} and H^{3' or 4'}), and 6.92 and 6.62 $(H^{3} \text{ or } 4 \text{ and } H^{3' \text{ or } 4'})$. At higher temperatures the doublet H^{2} and H2' resonances coalesce to a doublet and the H5 and H5' doublets coalesce to a broad singlet; a similar effect is appar-ent in the triplet $H^{3 \text{ or } 4}$ and $H^{3' \text{ or } 4'}$ resonances. These data provide evidence for a rapid intramolecular scrambling involving interconversion of unidentate and bidentate heterocyclic ligands *i.e.* (A) \Longrightarrow (A'). The alternative explanation for this behaviour is the existence of a five-co-ordinate structure for (7a) containing two inequivalent bidentate pyridine-2thiolato-ligands which undergoes a rapid intramolecular rearrangement without metal-oxygen bond cleavage. The latter interpretation seems less probable in view of the fact that related Pd¹¹ and Pt¹¹ dithiolato-complexes [M(S-S)₂L] $(S-S = S_2PR_2, S_2CNR_2, \text{ or } S_2COR; L = \text{tertiary phosphine},$ $\mathbf{R} = alkyl$ or aryl) consistently exhibit four-co-ordinate



square planar Pt¹¹ molecule.¹¹ In contrast to this behaviour, the oligomer (4) was transformed by dimethylphenylphosphine into a bis-phosphine complex [PtCl(SONC₅H₄)(PMe₂Ph)₂] (6). It seems probable that the different stoicheiometries of the products (5) and (6) isolated from the reactions of complex (4) with PPh₃ and PMe₂Ph reflect the large excess of phosphine employed in the latter case. The ³¹P n.m.r. spectrum of (6) indicates two phosphorus magnetic environments with $J(^{31}P^{-195}Pt)$ values (3 002 and 3 415 Hz) in accord with a cis square planar Pt¹¹ structure.¹¹ Accordingly two doublets are observed in the ¹⁹⁵Pt n.m.r. spectrum. Two structures are possible on this basis (6a) and the ionic form (6b). Unlike (5) which exhibits a v(Pt-Cl) mode at 320 cm⁻¹ no such band is present in the i.r. spectrum of (6) which argues in favour of (6b). Related 1,1-dithiolato-complexes [M(PR¹₃)₂(S₂PR²₂]⁺X⁻ (e.g. $R^1 = Ph$, $R^2 = OEt$) have been reported previously ¹² and in one particular case the complex [Pd(PEt₃)₂(S₂PPh₂)]-[S₂PPh₂] was structurally characterised by X-ray diffraction methods. We favour such a structure for complex (6) but the small quantities of complex available precluded accurate

the species. In an attempt to extend the range of available mixed phosphine–N-oxide complexes, compound (3a) was treated with triphenylphosphine in warm acetone. From analytical data, and from integral values in 360 MHz ¹H n.m.r. spectra of the product, a stoicheiometry of [Pt(SONC₅H₄)₂(PPh₃)]

conductivity measurements, to confirm the ionic nature of

square planar geometries with one chelating and one terminal dithiolate ligand.^{12b,13} Moreover ⁷⁷Se n.m.r. studies of [Pt(Se₂-CNBu¹₂)₂(PR₃)] (R = Et or Ph) provide evidence for a fourco-ordinate molecule in which the chelating and terminal Se₂-CNBu¹₂ ligands undergo exchange *via* a five-co-ordinate intermediate.¹³

In an attempt to prepare the analogues (7b) and (7c) of complex (7a) an alternative synthesis was devised: thus $[PtCl_2(PPh_3)_2]$ and the thallium salt (1b) gave the desired (7a) in moderate yield (52%) and this procedure was elaborated to provide the palladium analogue (7b) in 69% yield. It did not prove possible to synthesise the nickel compound (7c) by this route since $[NiBr_2(PPh_3)_2]$ and $Tl(SC_5H_4NO)$ gave only the known complex (3c) in moderate yield (45%). The spectroscopic features of (7b) are very similar to those of (7a) except that four resonances due to the SC5H4NO ligands are observed in the ¹H n.m.r. spectra down to 217 K where slight broadening is observed. This again suggests rapid exchange between the chelating and terminal SC5H4NO ligands but in this case with a significantly lower energy barrier for the process. This difference has been observed previously in intramolecular exchange processes involving platinum and palladium complexes ¹⁴ and is not unexpected in view of the greater lability generally associated with the latter metal.

^{*} See footnote on p. 783.



In an effort to prepare deoxy analogues of (7a) and (7b), the complexes $[MCl_2(PPh_3)_2]$ (M = Pt or Pd) were separately treated with the thallium(I) salt, thallium pyridine-2-thiolate to give complexes of stoicheiometry [M(SC₅H₄N)₂(PPh₃)₂] [M = Pt (8a), Pd (8b)]. Interestingly the ³¹P n.m.r. spectrum contains two phosphine resonances (integrated ratio 1:1) one of which has a chemical shift identical to that of the free phosphine and in the case of (8a) does not exhibit platinum satellites. Apparently complete phosphine dissociation occurs in solution to generate free PPh3 and (9a) and (9b), the deoxy analogues of (7a) and (7b). This is further supported by the presence of a doublet in the ¹⁹⁵Pt n.m.r. spectrum of (8a), J(Pt-P) = 3.872 Hz. Variable temperature ¹H n.m.r. spectra again suggest fluxional behaviour but in both cases the spectra were insufficiently resolved (even at 360 MHz) clearly to identify the separate resonances undergoing coalescence. Presumably a process related to that involving (7a) and (7b) is involved.

Experimental

Unless otherwise stated i.r. spectra were recorded as CHCl₃ solutions on a Perkin-Elmer 580 spectrophotometer. N.m.r. spectra were recorded in CDCl₃ (unless stated otherwise) on JOEL MH 100, Bruker WP 200 SY, or Bruker WH 360 spectrometers; ¹H (200 or 360 MHz), ³¹P (81 MHz), and ¹⁹⁵Pt (43 MHz) chemical shifts are measured (293 K) relative to SiMe₄, H₃PO₄, and K₂[PtCl₄] respectively. Melting points are uncorrected. Solvents were refluxed over sodium (tetrahydrofuran) or calcium hydride (CH₂Cl₂, Et₂O, and hexane) and distilled under nitrogen prior to use. Acetone was dried over molecular sieves.

Reaction of Zeise's Salt with Quinoline N-Oxide; Preparation of [PtCl₂(CH₂=CH₂)(ONC₉H₇)].—To a stirred solution of Zeise's salt (0.50 g, 1.36 mmol) in water (20 cm³) was added quinoline N-oxide (0.59 g, 4.07 mmol) at room temperature. The yellow-green precipitate was filtered off and recrystallised from chloroform-ethanol to give the pure title compound (0.32 g, 54%), m.p. 158—160 °C (Found: C, 30.05; H, 2.55; Cl, 16.30. C₁₁H₁₁Cl₂NOPt requires C, 30.10; H, 2.50; Cl, 16.15%).

Preparation of the Thallium Salt of 1-Oxido-pyridinium-2thiol, (1b).—The title salt precipitated quantitatively when 1oxidopyridinium-2-thiol (1 mol equivalent) in the minimum amount of methanol or methanol-water was treated dropwise with thallium(1) acetate (1 mol equivalent) in methanol or methanol-water. The precipitate was washed (H₂O then MeOH) and dried to give the yellow salt (1b), m.p. 167— 168 °C (Found: C, 18.40; H, 1.30; N, 4.35. C₃H₄NOSTI requires C, 18.20; H, 1.20; N, 4.25%). I.r. v_{max.} (KBr): 1 580, 1 530, 1 450, 1 405, 1 260, 1 180, 1 125, 1 075, 825, 750, and 695 cm⁻¹.

Reaction of 1-Oxidopyridinium-2-thiol with $K_2[PtCl_4]$ in Acetone; Preparation of Bis(1-oxidopyridinium-2-thiolato)platinum(II) (3a).—Dipotassium tetrachloroplatinate (0.55 g, 1.3 mmol) and 1-oxidopyridinium-2-thiol (0.305 g, 2.4 mmol) 785

were heated under reflux in acetone (25 cm³) for 1 h. The product was filtered off and diethyl ether (20 cm³) was added to the filtrate to precipitate orange (3a) (0.15 g, 27%), m.p. 248—250 °C (from CHCl₃) (Found: C, 27.20; H, 1.75; N, 6.30; O (direct procedure), 6.40. $C_{10}H_8N_2O_2PtS_2$ requires C, 26.85; H, 1.80; N, 6.25; O, 7.15%). I.r. v_{max} : 3 010, 1 610, 1 550, 1 465, 1 425, 1 260, 1 180, 1 145, 1 095, 1 040, 910, and 825 cm⁻¹. N.m.r.: ¹H δ 8.24 (double multiplet, *J ca.* 8, H²), 7.54 (double multiplet, *J ca.* 8, H⁵), 7.24 (triple multiplet, *J ca.* 8, H³ or H⁴), 6.85 p.p.m. (t of d, *J ca.* 8 Hz, H³ or H⁴); ¹⁹⁵Pt δ -1 039 p.p.m.

Reaction of 1-Oxidopyridinium-2-thiol with $K_2[PtCl_4]$ in Water; Preparation of the Oligomer (4).-Dipotassium tetrachloroplatinate (0.76 g, 1.8 mmol), 1-oxidopyridinium-2-thiol (0.68 g, 5.4 mmol), and water (25 cm³) were stirred at room temperature for 1 h. The precipitate was washed with acetone (20 cm³) and then diethyl ether (20 cm³). Evaporation of the filtrate gave orange bis(1-oxidopyridinium-2-thiolato)platinum(II) (3a) (0.15 g) which was spectroscopically (i.r. and n.m.r.) identical with the sample prepared previously. The insoluble material was washed successively with chloroform, acetone, and water to give a yellow-brown insoluble oligomeric material (4) (0.25 g), m.p. ca. 350 °C (decomp.). Samples from different preparative runs gave different analytical data (ranges given are data acquired from four samples unless otherwise stated) (Found: C, 19.10-21.05; H, 1.35-1.70; Cl, 9.10-13.50; N, 4.05-4.35; O (one sample, direct method) 5.65; S (one sample), 9.75. C₂₅H₂₀Cl₃N₅O₅Pt₄S₅ requires C, 22.3; H, 1.50; Cl, 4.40; N, 5.20; O, 5.25; S, 11.90%). I.r. v_{max.} (Nujol mull): 1 610, 1 560, 1 430, 1 280, 1 260, 1 180, 1 150, 1 100, 1 050, 835, 770, and 720 cm⁻¹.

Reaction of Zeise's Salt with 1-Oxidopyridinium-2-thiol.—1-Oxidopyridinium-2-thiol (0.35 g, 2.76 mmol) was added to a stirred solution of Zeise's salt (0.50 g, 1.36 mmol) in acetone (20 cm³) at room temperature. A yellow-brown precipitate was washed with hot chloroform to give a sample of the oligomeric material (4) (0.21 g) described above. The initial filtrate was evaporated to leave bis(1-oxidopyridinium-2thiolato)platinum(II) (0.04 g), m.p. 248—250 °C (from CHCl₃).

Reaction of Oligomeric Material (4) with Triphenylphosphine; Preparation of Chloro(1-oxidopyridinium-2-thiolato)(triphenylphosphine)platinum(II) (5).—The oligomer (4) (0.15 g) and triphenylphosphine (0.16 g) were heated under reflux in ethanol (30 cm³) for 10 min. The solution was evaporated to low volume to precipitate a yellow product, recrystallisation of which from benzene-ethanol gave yellow microcrystalline (5) (0.05 g), m.p. 239—240 °C (Found: C, 44.65; H, 3.10; Cl, 6.20. C₂₃H₁₉CINOPPtS requires C, 44.60; H, 3.05; Cl, 5.75%). I.r. v_{max.}: 3 020, 1 610, 1 552, 1 485, 1 465, 1 440, 1 255, 1 180, 1 145, and 1 100 cm⁻¹. N.m.r.: ¹H δ 8.31 (d, J 7), 7.75 (m), 7.42 (m), 7.20 (m), 6.87 p.p.m. (t of d, J 1.6 and 7 Hz); ³¹P-{¹H} δ 4.25 p.p.m. [J(³¹P-¹⁹⁵Pt) 3 884 Hz]; ¹⁹⁵Pt δ -2 253 p.p.m. [d, J(³¹P-¹⁹⁵Pt) 3 866 Hz].

Reaction of (4) with Dimethylphenylphosphine; Preparation of Chlorobis(dimethylphenylphosphine)(1-oxidopyridinium-2thiolato)platinum(11) (6).—The oligomer (4) (0.125 g) and dimethylphenylphosphine (0.5 g) in acetone (30 cm³) were stirred at room temperature for 1 h. The solvent and excess dimethylphenylphosphine were evaporated to leave an impure white material, recrystallisation of which from acetonediethyl ether gave white microcrystalline (6) (0.08 g), m.p. 127—129 °C (Found: C, 39.90; H, 4.10; Cl, 5.55; S, 5.10. C₂₀H₂₆CINOP₂PtS requires C, 38.70; H, 4.20; Cl, 5.70; S, 5.15%). I.r. v_{max}: 3 020, 2 940, 1 605, 1 555, 1 490, 1 465, 1 440, 1 420, 1 310, 1 295, 1 255, 1 210, 1 185, 1 150, 1 115, 1 095, 950, 915–930, 850, and 830 cm⁻¹. N.m.r.: ¹H δ 8.52 (d, 1 H, J 7), 7.78 (d, 1 H, J 8), 7.45 (m, 12 H, aryl H), 1.88 [d, 6 H, J(P-H) 11.3], 1.79 p.p.m. [d, 6 H, J(P-H) 11.2 Hz]; ³¹P-{¹H} δ 6.78 [J(³¹P-³¹P) 26, J(³¹P-¹⁹⁵Pt) 3 002], -24.83 p.p.m. [J(³¹P-³¹P) 26, J(³¹P-¹⁹⁵Pt) 3 415 Hz].

Preparation of Bis(1-oxidopyridinium-2-thiolato)(triphenylphosphine)platinum(11) (7a).—(a) From (3a) and triphenylphosphine. Complex (3a) (0.14 g, 0.31 mmol) and triphenylphosphine (0.16 g, 0.62 mmol) were warmed in acetone (20 cm³) until all the triphenylphosphine had dissolved. The solution was evaporated to low volume and the yellow precipitate was separated and recrystallised (CHCl₃-EtOH) to give pure (7a) (0.05 g, 23%), m.p. 245-247 °C (Found: C, 47.80; H, 3.40; S, 8.85. C₂₈H₂₃N₂O₂PPtS₂ requires C, 47.40; H, 3.25; S, 9.05%). I.r. v_{max} : 2 970, 1 605, 1 590, 1 551, 1 485, 1 452, 1 440, 1 420, 1 262, 1 250, 1 180, 1 140, 1 100, and 1 092 cm⁻¹. N.m.r. (360 MHz, 220 K); ¹H δ 8.25 (d, 1 H, H²) 8.06 (d, 1 H, H⁵), 7.98 (d, 1 H, H²), 7.63 (m, 6 H, PPh₃), 7.57 (d, 1 H, H^{5'}), 7.35 (m, 9 H, PPh₃), 7.31 (d, 1 H, H³ or H⁴), 6.92 (t, 1 H, H^{3 or 4}), 6.65 (t, 1 H, H^{3' or 4'}), 6.62 (t, 1 H, H^{3' or 4'}). N.m.r. (360 MHz, 298 K): ¹H & 8.07 (d, 2 H, H^{2,2'}), 7.82 (br, 2 H, H^{5,5'}), 7.68 (m, 6 H, PPh₃), 7.35 (m, 9 H, PPh₃), 6.93 (br, 2 H, H^{3,3' or 4,4'}), 6.73 p.p.m. (t, 2 H, H^{3,3' or 4,4'}). N.m.r.: ³¹P-{¹H} δ 6.30 p.p.m [$J(^{13}P^{-195}Pt)$ 3 926 Hz]; ¹⁹⁵Pt $\delta - 1$ 964 p.p.m. [$J(^{195}Pt^{-31}P)$ 3 925 Hz].

(b) From dichlorobis(triphenylphosphine)platinum(II) and the thallium salt (1b). Dichlorobis(triphenylphosphine)platinum(II) (0.50 g, 0.63 mmol) and the thallium salt (1b) (0.42 g, 1.26 mmol) were heated under reflux in chloroform (30 cm³) for 1.5 h. Addition of diethyl ether to the cooled solution gave a precipitate of yellow bis(1-oxidopyridinium-2-thiolato)(triphenylphosphine)platinum(II) (0.23 g, 52%), m.p. 233-236 °C (from CHCl₃-Et₂O). This material was spectroscopically (i.r., ¹H n.m.r.) identical with that described in (a) above.

Preparation of Bis(1-oxidopyridinium-2-thiolato)(triphenylphosphine)palladium(11) (7b).-Dichlorobis(triphenylphosphine)palladium(II) (0.3 g, 0.43 mmol) and the thallium salt (1b) (0.33 g, 1.0 mmol) were stirred in tetrahydrofuran (thf) (40 cm³) at 20 °C for 6 h under nitrogen. Volatiles were evaporated under reduced pressure and the residue extracted with dichloromethane and filtered. Addition of hexane and cooling gave orange-red crystalline (7b) (0.18 g, 69%), m.p. 163-165 °C (darkens at 158 °C) (Found: C, 53.95; H, 4.15; N, 4.80. C₂₈H₂₃N₂O₂PPdS₂ requires C, 54.15; H, 3.70; N, 4.50%). I.r. v_{max} : 3 000, 1 605, 1 590, 1 551, 1 485, 1 455, 1 440, 1 420, 1 268, 1 247, 1 187, 1 167, 1 140, 1 100, and 1 095 cm⁻¹. N.m.r. ¹H & 6.75 (t of d, 2 H, J 2 and 7), 6.90 (t of d, 2 H, J 2 and 7), 7.37 (m, 9 H, PPh₃), 7.63 (dd, 2 H, J 2 and 7), 7.70 (m, 6 H, PPh₃), 8.00 p.p.m. (d, 2 H, J 7 Hz); ³¹P-{¹H} δ 33.46 p.p.m.

Preparation of Bis(pyridine-2-thiolato)bis(triphenylphosphine)platinum(II) (8a).—Dichlorobis(triphenylphosphine)platinum(II) (0.4 g, 0.5 mmol) and thallium pyridine-2thiolate (0.35 g, 1.1 mmol) were stirred under nitrogen in thf (40 cm³) for 15 h. Volatiles were evaporated under reduced pressure and the residue was extracted with dichloromethane and filtered. Addition of diethyl ether and cooling gave yellow (8a) (0.28 g, 66%), m.p. ca. 235 °C (Found: C, 59.15; H, 4.20; N, 3.00; S, 7.20. C₄₆H₃₈N₂P₂PtS₂ requires C, 58.80; H, 4.05; N, 3.00; S, 6.80%). I.r. v_{max}: 3 025, 2 980, 1 585, 1 575, 1 482, 1 450, 1 447, 1 430, 1 410, 1 260, 1 120, and 1 100 cm⁻¹. N.m.r. (CD₂Cl₂): ¹H δ 6.73(br), 7.33(br), 7.64(br), 8.10 p.p.m. (br); J. CHEM. SOC. DALTON TRANS. 1983

³¹P-{¹H} δ 7.96 [J(³¹P-¹⁹⁵Pt) 3 882 Hz], -4.28 p.p.m. (free PPh₃); ¹⁹⁵Pt δ - 2 413 p.p.m. [J(¹⁹⁵Pt-³¹P) 3 872 Hz].

Preparation of Bis(pyridine-2-thiolato)bis(triphenylphosphine)palladium(II) (8b).—Dichlorobis(triphenylphosphine)palladium(II) (0.3 g, 0.43 mmol) and thallium pyridine-2-thiolate (0.3 g, 0.95 mmol) were stirred under nitrogen in thf (40 cm³) for 15 h. Volatiles were evaporated and the residue was extracted with dichloromethane and filtered. Addition of hexane and cooling gave orange (8b) (0.28 g, 76%), m.p. 185—186 °C (Found: C, 63.70; H, 4.45; N, 3.45; P, 7.20; S, 8.85. C₄₆H₃₈N₂P₂PdS₂ requires C, 64.90; H, 4.45; N, 3.30; P, 7.30; S, 7.50%). I.r. v_{max}: 1 585, 1 575, 1 480, 1 450, 1 438, 1 430, 1 410, 1 320, 1 125, and 1 100 cm⁻¹. N.m.r.: ¹H aromatic multiplets at δ 7.77, 7.62, 7.25, and 6.56 p.p.m. (t of d); ³¹P-{¹H} δ 33.4 and -4.38 p.p.m. (free PPh₃).

Preparation of Bis(1-oxidopyridinium-2-thiolato)nickel(II) (3c).—Dibromobis(triphenylphosphine)nickel(II) (0.2 g 0.27 mmol) and thallium pyridine-2-thiolate (0.2 g 0.6 mmol) were stirred under nitrogen in thf (40 cm³) for 2 h. Volatiles were evaporated and the residue was extracted with dichloromethane and filtered. Addition of diethyl ether and cooling gave brown (3c) (38 mg, 45%), m.p. 258—260 °C (decomp.) [lit.,¹⁰ 260 °C (decomp.)] (Found: C, 38.90; H, 2.70; N, 9.20. C₁₀H₈N₂NiO₂S₂ requires C, 38.60; H, 2.25; N, 9.00%). I.r. v_{max} : 3 020, 1 605, 1 565, 1 465, 1 430, 1 250, 1 190, and 1 150 cm⁻¹. N.m.r.: ¹H δ 8.01 (d of m, J 7), 7.36 (d of m, J 7), 7.11 (t of d, J 7 and 1), 6.79 p.p.m. (J 7 and 1 Hz).

Acknowledgements

We thank the S.E.R.C. and the British Council for financial support. One of us (M. V. R.) thanks Professor A. Furlani for arranging leave of absence from the University of Rome and for helpful discussions.

References

- 1 See R. Uson, L. A. Oro, M. A. Ciriano, and F. Lahoz, J. Organomet. Chem., 1981, 217, 251 and refs. therein.
- 2 Y. Ozaki, S. Matsuoka, and M. Magami, Jap. P. 26 768/ 1973; Chem. Abstr., 1973, 79, P42694 r.
- 3 C. Bouillon and G. Rosenbaum, G. P. 2 832 010/1979; Chem. Abstr., 1979, 90, 168466.
- 4 R. E. Muntwyler, G. P. 2717 041/1977; Chem. Abstr., 1978, 88, 37623 y.
- 5 J. L. Davidson, I. E. P. Murray, P. N. Preston, M. V. Russo, Lj. Manojlovic-Muir, and K. W. Muir, J. Chem. Soc., Chem. Commun., 1981, 1059.
- 6 L. Garcia, S. I. Shupeck, and M. Orchin, *Inorg. Chem.*, 1962, 1, 893.
- 7 A. R. Katritzky and J. M. Lagowski, 'Chemistry of the Heterocyclic N-Oxides,' Academic Press, New York, 1971, p. 443.
- 8 R. A. Jones and A. R. Katritzky, J. Chem. Soc., 1960, 2937.
- 9 M. A. Robinson, J. Inorg. Nucl. Chem., 1964, 26, 1277.
- 10 J. E. Lydon, M. R. Truter, and C. R. Watling, Proc. Chem. Soc., 1964, 193; J. E. Lydon and M. R. Truter, J. Chem. Soc., 1965, 6899.
- 11 P. S. Pregosin and R. S. Kunz, in 'N.m.r. Basic Principles and Progress No. 16,' Springer, Berlin, 1979, p. 94.
- 12 (a) J. M. C. Alison and T. A. Stephenson, J. Chem. Soc., Dalton Trans., 1973, 254 and refs. therein; (b) M. C. Corncock and T. A. Stephenson, *ibid.*, 1977, 501 and refs. therein.
- 13 J. P. Fackler, jun. and W.-H. Pan, J. Am. Chem. Soc., 1979, 101, 1607.
- 14 K. R. Dixon, Inorg. Chem., 1977, 16, 2618; J. B. Brandon, M. Collins, and K. R. Dixon, Can. J. Chem., 1978, 56, 950.

Received 21st June 1982; Paper 2/1041