


The Bartlett mechanism has been widely used to rationalize the results of olefin epoxidations by peroxides, and this determination of the disposition of the atoms in the transition structure can be reasonably extended to related oxygen-transfer reactions.



Acknowledgment. We are grateful to the National Science Foundation and the National Institutes of Health for support of this work.

Supplementary Material Available: Experimental details and spectral data for compounds 1, 2, 7, and 8 and experimental procedures for double-label crossover experiments and control studies to rule out label scrambling (6 pages). Ordering information is given on any current masthead page.

(9) We note that our work does not distinguish between spiro and planar arrangements of the olefin, a topic discussed in detail by Rebek. See: Rebek, J., Jr.; Marshall, L.; McManis, J.; Wolak, R. *J. Org. Chem.* 1986, 51, 1649.

Expanded Heterohelices: Molecular Coils That Form Chiral Complexes

Thomas W. Bell* and Hélène Jousselin

Department of Chemistry, State University of New York
Stony Brook, New York 11794-3400

Received April 11, 1991

Helical structures are ubiquitous in nature and are essential to life itself (e.g., polypeptides and nucleic acids). In recent years chemists have discovered several metal-ligand combinations that lead to spontaneous assembly of multicomponent helical and double-helical structures.¹ In this communication, we report the first examples of a new class of preorganized, monohelical polypyridine ligands. These "molecular coils" composed of fused six-membered rings wrap around guest ions, forming helical 1:1 complexes. Helicenes² such as [6]helicene (**2**) consist only of angularly fused rings; their ideal, planar analogue is coronene (**1**). Combination of angular with linear fusion of benzene rings leads to planar cycloarenes³ (e.g., kekulene, **3**) and conceptually to unknown, "expanded" carbocyclic helicenes, such as **4**. Re-




Figure 1. Syntheses of expanded heterohelices: (a) $\text{Me}_2\text{NCH}_2\text{Cl}$, CH_3CN (95%); (b) NaOH , H_2O , $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{I}/\text{Et}_3\text{N}$, CH_2Cl_2 (78%); (c) DMSO (55%); (d) O_3 , CH_2Cl_2 , $\text{CH}_3\text{OH}/\text{Me}_2\text{S}$ (92%); (e) NH_4OAc , DMSO , 95 °C (22%); (f) $t\text{-BuOCH}(\text{NMe}_2)_2$, 80 °C (82%); (g) NH_4BF_4 , DMF , 150 °C.




Figure 2. UV-visible absorption spectra of **12** and **12.Na⁺**.

placement of internal C-H groups in cycloarenes with heteroatoms gives torands,⁴ such as **5**,^{4b-e,5} which tightly bind alkali metals.^{4c}

Both of the new "expanded" heterohelices (**6** and **12**) were synthesized from 9-*n*-butyl-2,3,5,6,7,8-hexahydro-4(1*H*)-acridinone (**7**) and 5-benzylidene-9-*n*-butyl-2,3,5,6,7,8-hexa-


(1) (a) Constable, E. C.; Holmes, J. M.; Raithby, P. R. *Polyhedron* 1991, 10, 127-132. (b) Lehn, J.-M. *Angew. Chem., Int. Ed. Engl.* 1990, 29, 1304-1319. (c) Constable, E. C.; Elder, S. M.; Healy, J.; Ward, M.; Tocher, D. A. *J. Am. Chem. Soc.* 1990, 112, 4590-4592. (d) Constable, E. C.; Ward, M. D.; Tocher, D. A. *J. Am. Chem. Soc.* 1990, 112, 1256-1258. (e) Koert, U.; Harding, M. M.; Lehn, J.-M. *Nature* 1990, 346, 339-342. (f) Constable, E. C. *Nature* 1990, 346, 314-315. (g) Dietrich-Buchecker, C. O.; Sauvage, J.-P.; *Angew. Chem., Int. Ed. Engl.* 1989, 28, 189-192. (h) Lehn, J.-M.; Rigault, A. *Angew. Chem., Int. Ed. Engl.* 1988, 27, 1095-1097. (i) Lehn, J.-M.; Rigault, A.; Siegel, J.; Harrowfield, J.; Chevrier, B.; Moras, D. *Proc. Natl. Acad. Sci. U.S.A.* 1987, 84, 2565-2569. (j) Vögtle, F.; Weber, E. *Angew. Chem., Int. Ed. Engl.* 1979, 18, 753-776.

(2) (a) Laarhoven, W. H.; Prinsen, W. J. C. In *Topics in Current Chemistry, Vol. 125, Stereochemistry*; Vögtle, F., Weber, E., Eds.; Springer-Verlag: New York, 1984; pp 63-130. (b) Meurer, K. P.; Vögtle, F. In *Topics in Current Chemistry, Vol. 127, Organic Chemistry*; Boschke, F. L., Ed.; Springer-Verlag: New York, 1985; pp 1-76. See also: (c) Gupta, R. B.; Kaloustain, M. K.; Franck, R. W.; Blount, J. F. *J. Am. Chem. Soc.* 1991, 113, 359-361.

(3) (a) Funhoff, D. J. H.; Staab, H. A. *Angew. Chem., Int. Ed. Engl.* 1986, 25, 742-744. (b) Staab, H. A.; Diederich, F.; Caplar, V. *Liebigs Ann. Chem.* 1983, 2262-2273. (c) Staab, H. A.; Diederich, F.; Krieger, C.; Schweitzer, D. *Chem. Ber.* 1983, 116, 3504-3512. (d) Staab, H. A.; Diederich, F. *Chem. Ber.* 1983, 116, 3487-3503. (e) Diederich, F.; Staab, H. A. *Angew. Chem., Int. Ed. Engl.* 1978, 17, 372-374.

(4) (a) Bell, T. W.; Liu, J. *Angew. Chem., Int. Ed. Engl.* 1990, 29, 923-925. (b) Bell, T. W.; Firestone, A.; Liu, J.; Ludwig, R.; Rothenberger, S. D. In *Inclusion Phenomena and Molecular Recognition*; Atwood, J. L., Ed.; Plenum Press: New York, 1990. (c) Bell, T. W.; Firestone, A.; Ludwig, R. *J. Chem. Soc., Chem. Commun.* 1989, 1902-1904. (d) Bell, T. W.; Firestone, A.; Guzzo, F.; Hu, L.-Y. *J. Inclusion Phenom.* 1987, 5, 149-152. (e) Bell, T. W.; Firestone, A. *J. Am. Chem. Soc.* 1986, 108, 8109-8111.

(5) Dodecahydrohexaazakekulene: Ransohoff, J. E. B.; Staab, H. A. *Tetrahedron Lett.* 1985, 26, 6179-6182.



hydro-4(1*H*)-acridinone (**8**) by variations of previously reported methods^{4b,d,e} (Figure 1). These intermediates were synthesized in several steps from cyclohexanone and valeraldehyde by published methods.^{4b,d,e,6} Ketone **8** was converted to enone **9** by a new coupling procedure⁷ using dimethylmethyleammonium chloride.⁸ Reaction of ketone **7** with enone **9** gave unsymmetrical heptacyclic terpyridyl **10**.⁹ Ozonolytic cleavage of the benzylidene group in **10** gave heptacyclic ketone **11**,⁹ the key intermediate for synthesis of expanded heterohelices. Homologation of **11** by reaction with ammonium acetate and **9** and chromatography of the crude product on basic alumina gave free ligand **12** (22%)⁹ and a later fraction, which proved to be a sodium complex (**12**·Na⁺).¹⁰ Reaction of **11** with Brederick's reagent¹¹ gave β -dimethylamino enone **13**, which was coupled with **11** by the method of Firestone.¹² Chromatography on basic alumina again gave two fractions, apparently containing free ligand **6** and **6**·Na⁺.¹³

Geminal protons of **6** and **12** become diastereotopic when their enantiomeric helical conformations interconvert slowly. In free host **12** various methylene groups are observed as distinct two-proton multiplets in the 600-MHz ¹H NMR spectrum, indicating that helix inversion is rapid on the NMR time scale. The ¹H NMR spectrum of **12**·NaCF₃SO₃, prepared by treatment of **12** with sodium triflate in methanol, shows broadened CH₂ peaks due

to slow helix inversion; the benzylidene peak remains sharp but is shifted upfield by 2.0 ppm. These results show not only that helix inversion is slow in **12**·Na⁺ but also that complexation alters the helix conformation. Figure 2 shows that a large bathochromic shift of the longest wavelength UV absorption also occurs upon binding sodium.

We conclude that expanded helix **12** consists of rapidly interconverting enantiomers in solution; complexation of sodium increases the barrier to helix inversion. This potentially useful conformational change and the resulting optical response are more extreme than those observed in expanded systems comprising less than one full turn of a helix.^{4a} Complexation and conformational properties of **6** are currently under investigation.¹⁴

Acknowledgment. We thank Richard Ludwig for recording 600-MHz ¹H NMR spectra and for assistance with preparing figures. This project was supported by the National Institutes of Health (GM 32937). The Bruker AMX-600 NMR instrument was obtained with grants from the NIH (RR05547A) and the NSF (CHE 8911350) and with support from the Center for Biotechnology and SUNY, Stony Brook.

(14) During the course of this research we learned of efforts by G. Balavoine et al. (Université de Paris-Sud, Orsay) to prepare compounds of related structure.

Traveling Fronts of Methacrylic Acid Polymerization

John A. Pojman

Department of Chemistry and Biochemistry
University of Southern Mississippi
Hattiesburg, Mississippi 39406-5043

Received April 29, 1991

We report here the preliminary investigation of traveling fronts in methacrylic acid polymerization in unstirred solutions of the monomer and benzoyl peroxide, with and without a promoter.

An autocatalytic reaction in an unstirred vessel can support a constant-velocity wave front resulting from the coupling of diffusion to the chemical reaction. Numerous reactions in solution have been described in which a front of chemical reactivity propagates through the medium from the site of an initial concentration perturbation.¹⁻¹¹ Traveling wave fronts in populations of short self-replicating RNA variants have been created in thin capillary tubes,¹² but fronts have not been studied in synthetic polymerization reactions. We have set out to do so in an attempt to observe ultimately the sorts of nonlinear propagation phenomena seen in solid-fuel, gasless combustion such as pulsating and spinning fronts.¹³⁻¹⁵

Coupling the exothermic addition polymerization of methacrylic

(6) (a) Bell, T. W.; Cho, Y.-M.; Firestone, A.; Healy, K.; Liu, J.; Ludwig, R.; Rothenberger, S. D. *Org. Synth.* **1990**, *69*, 226-237. (b) Bell, T. W.; Rothenberger, S. D. *Tetrahedron Lett.* **1987**, *28*, 4817-4820. (c) Bell, T. W.; Firestone, A. J. *Org. Chem.* **1986**, *51*, 764-765.

(7) Bell, T. W.; Firestone, A.; Kwok, D.-I.; Ludwig, R., manuscript in preparation.

(8) Risch, N.; Esser, A. *Synthesis* **1988**, 337-339.

(9) Satisfactory ¹H NMR, ¹³C NMR, IR, MS, UV-visible, and C, H, N microanalytical data were obtained.

(10) The ¹H NMR spectrum of this material was identical with that of **12**·NaCF₃SO₃. Continuous extraction of its solution in CHCl₃ with pure water gave free **12**.

(11) Brederick, H.; Simchen, G.; Rebsdat, S.; Kantlehner, W.; Horn, P.; Wahl, R.; Hoffman, H.; Grieshaber, P. *Chem. Ber.* **1968**, *101*, 41-50.

(12) Firestone, A. Ph.D. Thesis, SUNY, Stony Brook, 1988.

(13) As observed for **12** and **12**·Na⁺, the ¹H NMR spectrum of the first-eluted fraction was less broad than that of the later fraction. The principal high-mass peak in the FAB-MS spectra of both fractions was *m/z* 1052 (*M* + 23).

(1) Zaikin, A. N.; Zhabotinskii, A. M. *Nature* **1970**, *225*, 535-537.
(2) Winfree, A. T. *Science* **1973**, *181*, 937-939.
(3) Field, R. J.; Noyes, R. M. *J. Am. Chem. Soc.* **1974**, *96*, 2001-2006.
(4) Reusser, E. J.; Field, R. J. *J. Am. Chem. Soc.* **1979**, *101*, 1063-1071.
(5) Gribshaw, T. A.; Showalter, K.; Banville, D. L.; Epstein, I. R. *J. Phys. Chem.* **1981**, *85*, 2152-2155.
(6) Baza, G.; Epstein, I. R. *J. Phys. Chem.* **1985**, *89*, 3050-3053.
(7) Hanna, A.; Saul, A.; Showalter, K. *J. Am. Chem. Soc.* **1982**, *104*, 3838-3844.
(8) *Oscillations and Traveling Waves in Chemical Systems*; Field, R. J., Burger, M., Eds.; Wiley: New York, 1985.
(9) Harrison, J.; Showalter, K. *J. Phys. Chem.* **1986**, *90*, 225-226.
(10) Szivoczka, L.; Nagypal, I.; Boga, E. *J. Am. Chem. Soc.* **1989**, *111*, 2842-2845.
(11) Ross, J.; Müller, S. C.; Vidal, C. *Science* **1988**, *240*, 460-465.
(12) Bauer, G. J.; McCaskill, J. S.; Otten, H. *Proc. Natl. Acad. Sci. U.S.A.* **1989**, *86*, 7937-7941.
(13) Kaper, H. G.; Leaf, G. K.; Margolis, S. B.; Matkowsky, B. J. *Combust. Sci. Technol.* **1987**, *53*, 289-314.
(14) Anselm-Tamburini, U.; Munir, Z. A. *J. Appl. Phys.* **1989**, *66*, 5039-5045.
(15) Matkowsky, B. J.; Sivashinsky, G. I. *SIAM J. Appl. Math.* **1978**, *35*, 465-478.