


SYNTHETIC STUDIES ON CLERODANE DITERPENOIDS. 3. AN ALTERNATIVE SYNTHESIS OF AN UNNATURAL ANTIFEEDANT^{1,2}

Hsing-Jang Liu* and Naim H. Al-Said

Department of Chemistry, University of Alberta, Edmonton, Alberta,
Canada T6G 2G2


Abstract – An alternative synthesis of (1S*,2S*,6R*,10S*)-2-acetoxy-1-acetoxymethyl-5,5-dimethylbicyclo[4.4.0]decane-10-spiro-2'-oxirane (**1**), an unnatural antifeedant, has been effected *via* an intermolecular Diels-Alder approach.

During the investigation of clerodane diterpenoids, Ley *et al.*³ observed that (1S*,2S*,6R*,10S*)-2-acetoxy-1-acetoxymethyl-5,5-dimethylbicyclo[4.4.0]decane-10-spiro-2'-oxirane (**1**), a simple analogue of the natural antifeedant clerodin,⁴ also showed interesting insect antifeedant activity. They prepared this compound in eight synthetic operations starting from 4,4-dimethyl-2-cyclohexenone.³ We wish to report herein an alternative synthesis of this unnatural antifeedant.




Our approach to the requisite *cis*-decalin system possessing appropriate oxidation level at various key centers was based on an intermolecular Diels-Alder reaction. This was effected by the addition of dienone ester (**2**)⁵ to diene (**3**).⁶ Under zinc chloride catalysis the reaction

occurred readily and efficiently. After 48 h at room temperature in methylene chloride using one part each of the diene and dienophile, adduct (**4**) was isolated in 71% yield. Subsequent reduction of adduct (**4**) with lithium aluminum hydride gave a 65% yield of the desired diol (**5**) which was acetylated with acetic anhydride in tetrahydrofuran in the presence of pyridine and 4-dimethylaminopyridine. Diacetate (**6**) thus obtained in quantitative yield was subjected to debenzoylation with 10% palladium on carbon under an atmosphere of hydrogen.^{8,9} Further hydrogenation of the resulting alcohol (**7**) using platinum as a catalyst resulted in the formation of the corresponding saturated alcohol (**8**) in 97% yield over two steps.



Scheme 1


An alternative approach to this key intermediate (**8**) has also been developed (Scheme 2). In this approach, 2,4-pentadienol (**9**)⁶ was used as the diene in the initial Diels-Alder reaction. Its addition to dienone (**2**) under zinc chloride catalysis gave adduct (**10**) in nearly



quantitative yield. Interestingly, when compound (**10**) was treated with a trace amount of *p*-toluenesulfonic acid in refluxing methanol, the corresponding methyl ether (**11**) was formed in 85% yield, presumably *via* the intermediacy of oxonium ion (**11a**). Reduction of **11** with

lithium aluminum hydride followed by catalytic hydrogenation of the resulting diol (**12**) using platinum gave rise to saturated diol (**13**) (66% overall yield). Diol (**13**) was acetylated with acetic anhydride in pyridine in the presence of a catalytic amount of 4-dimethylaminopyridine to give the corresponding diacetate (**14**) (82% yield) which was subjected to demethylation with *B*-bromo-9-borobicyclo[3.3.1]nonane¹⁰ followed by brief treatment with aqueous sodium hydroxide to furnish the desired alcohol (**8**) (85% yield).

The conversion of alcohol (**8**) to the target compound (**1**) was carried out as follows (Scheme 3). Treatment of **8** with *tri-n*-butylphosphine and *N*-(phenylselenenyl)phthalimide (PSP)¹¹ in tetrahydrofuran effected the replacement of its hydroxy group with a phenylselenenyl group, enroute to the installation of the required double bond. Subsequent oxidation of the resulting selenide (**15**) (94% yield) using *m*-chloroperbenzoic acid followed by an elimination reaction carried out in refluxing carbon tetrachloride containing a small amount of diethylamine gave rise to the desired olefin (**16**). Finally, epoxidation of **16** with *m*-chloroperbenzoic acid furnished the target compound (**1**), of which the spectroscopic data were found to be in good agreement with those reported.³ In addition to providing an effective alternative synthesis of an interesting insect antifeedant, the current work also served as a model study for the construction of clerodin and related clerodane diterpenoids.



Scheme 3

ACKNOWLEDGEMENTS

We are grateful to the Natural Sciences and Engineering Research Council of Canada and the University of Alberta for financial support.

REFERENCES AND NOTES

1. For Part 2, see H. J. Liu, K.S. Shia, Y. Han, D. Sun, and Y. Wang, *Can. J. Chem.*, in press.
2. This paper is dedicated in memory of Professor Shun-ichi Yamada.
3. W. P. Jackson and S. V. Ley, *J. Chem. Soc., Perkin Trans. I*, 1981, 1516.
4. D. Roger, G. G. Unal, D. J. Williams, S. V. Ley, G. A. Sim, B. S. Joshi, and K. R. Ravindranath, *Chem. Commun.*, 1979, 97.
5. H. J. Liu and E. N. C. Browne, *Can. J. Chem.*, 1987, **65**, 1262.
6. This compound was prepared in 62% yield by benzylation (NaH, BnBr, THF, rt, 24 h) of the corresponding alcohol which, in turn, was prepared in 70% yield by esterification (MeOH, H₂SO₄, rt, 16 h) of 2,4-pentadienoic acid⁷ followed by reduction (LiAlH₄, Et₂O, rt, 16 h).
7. A. D. Mebane, *J. Am. Chem. Soc.*, 1952, **74**, 5227.
8. C. H. Heathcock and R. Ratcliffe, *J. Am. Chem. Soc.*, 1971, **93**, 1746.
9. It is noteworthy that, under these conditions, the neopentyl double bond was intact.
10. M. V. Bhay, *J. Organomet. Chem.*, 1978, **156**, 221.
11. P. A. Grieco, J. Y. Jaw, D. A. Clare, and K. C. Nicolaou, *J. Org. Chem.*, 1981, **46**, 1215.

Received, 16th May, 1997