## Nonconcerted Paths for Reactions of Alkene–Zirconocene Complexes

Ei-ichi Negishi,\* Danièle Choueiry, Thinh B. Nguyen, and Douglas R. Swanson

> Department of Chemistry, Purdue University West Lafayette, Indiana 47907

Noriyuki Suzuki and Tamotsu Takahashi\*

Coordination Chemistry Laboratory Institute for Molecular Science Myodaiji, Okazaki 444, Japan

Received September 10, 1993

Concerted reaction paths have been implicated in various reactions of alkyne-zirconocene derivatives including benzynezirconocenes.<sup>1</sup> Thus, for example,  $Cp_2ZrPh_2$ , where  $Cp = \eta^5$ - $C_5H_5$ , was reported to react with (E)- and (Z)-stilbenes to give stereospecifically 1a and 1b, respectively.<sup>2</sup> Similarly, the ZrCp<sub>2</sub>promoted bicyclization reaction of 2 proceeds with stereochemical retention to give the corresponding isomers of 3.3 On the other



hand, the stereochemistry of reactions of alkene-zirconocene derivatives with respect to the E-Z configuration of the starting alkenes has remained essentially unknown.

We now report that, although the reactions of alkenezirconocenes can, in some cases, proceed stereospecifically, they can also undergo stereoisomerization leading to nonstereospecific but often highly stereoselective processes, which most likely proceeds via novel dipolar zirconate species represented by 4.

$$c_{\overline{ZrCp_2L_n}}$$
 4 (L<sub>n</sub> = ligands and ZrCp<sub>2</sub>-containing groups)

The reaction at -78 to 23 °C of **5a** and **5b** with *n*-Bu<sub>2</sub>ZrCp<sub>2</sub>, generated in situ by treating Cl<sub>2</sub>ZrCp<sub>2</sub> with n-BuLi in THF at -78 °C,<sup>4</sup> produced 6a and 6b, respectively. Compound 6a was obtained in 85% NMR yield as a >96% stereoisomerically pure species, while 6b was obtained in 93% NMR yield, as a 10:1 mixture of 6b and 6a after 1 h at 23 °C. Carbonylation of 6a and 6b with 1.1 atm of CO at 0 °C followed by iodinolysis<sup>5</sup> gave 7a and 7b, respectively. Compound 7a was isolated in 70% yield as a >97% stereoisomerically pure species, while the product derived from 6b was 90-95% isomerically pure 7b contaminated with 7a (5-10%). The stereochemical assignments of these

(2) Kropp, K.; Erker, G. Organometallics 1982, 1, 1246. See, however,
(2) Kropp, K.; Erker, G. Organomet. Chem. 1982, 224, 29.
(3) Negishi, E.; Swanson, D. R.; Cederbaum, F. E.; Takahashi, T. Tetrahedron Lett. 1987, 28, 917. See also McDade, C.; Bercaw, J. E. J.

Organomet. Chem. 1985, 279, 281.

(4) (a) Negishi, E.; Cederbaum, F. E.; Takahashi, T. Tetrahedron Lett. 1986, 27, 2829. (b) Negishi, E.; Holmes, S. J.; Tour, J. M. Miller, J. A.; Cederbaum, F. E.; Swanson, D. R.; Takahashi, T. J. Am. Chem. Soc. 1989, 1, 3336

(5) Swanson, D. R.; Rousset, C. J.; Negishi, E.; Takahashi, T.; Seki, T.; Saburi, M.; Uchida, Y. J. Org. Chem. 1989, 54, 3521.

compounds were in part based on NMR spectroscopy. The identity of 7a was confirmed by its X-ray analysis. The unusual trans stereochemistry for ring fusion<sup>6</sup> was further established by synthesizing the two cis-fused isomers 7c and 7d from previously reported 2-phenylbicyclo[3.3.0]oct-1-en-3-one4b by its reduction with 3 equiv of LiAlH(OMe)<sub>3</sub> and 1.5 equiv of CuBr<sup>7</sup> and hydrogenation with  $H_2$  (10 psi) over Pd/BaSO<sub>4</sub> in EtOAc at 25 °C, respectively. These results clearly demonstrate that the formation of zirconacyclopentane derivatives via ZrCp<sub>2</sub>promoted alkene-alkene coupling can proceed stereospecifically.



Surprisingly, the reaction of  $Et_2ZrCp_2$  with (E)- or (Z)- $\beta$ methylstyrene was >98% stereoselective but nonstereospecific, producing 8a in high yields. Its deuterolysis gave 9a, which was  $\geq$ 96% erythro.<sup>8</sup> The sharply contrasting results in eqs 2 and 3 prompted us to further examine the reaction shown in eq 2, which indeed revealed a slow but significant isomerization of 6b into 6a. Over 48 h at 23 °C, the ratio of 6a to 6b changed from 1:10 to 9:1. These results clearly indicate that the thermodynamically less stable stereoisomer of zirconacyclopentane derivatives can isomerize to the more stable stereoisomer. To our knowledge, this kind of stereoisomerization of metallacyclopentanes appears to be unprecedented.

$$Et_{2}ZrCp_{2} \xrightarrow{(Z) \cdot or (E) - PhCH = CHMe} ZrCp_{2} \xrightarrow{(Z) \cdot or (E) - PhCH = CHMe} Me_{A} \xrightarrow{Ph}_{D} D (3)$$

$$8a \qquad 9a \qquad (3)$$

We have further found that, in addition to (Z)-stilbene,<sup>3</sup> various olefins, such as (Z)-PhCH=CHMe, (Z)-PhCH=CHSiMe<sub>3</sub>, and (E)-cyclooctene,<sup>9</sup> undergo stereoisomerization catalyzed by n-Bu<sub>2</sub>ZrCp<sub>2</sub>. A 1:1 mixture of (Z)-PhCH=CHPh and (Z)-PhCD=CDPh gave a 1:1 mixture of the corresponding E isomers under the influence of n-Bu<sub>2</sub>ZrCp<sub>2</sub> without a sign of H-D scrambling. Coupled with the fact that there is no allylic hydrogen, two commonly observed transition-metal-mediated stereoisomerization processes, i.e., hydrometalation-dehydrometalation and allylic C-H oxidative addition-reductive elimination,<sup>10</sup> are ruled

- (6) (a) Rousset, C. J.; Swanson, D. R.; Lamaty, F.; Negishi, E. Tetrahedron Lett. 1989, 30, 5105. (b) Nugent, W. A.; Taber, D. F. J. Am. Chem. Soc. 1989, 111, 6435.
- (7) Semmelhack, M. F.; Stauffer, R. D.; Yamashita, A. J. Org. Chem. 1977. 42. 3180.
- (8) The threo isomer of 9a was prepared as follows for spectral comparison.



(9) For its preparation, see: Carbally, R. P.; Perkins, M. J.; Elnitski, A. P. J. Chem. Soc., Perkin Trans. 1 1979, 793.

(10) For papers presenting such mechanisms for a different Ti- or Zrcatalyzed alkene isomerization reaction, see: (a) Akita, M.; Yasuda, H.; Nagasuna, K.; Nakamura, A. Bull. Chem. Soc. Jpn. 1983, 56, 554. (b) Maye, J. P.; Negishi, E. Tetrahedron Lett. 1993, 34, 3359.

© 1994 American Chemical Society

<sup>(1)</sup> For reviews, see: (a) Negishi, E.; Takahashi, T. Aldrichimica Acta 1985, 18, 31. (b) Negishi, E.; Takahashi, T. Synthesis 1988, 1. (c) Negishi, E. Chem. Scr. 1989, 29, 457. (d) Buchwald, S. L.; Nielsen, R. B. Chem. Rev. 1988, 88, 1047

out. However, isomerization of simple alkyl-substituted alkenes is indeed complicated and even dominated by a more conventional regio- and stereoisomerization process.<sup>10</sup> A detailed NMR study of the isomerization reaction of (Z)-stilbene in the presence of 1 equiv of preformed (E)-(PhCH=CHPh)ZrCp<sub>2</sub> (10) has revealed that the stereoisomerization reaction is first order in (Z)-stilbene and second order in 10. Critically important here is that this isomerization proceeds associatively.



In principle, alkene stereoisomerization may involve one or more of C-H and  $\sigma$  and/or  $\pi$  C-C bond cleavages. To rigorously probe this point, we chose (Z)-PhCD=CHMe<sup>11</sup> (11) as a readily accessible, regio- and stereodefined Z olefin containing four different groups around the C=C bond. Its stereoisomerization using 10 mol % of n-Bu<sub>2</sub>ZrCp<sub>2</sub> produced the E isomer 12 without a sign of the formation of any species arising from changes in atom-linking sequences. The results rule out all but the  $\pi$ -bond cleavage mechanisms, for which the homolytic and heterolytic extremes may be considered. Neither Me<sub>2</sub>ZrCp<sub>2</sub> nor tetramethylenezirconocene in the presence of an excess of ethylene catalyzed Z-to-E isomerization of stilbene at 23 °C. In fact, no reaction was observed. These results indicate that as such neither dialkylzirconocenes nor zirconacyclopentanes are isomerization catalysts. Since <sup>1</sup>H NMR analysis of the isomerization reaction mixtures revealed the presence of alkene-ZrCp<sub>2</sub> complexes, e.g., 10, as the predominant  $ZrCp_2$  derivatives, they most likely serve as the active catalysts. In order for dialkylzirconocenes and zirconacyclopentanes to act as catalyst precursors, they must therefore be convertible to alkene-ZrCp<sub>2</sub> complexes. To further probe the mode of interaction between an alkene and an alkene-ZrCp<sub>2</sub>, especially the sign and extent of charge buildup on an alkenyl C and Zr, p,p'-disubstituted (Z)-stilbenes (>98% Z) containing Me, F, Cl, and  $CF_3$  (13) were prepared by partial reduction of the corresponding tolans which, in turn, were prepared by the  $Pd(PPh_3)_4$ -catalyzed reaction of  $(-ZnC=C-)_n$  with any l iodide (2 equiv).<sup>12</sup> Isomerization of 13 was carried out in 2:1 THF-C<sub>6</sub>D<sub>6</sub> at 23 °C using the corresponding stilbene-ZrCp<sub>2</sub> generated in situ from 1 molar equiv each of 13 and (n-Bu)<sub>2</sub>ZrCp<sub>2</sub> as the isomerization agent, and its progress was monitored by <sup>1</sup>H NMR spectroscopy. Under these conditions 13e remained essentially unchanged over several hours. In the other cases, the signals for the E isomer emerged and grew at the expense of those of the Z isomer, while those for the (E)-stilbene-ZrCp<sub>2</sub> remained essentially constant. No other stilbene-containing or -derived species were detectable. The pseudo-first-order rate constants observed with 13a-d are  $1.19 \times 10^{-3}$  (13a),  $1.84 \times 10^{-3}$  (13b),  $4.56 \times 10^{-4}$  (13c), and  $2.79 \times 10^{-5}$  (13d) s<sup>-1</sup>. Although a clearcut linear free energy relationship is not observed, it is not surprising, considering the reactant structures and the nature of interactions. Nonetheless, the overall trend in substituent effects seems unmistakable and points to the involvement of benzyl cationic species. One associative process we initially considered involves 14 as a transient species. However, the second-order rate dependence<sup>13</sup> on Cp<sub>2</sub>Zr(PhCH=CHPh) is clearly inconsistent with this view. Our recent elucidation of the structure of  $(Cp_2ZrMe)_2(CH_2=CH_2)$  (15) by NMR and X-ray analyses<sup>14</sup> prompts us to propose 16 as an active species consistent with all of the data presented herein.<sup>15</sup> The modes of alkene-ZrCp<sub>2</sub> interaction and alkene stereoisomerization reflected in 14 and 16 are essentially the same. Significantly, however, dimerization of Cp<sub>2</sub>Zr(PhCH=CHPh) as depicted in 17 represents a plausible, novel mode of activation of one Zr center as an electrophile by the other. The opportunity for possible, extensive electron delocalization in 16 may also favor the proposed 2:1 interaction.

In summary, the reaction of alkene-zirconocenes with alkenes can, in some cases, lead to the formation of zirconacyclopentanes via concerted C-Zr bond addition to alkenes, which is thought to involve donation of alkene  $\pi$ -electrons and back-donation of C-Zr bonding electrons.<sup>1</sup> In cases where the back-donation is hindered due to steric and other factors, the alkene-zirconocenes presumably act merely as electron-accepting Lewis acids, leading to the formation of zirconates represented by 4. This process can be facilitated through dimeric interaction as in 16. In addition to inducing E-Z isomerization, 4 could, in some cases, ringexpand to provide zirconacyclopentanes via a polar path, for which the reaction shown in eq 3 is a possible candidate. These polar processes represent yet another novel reaction pattern that ZrCp<sub>2</sub> derivatives display and show some striking contrasts between alkene-zirconocenes and their alkyne counterparts.

Acknowledgment. We thank the National Science Foundation (CHE-9402288) for support of this research. We also thank Dr. P. E. Fanwick for X-ray analysis and the NMR Center of the Chemistry Department for technical assistance. D.C. is a graduate research fellow (1992-1993) on funds provided by the Purdue Research Foundation. We acknowledge that Professors Y. Uchida and M. Saburi as well as Mr. M. Murakami participated in some related aspects of this research.

Supplementary Material Available: Experimental details including the stereoisomerization of (Z)-stilbenes and IR and NMR data for 5a,b, 6a,b, 7a-d, 9a, and other compounds (11 pages). This material is contained in many libraries on microfiche, immediately follows this article in the microfilm version of the journal, and can be ordered from the ACS; see any current masthead page for ordering information.

(13) As suggested by a referee, the apparent second-order dependence on  $Cp_2Zr(PhCH \longrightarrow CHPh)$  may be due to LiCl generated as a byproduct in the reaction of  $Cp_2ZrCl_2$  with *n*-BuLi. However, addition of LiCl (4 equiv) did not affect the rate of reaction.

<sup>(11)</sup> Compound 11 was prepared as a 9:1 mixture of the Z and  $\overline{E}$  isomers (11) Composed a Value property of the provide a sector of the provide the provide the provided of the provided of

<sup>(14)</sup> Takahashi, T.; Kasai, K.; Suzuki, N.; Nakajima, K.; Negishi, E. Organometallics, in press. See also: (a) Kaminsky, W.; Kopf, J.; Sinn, H.; Volimer, H.-J. Angew. Chem., Int. Ed. Engl. 1976, 15, 629. (b) Cotton, F. A.; Kibala, P. A. Inorg. Chem. 1990, 29, 3192.

<sup>(15)</sup> A search for bimetallic intermediates was urged by a referee, whom we thank.