

PII: S0040-4039(96)02028-X

Novel Palladium Catalyzed Dehydrogenation of α-Hydrazinonitriles to Hydrazonoyl Cyanides Using Cyclopentene. Synthesis of 1H-Pyrazole-4-Carboxylate

Yong Hae Kim^{*} and Jun Young Choi

Department of Chemistry, Korea Advanced Institute of Science and Technology, 373-1, Kusong Dong, Yusong Gu, Taejon 305-701, Korea

Abstract. Palladium charcoal catalyzed dehydrogenation of α -hydrazinonitriles has been found to give hydrazonoyl cyanides in the presence of cyclopentene in high yields, which was applied to the synthesis of 1*H*-pyrazole-4-carboxylates.Copyright © 1996 Elsevier Science Ltd

Hydrazones have been extensively utilized as important substrates in heterocyclic synthesis¹ and asymmetric synthesis.² Especially, hydrazonoyl halides, aldehyde hydrazones and α -nitrohydrazones have been used for the generation of nitrile imines, which are well used for 1,3-dipolar cycloadditions.³ But, there are few reports⁴ concerning general syntheses of hydrazonoyl cyanides for the utilization of heterocyclic synthesis.

In this paper, we report a novel palladium⁵ charcoal catalyzed dehydrogenation of α -hydrazinonitriles to hydrazonoyl cyanides in the presence of cyclopentene in high yields. There are many reports concerning stoichiometric⁶ or catalytic^{5a,7} oxidations of secondary amines to imines but no reports concerning stoichiometric or catalytic oxidation of any kind of hydrazines to hydrazones to our knowledges. During the research for N-N bond cleavage of a chiral α -hydrazinonitrile 1^{2b,8}, it was unexpectedly found that 1 was dehydrogenated in the presence of Pd/C and H₂ to afford a hydrazonoyl cyanide 2 instead of reductive cleavage of N-N bond.



To examine generality of the present reaction, various α -hydrazinonitriles 3 were tested as substrates (Table 1). α -Hydrazinonitriles were prepared from hydrocyanation of aldehyde hydrazones with HCl-NaCN in MeOH-H₂O at room temperature in 85-90% yields.⁹ α -N,N-Methylphenylhydrazinonitrile 3a reacted with equimolar amount of commercially available 10 % Pd/C in MeOH at room temperature for 3 h to give smoothly a hydrazonoyl cyanide 4a in 85 % yield. But, the stoichiometric use of Pd/C is undesiable from economic and environmental viewpoints. In the case of use of catalytic amount (10 mol %) of Pd/C, most of starting material remained even after 48 h stirring to afford 4a of trace amounts (below 10 %). However, when excess cyclopentene, which is good hydrogen acceptor, was added to the above reaction mixture,

Table 1. Palladium Catalyzed Oxidation of α-Hydrazinonitriles to Hydrazonoyl Cyanides Using Cyclopentene.



Run	Substrate	R ¹	R ²	Pd/C ^a (mol	%)Cyclopentene (eq.)	Time (h)	Product	Yield ^b (%)
1	3a	Me	(CH ₃) ₂ CH	100	-	3	4 a	85
2	3a	Me	(CH ₃) ₂ CH	10	-	48	4 a	trace
3	3a	Me	(CH ₃) ₂ CH	10	2	3	4a	82
4	3b	Me	CH3	100	-	3	4b	83
5	3b	Me	CH ₃	10	2	3	4b	80
6	3c	Me	PhCH ₂	100	-	3	4 c	82
7	3c	Me	PhCH ₂	10	2	3	4c	81
8	3d	Me	CH ₃ (CH ₂) ₃	10	2	3	4d	85
9	3e	н	(CH ₃) ₂ CHCH	2 10	2	10	4 e	75
10	3f	Н	CH ₃ (CH ₂) ₃	10	2	10	4f	71

a. When Pd(OH)₂ was used, trace amounts of product were obtained.

b. Isolated yield.

surprisingly starting material disappeared completely after 3 h stirring, while cyclohexene had no effect under the same reaction conditions. α -Substituted 3 (R²:Me (3b), PhCH₂ (3c) and n-Bu (3d)) resulted in high yields (80-85%) of 4. α -N-Phenylhydrazinonitriles (3e and 3f) were converted into 4 in 75% and 71% respectively without forming any azocompound. Hydrazonoyl cyanides exist as a mixture of *E* and *Z* isomers.¹⁰ Both isomers were observed by ¹H NMR spectroscopy¹¹ (4b,4e and 4f) and a single isomer was observed for others (4a,4c and 4d). General experimental procedure is as following:

To α -hydrazinonitrile (1mmol) in MeOH (2 ml) was added 0.1 mmol of commercially available 10 % Pd/C and 2 mmol of cyclopentene at room temperature and stirred for given hours (Table 1). And palladium charcoal was filtered through Celite and the filtrate was condensed under reduced pressure and purified with silicagel chromatography to give the corresponding hydrazonoyl cyanide.¹¹



Figure 1. Oxidative Dehydrogenation of α -Hydrazinonitriles.

The present palladium catalyzed dehydrogenation reactions may be rationalized by Figure 1.⁵ The reaction appears to be initiated by insertion of palladium into the N-H bond of an α -hydrazinonitrile. Reductive elimination of 3' gives product 4 and [H-Pd-H], which affords hydrogens to the cyclopentene to reproduce Pd for a catalyst. Similar ruthenium catalyzed activation of C-H bond of the α -hydrogen of a cyano group has been reported.¹²

 α -N-phenylhydrazonoyl cyanide (4e and 4f) turned out to be good substrates for the synthesis of heterocyclic compounds. As one of the applications, N-phenylhydrazonovl cyanides (4e and 4f) reacted with ethyl acrylate in the presence of NaH to give the corresponding ethyl 3-alkyl-1-phenyl-1H-pyrazole-4carboxylates (5a and 5b) respectively in high yields (5a: 84%, 5b: 85%)¹³, while formation of 1H-pyrazole-5carboxylates 7 were not detected. It was reported that 1,3-dipolar cycloaddition from nitrile imine and alkylpropiolate afforded 7 as major product.¹⁴ The cycloadditions may be rationalized by a stepwise path rather than a concerted path¹⁵ which is 1,3-dipolar cycloaddition mechanism of nitrile imines. The reaction appears to involve the conjugate addition of sodium salt of 4e or 4f by NaH to ethyl acrylate following by the intramolecular cyclization of the initially formed enolate, accompanying leaving of CN group, and finally following aromatization of pyrazoline to afford thermodynamic stable pyrazoles.^{16a} An intermediate 6 of the stepwise reaction was actually isolated in 75% yield and confirmed when the reaction was carried out under mild conditions (reaction temperature: -78 °C-0 °C). The structure of 5 could be characterized by comparison of the chemical shifts (5a: 8.33 ppm, 5b: 8.32 ppm) of proton at the 5 position with those from the similar compounds in ¹H NMR spectrum.¹⁶



In conclusion, the present reaction provides novel general method for the synthesis of hydrazonoyl cyanide through Pd catalized dehydrogenation reaction of α -hydrazinonitriles in the presence of cyclopentene and thus hydrazonoyl cyanides could be used as the good substrates for the synthesis of 1*H*-pyrazole-4-carboxylate. Further studies on synthetic application of the present method are in progress.

Acknowledgement : This work was supported by CBM from Korea Science and Engineering Foundation.

References and Notes

- 1. Review : Hafez, E. A. A; Abed, N. M.; Elmoghayer, M. R. H.; El-Agamey, A. G. Heterocycles, 1984, 22, 1821-1877.
- (a) Review : Enders, D., Asymmetric Synthesis, Ed. J. D. Morrison, Academic press, New York, Vol3, 1984.
 (b) Kim, Y. H.; Choi, J. Y. Tetrahedron Lett., 1996, 37, 5543-5546.
- Review: (a) Huisgen, R. Angew. Chem. Int. Ed. Engl., 1963, 2, 565-632. (b) Caramella, P.; Grunanger, P 1,3-Dipolar Cycloaddition Chemisry, Ed by Padwa, A., John Wiley and Sons, New York, Vol. 1-2, 1984.
- 4. (a) Shawali, A. S.; Hassaneen, H. M.; Sherif, S. M. J. Heterocyclic Chem., **1980**, 17, 1745-1749. (b) Ranganathan, D.; Barnezai, S. Tetrahedron Lett. **1983**, 24, 1067-1070.
- 5. Examples of Pd catalyzed dehydrogenations : (a) Murahashi, S. I.; Yoshimura, N.: Tsumiyama, T.;

Kojima, T. J. Am. Chem. Soc., 1983, 105, 5002-5011. (b)Murahashi, S. I.; Mitsui, H.; Watanabe, T.; Zenki, S. Tetrahedron Lett., 1983, 24, 1049-1052. (c) Amemiya, Y.; Miller, D. D.; Hsu, F. L. Synth. Comm., 1990, 20, 2483-2489.

- (a) Pratt, E. F.; Mcgovern, T. P. J. Org. Chem., 1964, 29, 1540-1543. (b) Bonnett, R.; Clark, V. M.; Giddey, A.; Todd, A. J. Chem. Soc. 1959, 2087-2093. (c) Grundon, M. F.; Reyhold, B. E. ibid, 1964, 2445-2450. (d) Fryer, R. I.; Archer, G. A.; Brust, B.; Zally, W.; Stembach, L. H. J. Org. Chem., 1965, 30, 1308-1310.(e) Muller, P.; Gilabert, P. A. Tetrahedron, 1988, 44, 7171-7175. (f) Porta, F.; Crotti, C.; Cenini, S. J. Mol. Catal., 1989, 50, 333-341. (g) Wehrli, P. A.; Schaer, B. Synthesis, 1974, 288-289.
- (a) Murahashi, S.; Naota, T.; Taki, H. J. Chem. Soc., Chem. Commun., 1985, 613-614. (b) Nishinaga, A., Yamazaki, S.; Matsuura, T. Tetrahedron Lett., 1988, 29, 4115-4118. (c) Maruyama, K.; Kusukawa, T.; Higuchi, Y.; Nishinaga, A. Chem. Lett., 1991, 1093-1096. (d) Yamazaki, S. Chem. Lett., 1992, 823-826.
 (e) Yamaguchi, T.; Takeda, T. Chem. Lett., 1992, 1933-1936.
- 8. The cleavage of N-N bond of hydrazines α -substituted with alkyl or aryl groups readily occurs by Pd/C system. However, indoline hydrazine 1 containing CN group was oxidized to 2 in 80% yield.
- 9. Hydrocyanation reaction of benzaldehyde hydrazone did not occur even under reflux condition.
- 10. (a) Karabatsos, G. J.; Taller, R. A. *Tetrahedron*, **1968**, 24, 3923-3927. (b) Karabatsos, G. J.; Krumel, K. L. *ibid*, **1967**, 23, 1097-1105.
- 4a : yellow oil; ¹H NMR (CDCl₃, 200 mHz) δ 1.24 (6H, dd), 2.83 (1H, m), 3.80 (6H, s), 7.02 (1H, m), 7.28 (4H, m).; IR (NaCl) 2191 (CN) cm⁻¹; MS (EI) m/z 201.

4b : mp 49-51 °C; ¹H NMR (CDCl₃, 200 mHz) δ 1.96 and 2.29 (3H, 2s, E/Z), 3.47 and 3.81 (3H, 2s, E/Z), 7.04 (1H, m), 7.29 (4H, m).; IR (NaCl) 2192 (CN) cm⁻¹; MS (EI) m/z 173.

4c : mp 70-72; ¹H NMR (CDCl₃, 200 mHz) δ 3.82 (3H, s), 3.83 (2H, s), 7.09 (1H, m), 7.30 (4H, m).; IR (NaCl) 2190 (CN) cm⁻¹; MS (EI) m/z 249.

4d : yellow oil; ¹H NMR (CDCl₃, 200 mHz) δ 0.958 (3H, t), 1.41 (1H, m), 1.65 (1H, m),2.56 (2H, t), 3.81 (3H, s), 7.04 (1H, m), 7.29 (4H, m).; IR (NaCl) 2191 (CN) cm⁻¹; MS (EI) m/z 215.

4e : mp 48-50 °C; ¹H NMR (CDCl₃, 200 mHz) δ 0.992 and 1.07 (3H, 2d, E/Z), 2.02 (1H, m), 2.25 and 2.36 (2H, 2d, E/Z), 6.97-7.34 (5H, m), 8.06 and 8.47 (1H, 2s, NH E/Z).; IR (NaCl) 2199 (CN) cm⁻¹; MS (EI) m/z 201.

4f : mp 60-62 °C; ¹H NMR (CDCl₃, 200 mHz) δ 0.994 and 0.978 (3H, 2t, E/Z), 1.44 (1H, m), 1.65 (1H, m), 2.32 and 2.51 (2H, 2t, E/Z), 6.99-7.34 (5H, m), 7.98 and 8.55 (1H, 2s, NH E/Z).; IR (NaCl) 2199 (CN) cm⁻¹; MS (EI) m/z 201.

- 12. Naota, T.; Taki, H.; Mizuno, M; Murahashi, S. I. J. Am. Chem. Soc., 1989, 111, 5954-5955.
- 13. General experimental procedure is as following :

To a solution of N-phenylhydrazonoyl cyanide 3e or 3f (1 mmol) and ethyl acrylate (1.2 mmol) in THF (2ml) was NaH (3 eq) in one portion at room temperature. After 30 min, H₂O was added, and reaction mixture was extracted with CH₂Cl₂, dried over MgSO₄ and evaporated under reduced pressure. The residue was purified by silica column chromatography to give the corresponding ethyl 3-alkyl-1-phenyl-1*H*-pyrazole-4-carboxylates.

5a : yellow oil; ¹H NMR (CDCl₃, 200 mHz) δ 0.957 (6H, d), 1.35 (3H, t), 2.09 (1H, m), 2.82 (2H, d), 4.28 (2H, q), 7.25-7.69 (5H, m), 8.33 (1H, s).; IR (NaCl) 1713, 1600, 1547 cm⁻¹; MS (EI) m/z 272. **5b** : yellow oil; ¹H NMR (CDCl₃, 200 mHz) δ 0.907 (6H, d), 1.49 (2H, m), 1.69 (2H, m), 2.99 (2H, m),

4.30 (2H, q), 7.29-7.69 (5H, m), 8.32 (1H, s).; IR (NaCl) 1716, 1599, 1548 cm⁻¹; MS (EI) m/z 272.

- 14. Huisgen, R.; Seidel, M.; Wallbillich, G.; Knupfer, H. Tetrahedron ,1962, 17, 3-29.
- 15. Huisgen, R. J. Org. Chem. 1976, 41, 403-419.
- (a) Oida, T., Shimizu, T.; Hayashi, Y.; Teramura, K. Bull. Chem. Soc. Jpn. 1981, 54, 1429-1433. (b) Ashton, W. T.; Doss, G. A. J. Heterocyclic Chem. 1993, 30, 307-331.

(Received in Japan 21 August 1996; revised 12 October 1996; accepted 14 October 1996)